

Global School-based Student Health Survey



Guatemala 2015 Fact Sheet

The 2015 Guatemala GSHS was a school-based survey of students in Primero, Segundo, and Tercero basico, which are typically attended by students aged 13-17. A two-stage cluster sample design was used to produce data representative of all students in Primero, Segundo, and Tercero basico in Guatemala. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate.

The Guatemala GSHS measured alcohol use; dietary behaviors; drug use; hygiene; mental health; physical activity; sexual behaviors; and violence and unintentional injury. Students self-reported their responses to each question on a computer scanable answer sheet.

The school response rate was 96%, the student response rate was 85%, and the overall response rate was 82%. A total of 4,374 students participated in the Guatemala GSHS. Prevalence estimates (percentages) and 95% confidence intervals are presented below.

	Students Aged 13-15 Years			Students Aged 16-17 Years			Students Aged 13-17 Years		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Alcohol Use									
Percentage of students who currently drank alcohol (at least one drink of alcohol on at least one day during the 30 days before the survey)	16.8 (13.5-20.7)	18.3 (13.3-24.7)	15.1 (11.6-19.5)	25.4 (17.9-34.8)	27.4 (18.6-38.5)	21.6 (12.9-33.9)	18.3 (14.9-22.4)	20.1 (14.6-27.1)	16.1 (12.9-19.9)
Percentage of students who ever drank so much alcohol that they were really drunk one or more times during their life	11.7 (8.7-15.6)	11.9 (8.2-17.0)	11.2 (7.7-16.0)	19.8 (13.8-27.5)	22.3 (13.8-33.8)	14.9 (8.8-24.1)	13.1 (10.4-16.4)	14.0 (10.1-19.2)	11.7 (8.5-15.9)
Dietary Behaviors									
Percentage of students who were underweight (<-2SD from median for BMI by age and sex)	1.2 (0.5-2.9)	2.1 (0.8-5.2)	0.3 (0.1-0.9)	4.6 (1.5-13.6)	7.7 (2.1-24.4)	0.1 (0.0-0.3)	1.8 (0.9-3.6)	3.2 (1.5-6.5)	0.2 (0.1-0.8)
Percentage of students who were overweight (>+1SD from median for BMI by age and sex)	28.0 (21.5-35.7)	29.9 (22.2-38.9)	26.0 (19.0-34.5)	32.4 (23.2-43.3)	23.3 (15.1-34.2)	45.9 (25.8-67.5)	28.8 (22.1-36.6)	28.6 (21.6-36.7)	29.2 (21.1-38.7)

Global School-based Student Health Survey

Guatemala 2015 Fact Sheet



	Students Aged 13-15 Years			Students Aged 16-17 Years			Students Aged 13-17 Years		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Dietary Behaviors									
Percentage of students who were obese (>+2SD from median for BMI by age and sex)	7.7 (5.4-10.8)	8.4 (5.9-11.8)	7.0 (4.2-11.3)	9.5 (6.0-14.7)	6.3 (3.0-12.9)	14.1 (9.1-21.3)	8.0 (5.8-11.0)	7.9 (5.5-11.3)	8.1 (5.7-11.4)
Percentage of students who usually drank carbonated soft drinks one or more times per day during the 30 days before the survey	61.4 (55.5-67.0)	63.5 (57.1-69.5)	59.7 (52.9-66.2)	55.6 (48.6-62.4)	59.3 (50.4-67.7)	52.3 (40.5-64.0)	60.4 (55.2-65.3)	62.7 (57.0-68.1)	58.6 (52.2-64.8)
Drug Use									
Percentage of students who ever used marijuana one or more times during their life	5.1 (3.4-7.6)	6.7 (4.1-10.8)	2.9 (2.0-4.2)	8.6 (4.1-17.0)	13.1 (6.4-24.9)	2.4 (0.6-8.9)	5.7 (4.2-7.8)	8.0 (5.3-11.7)	2.8 (2.0-4.0)
Percentage of students who used drugs before age 14 years for the first time, among students who ever used drugs	77.7 (58.6-89.6)	81.7 (64.1-91.7)	67.4 (44.0-84.5)	43.1 (17.6-72.9)	*	*	70.8 (55.6-82.4)	71.6 (57.6-82.4)	65.8 (45.8-81.4)
Hygiene									
Percentage of students who usually cleaned or brushed their teeth one or more times per day during the 30 days before the survey	94.4 (91.5-96.3)	93.2 (87.4-96.5)	95.5 (93.3-97.1)	93.9 (90.6-96.1)	98.2 (93.3-99.5)	89.0 (82.2-93.4)	94.3 (91.7-96.1)	94.2 (90.4-96.6)	94.6 (91.8-96.5)
Percentage of students who never or rarely washed their hands after using the toilet or latrine during the 30 days before the survey	2.6 (1.3-5.1)	2.7 (1.3-5.7)	2.1 (1.1-4.1)	4.4 (2.2-8.6)	2.6 (0.9-7.3)	5.6 (3.0-10.1)	2.9 (1.8-4.6)	2.7 (1.6-4.5)	2.6 (1.7-4.0)

*Indicates data were not available.

Global School-based Student Health Survey



Guatemala 2015 Fact Sheet

	Students Aged 13-15 Years			Students Aged 16-17 Years			Students Aged 13-17 Years		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Mental Health									
Percentage of students who seriously considered attempting suicide during the 12 months before the survey	16.4 (13.4-20.0)	11.5 (6.6-19.3)	21.4 (18.5-24.6)	17.0 (11.2-24.9)	15.4 (8.8-25.8)	20.1 (8.5-40.6)	16.5 (13.4-20.2)	12.3 (8.1-18.3)	21.2 (18.0-24.8)
Percentage of students who attempted suicide one or more times during the 12 months before the survey	16.3 (12.9-20.3)	12.6 (8.7-17.9)	19.9 (14.8-26.2)	12.4 (7.2-20.7)	6.8 (2.5-17.5)	17.1 (6.3-38.7)	15.6 (12.6-19.1)	11.5 (8.2-15.8)	19.5 (15.5-24.1)
Percentage of students who did not have any close friends	6.4 (5.4-7.6)	6.7 (4.9-9.2)	6.1 (4.4-8.5)	7.2 (3.9-12.7)	10.9 (5.9-19.5)	1.7 (0.4-7.7)	6.5 (5.3-8.1)	7.6 (5.4-10.5)	5.5 (3.8-7.8)
Physical Activity									
Percentage of students were physically active at least 60 minutes per day on all 7 days during the 7 days before the survey	10.9 (7.8-15.1)	12.7 (8.0-19.6)	8.7 (6.2-12.2)	8.6 (4.2-16.8)	10.2 (4.2-22.5)	6.5 (3.8-10.9)	10.5 (7.5-14.6)	12.2 (7.8-18.7)	8.4 (5.9-11.8)
Percentage of students who attended physical education classes on three or more days each week during this school year	35.9 (28.3-44.4)	36.5 (28.6-45.1)	34.7 (26.1-44.4)	43.9 (29.5-59.5)	50.6 (35.0-66.2)	34.3 (17.5-56.3)	37.3 (29.1-46.3)	39.3 (30.9-48.4)	34.6 (26.0-44.3)
Percentage of students who spent three or more hours per day sitting and watching television, playing computer games, or talking with friends, when not in school or doing homework during a typical or usual day	23.1 (18.6-28.4)	20.9 (16.0-27.0)	25.6 (20.0-32.1)	19.2 (13.0-27.6)	19.8 (13.9-27.4)	18.7 (8.1-37.3)	22.5 (17.8-28.0)	20.7 (15.9-26.5)	24.7 (18.3-32.3)

Global School-based Student Health Survey



Guatemala 2015 Fact Sheet

	Students Aged 13-15 Years			Students Aged 16-17 Years			Students Aged 13-17 Years		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Sexual Behaviors									
Percentage of students who ever had sexual intercourse	14.5 (11.4-18.3)	21.2 (17.6-25.3)	7.6 (4.8-11.9)	40.2 (31.4-49.7)	57.0 (42.5-70.4)	16.3 (10.2-25.0)	19.0 (14.6-24.4)	28.5 (22.2-35.7)	8.9 (6.3-12.3)
Percentage of students who had sexual intercourse before age 14 years for the first time, among students who ever had sexual intercourse	63.2 (51.4-73.6)	61.4 (46.4-74.5)	*	22.4 (12.9-35.9)	24.6 (14.3-38.8)	*	48.1 (36.5-60.0)	47.0 (34.7-59.7)	49.8 (24.6-75.2)
Percentage of students who used a condom during last sexual intercourse, among students who ever had sexual intercourse	58.9 (46.7-70.1)	59.0 (48.5-68.8)	*	58.1 (46.7-68.6)	59.7 (45.9-72.2)	*	58.6 (50.1-66.6)	59.3 (50.8-67.2)	55.4 (36.5-72.9)
Violence and Unintentional Injury									
Percentage of students who were in a physical fight one or more times during the 12 months before the survey	22.8 (19.8-26.1)	31.2 (26.8-35.9)	14.1 (10.8-18.2)	25.1 (17.9-33.9)	32.8 (21.9-45.9)	12.7 (7.5-20.6)	23.2 (20.3-26.3)	31.5 (27.1-36.2)	13.9 (11.0-17.4)
Percentage of students who were seriously injured one or more times during the 12 months before the survey	32.7 (27.2-38.8)	38.3 (31.6-45.5)	27.0 (22.2-32.3)	30.6 (24.2-37.9)	34.5 (25.6-44.6)	21.7 (15.1-30.2)	32.3 (27.9-37.1)	37.6 (32.1-43.3)	26.2 (22.2-30.7)
Percentage of students who were bullied on one or more days during the 30 days before the survey	22.8 (19.6-26.3)	26.0 (21.2-31.4)	19.6 (16.4-23.2)	27.4 (22.3-33.1)	30.2 (18.7-44.8)	22.2 (12.1-37.1)	23.6 (20.3-27.2)	26.8 (21.2-33.3)	20.0 (17.5-22.7)

For additional information, please contact:
Olivia J. Brathwaite Dick,
OMS Guatemala, Guatemala, brathwaiteo@paho.org

For general information about the GSHS, please visit:
<http://www.who.int/chp/gshs/en/>
or
<http://www.cdc.gov/gshs>

*Indicates data were not available.