

GYTS Objectives

The Global Youth Tobacco Survey (GYTS), a component of the Global Tobacco Surveillance System (GTSS), is a global standard for systematically monitoring youth tobacco use (smoking and smokeless) and tracking key tobacco control indicators.

GYTS is a cross-sectional, nationally representative school-based survey of students in grades associated with ages 13 to 15 years. GYTS uses a standard core questionnaire, sample design, and data collection protocol. It assists countries in fulfilling their obligations under the World Health Organization (WHO) Framework Convention on Tobacco Control (FCTC) to generate comparable data within and across countries. WHO has developed MPOWER, a technical package of selected demand reduction measures contained in the WHO FCTC:



- Monitor tobacco use & prevention policies
- Protect people from tobacco smoke
- Offer help to quit tobacco use
- Warn about the dangers of tobacco
- Enforce bans on tobacco advertising, promotion, & sponsorship
- Raise taxes on tobacco

GYTS Methodology

GYTS uses a global standardized methodology that includes a two-stage sample design with schools selected with a probability proportional to enrollment size. The classes within selected schools are chosen randomly and all students in selected classes are eligible to participate in the survey. The survey uses a standard core questionnaire with a set of optional questions that countries can adapt to measure and track key tobacco control indicators. The questionnaire covers the following topics: tobacco use (smoking and smokeless), cessation, secondhand smoke (SHS), pro- and anti-tobacco media and advertising, access to and availability of tobacco products, and knowledge and attitudes regarding tobacco use. The questionnaire is self-administered; using scannable paper-based bubble sheets, it is anonymous to ensure confidentiality.

In the Kyrgyz Republic, GYTS was conducted in 2019 by the Department of Disease Prevention, and State Sanitary and Epidemiological Surveillance of the Ministry of Health of the Kyrgyz Republic and the Ministry of Education and Science of the Kyrgyz Republic. The overall response rate was 88.8%. A total of 6,145 eligible students in grades 7 - 9 completed the survey, of which 5,447 were aged 13-15 years. Data are reported for students aged 13-15 years.



The Ministry of Health
of the Kyrgyz Republic



World Health
Organization



The mark "CDC" is owned by the US Dept. of Health and Human Services and is used with permission. Use of this logo is not an endorsement by HHS or CDC of any particular product, service, or enterprise.

GYTS Highlights

TOBACCO USE

- 6.0% of students, 9.5% of boys, and 2.4% of girls currently used any tobacco products.
- 4.4% of students, 6.8% of boys, and 2.0% of girls currently smoked tobacco.
- 2.4% of students, 4.2% of boys, and 0.6% of girls currently smoked cigarettes.
- 1.8% of students, 2.4% of boys, and 1.2% of girls currently smoked hookah.
- 2.4% of students, 4.3% of boys, and 0.6% of girls currently used smokeless tobacco.

ELECTRONIC CIGARETTES

- 2.8% of students, 3.9% of boys, and 1.7% of girls currently used electronic cigarettes.

CESSATION

- Almost 6 in 10 (56.0%) students who currently smoke tobacco tried to stop smoking in the past 12 months.
- More than 7 in 10 (73.4%) students who currently smoke tobacco wanted to stop smoking now.

SECONDHAND SMOKE

- 16.8% of students were exposed to tobacco smoke at home.
- 14.2% of students were exposed to tobacco smoke inside enclosed public places.

ACCESS & AVAILABILITY

- 87.2% of students who currently smoke cigarettes bought cigarettes from a store, shop, street vendor, kiosk, or in the market (bazar).
- Among students who currently smoke cigarettes who tried to buy cigarettes, 58.5% were not prevented from buying them because of their age.

MEDIA


- Almost 5 in 10 (49.7%) students noticed anti-tobacco messages in the media.
- Almost 3 in 10 (26.7%) students noticed tobacco advertisements or promotions when visiting points of sale.
- 4.9% of students had something with a tobacco brand logo on it.

KNOWLEDGE & ATTITUDES

- 71.2% of students definitely thought other people's tobacco smoking is harmful to them.
- 77.7% of students favored prohibiting smoking inside enclosed public places.

TOBACCO USE

SMOKED TOBACCO	OVERALL (%)	BOYS (%)	GIRLS (%)
Current tobacco smokers ¹	4.4	6.8	2.0*
Current cigarette smokers ²	2.4	4.2	0.6*
Frequent cigarette smokers ³	0.2	0.3	0.0
Current smokers of other tobacco ⁴	2.8	4.1	1.6*
Current hookah smokers ⁵	1.8	2.4	1.2*
Ever tobacco smokers ⁶	17.2	25.5	8.9*
Ever cigarette smokers ⁷	13.2	20.9	5.7*
Ever smokers of other tobacco ⁸	9.6	13.4	5.8*
Ever hookah smokers ⁹	9.3	13.2	5.4*
SMOKELESS TOBACCO			
Current smokeless tobacco users ¹⁰	2.4	4.3	0.6*
Ever smokeless tobacco users ¹¹	5.1	8.2	2.0*
ANY TOBACCO USE (smoked and/or smokeless)			
Current tobacco users ¹²	6.0	9.5	2.4*
Ever tobacco users ¹³	18.7	27.4	10.1*
SUSCEPTIBILITY			
Never tobacco users susceptible to tobacco use in the future ¹⁴	11.5	13.5	9.8*
Never tobacco smokers who thought they might enjoy smoking a cigarette ¹⁵	12.0	17.1	7.9*



ELECTRONIC CIGARETTES

	OVERALL (%)	BOYS (%)	GIRLS (%)
Current electronic cigarette users ¹⁶	2.8	3.9	1.7*
Ever electronic cigarette users ¹⁷	5.1	7.4	2.8*

CESSATION

	OVERALL (%)	BOYS (%)	GIRLS (%)
Current tobacco smokers who tried to stop smoking in the past 12 months	56.0	58.2	--
Current tobacco smokers who wanted to stop smoking now	73.4	80.0	--
Current tobacco smokers who thought they would be able to stop smoking if they wanted to	89.2	90.2	--
Current tobacco smokers who have ever received help/advice from a program or professional to stop smoking	29.4	27.4	--

SECONDHAND SMOKE

	OVERALL (%)	BOYS (%)	GIRLS (%)
Exposure to tobacco smoke at home ¹⁸	16.8	19.9	13.6*
Exposure to tobacco smoke inside any enclosed public place ¹⁸	14.2	14.0	14.4
Exposure to tobacco smoke at any outdoor public place ¹⁸	14.8	15.4	14.2
Students who saw anyone smoking inside the school building or outside on school property ¹⁹	32.9	33.1	32.6

ACCESS & AVAILABILITY

	OVERALL (%)	BOYS (%)	GIRLS (%)
Current cigarette smokers who bought cigarettes from a store, shop, street vendor, kiosk, or in the market (bazar) ²⁰	87.2	88.9	--
Current cigarette smokers who were not prevented from buying cigarettes because of their age ²¹	58.5	60.0	--
Current cigarette smokers who bought cigarettes as individual sticks ²²	44.0	45.5	--

MEDIA

TOBACCO ADVERTISING	OVERALL (%)	BOYS (%)	GIRLS (%)
Students who noticed tobacco advertisements or promotions at points of sale ²³	26.7	29.6	23.9*
Students who saw anyone using tobacco on television, videos, or movies ²⁴	67.6	68.0	67.1
Students who were ever offered a free tobacco product from a tobacco company representative	4.5	5.8	3.2*
Students who had something with a tobacco brand logo on it	4.9	6.4	3.4*
ANTI-TOBACCO ADVERTISING			
Students who noticed anti-tobacco messages in the media ¹⁷	49.7	50.3	49.1
Students who noticed anti-tobacco messages at sporting or community events ²⁵	43.7	44.6	42.5
Current tobacco smokers who thought about quitting because of a warning label ²⁶	34.4	35.2	31.4
Students who were taught in school about the dangers of tobacco use in the past 12 months	70.1	67.7	72.6*

KNOWLEDGE & ATTITUDES

	OVERALL (%)	BOYS (%)	GIRLS (%)
Students who definitely thought it is difficult to quit once someone starts smoking tobacco	30.1	28.9	31.4
Students who thought smoking tobacco helps people feel more comfortable at celebrations, parties, and social gatherings	29.0	31.5	26.3*
Students who definitely thought other people's tobacco smoking is harmful to them	71.2	67.7	74.6*
Students who favored prohibiting smoking inside enclosed public places	77.7	74.9	80.5*
Students who favored prohibiting smoking at outdoor public places	74.5	71.5	77.6*

¹ Smoked cigarettes or other type of tobacco anytime during the past 30 days. ² Smoked cigarettes anytime during the past 30 days. ³ Smoked cigarettes on 20 or more days of the past 30 days. ⁴ Smoked tobacco other than cigarettes anytime during the past 30 days. ⁵ Smoked hookah anytime during the past 30 days. ⁶ Ever smoked cigarettes or other type of tobacco, even one or two puffs. ⁷ Ever smoked cigarettes, even one or two puffs. ⁸ Ever smoked tobacco other than cigarettes, even one or two puffs. ⁹ Ever smoked hookah, even one or two puffs. ¹⁰ Used smokeless tobacco anytime during the past 30 days. ¹¹ Ever used smokeless tobacco. ¹² Smoked cigarettes, smoked other type of tobacco, and/or used smokeless tobacco anytime during the past 30 days. ¹³ Ever smoked tobacco and/or used smokeless tobacco. ¹⁴ Susceptible to future tobacco use includes those who answered "definitely yes", "probably yes", or "probably not" to using tobacco if one of their best friends offered it to them, or "definitely yes", "probably yes", or "probably not" to using tobacco during the next 12 months. ¹⁵ Those who answered "Agree" or "Strongly Agree" to the statement: "I think I might enjoy smoking a cigarette". ¹⁶ Used electronic cigarettes anytime during the past 30 days. Current use of electronic cigarettes is assessed separately from cigarettes, other smoked tobacco products, and smokeless tobacco products and is not included in the current definition of current any tobacco use. ¹⁷ Ever used electronic cigarettes in their entire life. ¹⁸ During the past 7 days. ¹⁹ During the past 30 days. ²⁰ Outlet from which current cigarette smokers bought cigarettes the last time they smoked cigarettes in the past 30 days. ²¹ Among those who tried to buy cigarettes during the past 30 days. ²² Based on the last purchase, among those who bought cigarettes during the past 30 days. ²³ Among those who visited a point of sale in the past 30 days. ²⁴ Among those who watched television, videos, or movies in the past 30 days. ²⁵ Among those who attended sporting or community events in the past 30 days. ²⁶ Among those who noticed warning labels on cigarette packages in the past 30 days.

NOTE: Data are weighted to be nationally representative of all students aged 13-15 years who are enrolled in school. Percentages reflect the prevalence of each indicator in each group, not the distribution across groups. --Estimates based on unweighted cases less than 35 are not presented. *Gender comparisons are significant at p < 0.05.