


1993 CZECH REPUBLIC REPRODUCTIVE HEALTH SURVEY

Final Report



Czech Statistical Office

Factum, non Fabula

**WHO Collaborating Center for Perinatal Medicine/
Institute for the Care of Mother and Child, Prague**

Centers for Disease Control and Prevention, USA

March 1995

1993 CZECH REPUBLIC
REPRODUCTIVE HEALTH SURVEY
Final Report

Czech Statistical Office

Factum, non Fabula

**WHO Collaborating Center for Perinatal Medicine/
Institute for the Care of Mother and Child**

Centers for Disease Control and Prevention, USA

March, 1995

ACKNOWLEDGEMENTS

This report describes the 1993 Czech Republic Reproductive Health Survey (CRRHS) and its major findings. This nationwide undertaking could not have taken place without the involvement and cooperation of a large number of individuals and organizations in the Czech Republic and the United States. Funding for the survey was provided by the United States Agency for International Development (USAID/Washington). The survey was directed by Dr. Ivan Tomek and Mr. Jaroslav Kraus, who were responsible for coordinating many of the survey activities. Professor Zdenek Stembera, director of the World Health Organization Collaborating Center for Perinatal Medicine was the principal resource person for maternal and child health and administered the funds for the survey. Dr. Jitka Slavikova served as fieldwork coordinator. Drs. Howard Goldberg and Petr Velebil of the United States Centers for Disease Control and Prevention (CDC) provided technical assistance and consultation for all aspects of the survey. Mr. James Bednar and other staff members at USAID/Prague provided support to the survey in a number of ways. Besides these individuals and their organizations, there were many others that contributed to the many tasks necessary for successful completion of the survey, including administrative tasks, questionnaire development and review, training, field work/data collection, data processing, and analysis.

This report was prepared by Howard Goldberg, Petr Velebil, Zdenek Stembera, Ivan Tomek, and Jaroslav Kraus.

Contributors to the Survey

The following organizations and individuals made important contributions to the success of the 1993 Czech Republic Health Survey:

Dr. I. Tomek, Survey co-director Mr. J.
Kraus, Survey co-director Mrs. J.
Slavikova, Fieldwork coordinator

Czech Statistical Office

Mr. Edvard Outrata. President
Mr. J. Fischer, CSc, Vice president
Mrs. J. Ruzkova, CSc, Director, Vital Statistics Department
Mr. J. Kraus, Vital Statistics Department
Mr. J. Pecka, Programming Department
Mrs. L. Zacharova, Technical Support Department
Dr. B. Jungmann, CSc, Director, Public Opinion Research Institute
(PORI)
Dr. M. Vasickova, Interviewer supervisor, PORI
Mrs. E. Sochorova, Interviewer supervisor, PORI
Dr. J. Michalec, Survey Department, PORI
Ms. V. Kobylkova, Survey Department, PORI
J. Namestkova, A. Petraskova, M. Tyborova, PORI
Data entry staff
Interviewing staff
Printing office

FACTUM, *non fabula*

Dr. I. Tomek, CSc.
Dr. J. Slavikova Dr.
J. Herzmann, CSc.
Interviewing staff

WHO Collaborating Center for Perinatal Medicine/Institute for the Care of Mother and Child

Prof. MUDr. Z. Stembera, DrSc, Director, WHO Collaborating Center

Czech Family Planning Association

Dr. R. Uzel, CSc, Director, Czech Family Planning Association
Dr. V. Wynnyczuk, CSc, Czech Family Planning Association

Ministry of Health

Dr. M. Kosulicova, Chief, Department of Maternal and Child Health
Dr. K. Ciharova, Chief, Department of International Cooperation
Dr. I. Karaskova, Department of Maternal and Child Health
Dr. V. Albrecht, CSc, Chief, WHO Collaborating Centers Activity

Konsorcium Universitas

Dr. P. Kuchar, CSc.
Dr. J. Burianek, CSc.
Interviewers

Centers for Disease Control and Prevention. Division of Reproductive Health

Dr. H. Goldberg, PhD., Deputy Chief, Behavioral Epidemiology and
Demographic Research Branch
Dr. P. Velebil, MD., Visiting Scientist
Dr. L. Morris, PhD., Chief, Behavioral Epidemiology and Demographic
Research Branch
Ms. M. Watson, Chief, Computer Resources Section

United States Agency for International Development/Prague

Ms. L. Roussel, Country Representative
Mr. J. Bednar, Project Manager
Ms. E. Kostelkova, PhD., Project Specialist
Ms. S. Strbanova, PhD., Project Specialist

TABLE OF CONTENTS

	PAGE
ACKNOWLEDGEMENTS.....	iii
CONTRIBUTORS TO THE SURVEY	v
LIST OF TABLES.....	xi
LIST OF FIGURES.....	xxi
SUMMARY	1
CHAPTER I. INTRODUCTION/SURVEY METHODOLOGY.....	12
Survey Background	
Organizational Structure	
Questionnaire Content	
Sample Selection	
Data Collection	
Data Entry/Editing	
Response Rates	
CHAPTER II. RESPONDENT CHARACTERISTICS	20
CHAPTER III. CHILDBEARING/ABORTION.....	35
Children Ever Born	
Current Fertility	
Timing of First Births	
Desired Number of Children	
Planning Status of Pregnancies	
Induced Abortions	
CHAPTER IV. FAMILY PLANNING	59
Knowledge of Methods for Preventing Pregnancy	
Ever Use of Contraception	
Current Use of Contraception	
Source of Contraception	
Reasons for Not Using Contraception	
Contraceptive Failure/Discontinuation	

	PAGE
Use of Less Effective Methods	
Need for Family Planning Services	
Problems with Current Contraceptive Methods	
Contraceptive Sterilization	
Use of Infertility Services	
CHAPTER V. PREGNANCY, DELIVERY, AND INFANT HEALTH	97
Reaction to Pregnancy	
Physical Exertion/Stress during Pregnancy	
Prenatal Care	
Ultrasound Examinations	
Bed Rest/Hospitalization during Pregnancy	
Smoking during Pregnancy	
Birthweight/Pregnancy Outcomes	
Labor and Delivery	
Problems in the Post Partum Period	
Prenatal Counselling	
Breastfeeding	
CHAPTER VI. YOUNG ADULT SEXUAL BEHAVIOR.....	148
Initiation of Sexual Intercourse	
Contraception at First Intercourse	
Premarital Pregnancy Family Planning	
Discussions with Parents	
CHAPTER VII. REPRODUCTIVE HEALTH KNOWLEDGE AND ATTITUDES	161
Attitudes about Restrictions on Abortion	
Knowledge of the Menstrual Cycle	
Opinions about Oral Contraceptives	
Women's Family and Reproductive Roles	
CHAPTER VIII. WOMEN'S HEALTH	175
Gynecologic Examinations	
Breast Self Examination	
Cigarette Smoking	
Alcohol Consumption	
Overweight	

	PAGE
CHAPTER IX. TRANSMISSION OF HIV/AIDS.....	186
REFERENCES	190
APPENDIX A. SAMPLING ERRORS	192
APPENDIX B. CRRHS QUESTIONNAIRE	195

LIST OF TABLES

CHAPTER I: INTRODUCTION/METHODOLOGY

- I.1 Final Interview Status, by Size of Place
- I.2 Region of Residence and Size of Place of Residence, 1993 Czech Republic Reproductive Health Survey Respondents and Women 15-44 Years Old in the 1991 Czech Republic Census

CHAPTER II: RESPONDENT CHARACTERISTICS

- II.1 Comparison of Percent Distributions of Age and Marital Status according to the 1991 Czech Census and the 1993 Czech Republic Reproductive Health Survey
- II.2 Selected Sociodemographic Characteristics of Respondents, by Age Group
- II.3 Percent Distributions of Monthly Household Income, Employment Status of Respondents and Average Number of Hours Worked per Week
- II.4 Percent of Respondent Households with Various Possessions/Amenities and Percent Distribution of Number of Selected Possessions in Households
- II.5 Ownership of Residence, Number of Families Living in Residence and Number of People Living in Residence
- II.6 Husband's Education, Household Income, and Number of Selected Possessions, by Respondent's Education, Women 20 to 44 Years of Age
- II.7 Life Table Estimates of Percent of Women Ever in Union Before Ages 18, 20, and 25 and Median Age at First Union, by Selected Characteristics
- II.8 Percent of Women Ever in Union Who Are Still in Their First Marriage/Union, by Selected Characteristics, by Years since First Marriage/Union

CHAPTER III: CHILDBEARING/ABORTION

- III.1 Total Number of Live Births, by Age and Years since Marriage
- III.2 Mean Number of Children Ever Born, by Age and Selected Characteristics
- III.3 Age-Specific Fertility Rates and Total Fertility Rates (TFR), 1993 Czech Republic Reproductive Health Survey and 1991 Vital Statistics
- III.4 Total Fertility Rates, by Selected Characteristics
- III.5 Life Table Estimates of Percent of Respondents with Any Live Births before Selected Ages and Median Age at First Birth, by Selected Characteristics
- III.6 Life Table Estimates of Percent of Respondents with Any Live Births before Selected Marriage Durations and Median Duration at First Birth, according to Current Age and Education of Respondent
- III.7 Number of Children Planned When Respondent First Married, by Selected Characteristics, Women Ever Married
- III.8 Number of Children Planned When Respondent First Married, by Total Number of Children Currently Planned, Fecund Respondents Currently in Union
- III.9 Percent of Respondents Reporting That They Want to Have No More Children, by Selected Characteristics, Fecund Married Respondents
- III.10 Number of Additional Children Desired, according to Current Number of Living Children, Fecund Women 15 to 44 Years of Age
- III.11 Planning Status of Pregnancies Ending after 1987
- III.12 Outcomes of Reported Pregnancies Ending after 1987, according to Planning Status of Pregnancy

- III.13 Mean Number of Induced Abortions per Woman and Induced Abortions per 100 Known Conceptions, Based on Recent Official Statistics, Selected Countries of Central and Eastern Europe
- III.14 Reasons for Choosing to Have an Induced Abortion, for Abortions Taking Place after 1987

CHAPTER IV: FAMILY PLANNING

- IV.1 Percentage of Currently Married Respondents Familiar with Various Contraceptive Methods, by Education
- IV.2 Percentage of Currently Married Respondents Who Have Ever Used Various Contraceptive Methods, by Education
- IV.3 Percent of Currently Married Women Currently Using Any Contraceptive Method or Any Modern Method, by Selected Characteristics
- IV.4 Current Contraceptive Use, by Marital Status
- IV.5 Current Contraceptive Use, by Number of Living Children, Women Married or in Union
- IV.6 Current Contraceptive Use, by Age, Women Married or in Union
- IV.7 Current Contraceptive Use, by Education of Respondent, Women Married or in Union
- IV.8 Current Contraceptive Use, by Religion, Women Married or in Union
- IV.9 Current Contraceptive Use, by Whether Respondent Wants to Have Any More Children, Fecund, Nonpregnant Women Currently in Union
- IV.10 Source of Oral Contraceptives and IUDs, by Region, Current Users of Those Methods
- IV.11 Primary Reason for Not Using Contraception, by Marital Status
- IV.12 Percent of Women in Need of Family Planning Services, according to Two Definitions, by Selected Characteristics

- IV.13 Importance of Selected Factors in Couples' Decisions to Use Traditional Methods of Contraception, Current Users of Traditional Methods
- IV.14 Percent of Current Users of Withdrawal or Natural Family Planning Who Reported that Selected Factors Were Very Important or Somewhat Important in the Selection of Their Contraceptive Method, by Age, Education, and Method
- IV.15 Opinions of Current Users of Withdrawal or Natural Family Planning regarding the Effectiveness of Their Current Method Relative to Modern Methods, by Age, Education, and Method
- IV.16 Most Important Problem or Concern with Current Contraceptive Method, according to Method Used
- IV.17 Preference for Current or Other Method, Current Contraception Users and Method Preferred among Women Preferring Another Method
- IV.18 Contraceptive Failure Rates, in Percents, for Selected Methods after 12, 24, and 36 Months, for Segments of Use Beginning after 1987
- IV.19 Contraceptive Discontinuation Rates, in Percents, for Selected Methods after 12, 24, and 36 Months, for Segments of Use Beginning after 1987
- IV.20 Most Important Reason for Discontinuing Use of Contraceptive Method according to Method Used, All Reported Segments of Use Beginning and Ending after 1987
- IV.21 Interest in Surgical Sterilization among Fecund Women Wanting No More Children, by Selected Characteristics
- IV.22 Primary Reason for Lack of Interest in Surgical Sterilization among Fecund Women Wanting No More Children, by Education
- IV.23 Profile of Respondents Who Have Been Surgically Sterilized, Compared to All Respondents
- IV.24 Use of Infertility Services according to When Services Were Last Received and Selected Characteristics, Women Ever in Union

- IV.25 Percent of Respondents Receiving Selected Treatment/
Counselling among Those Who Visited a Physician or Clinic for
Infertility Problems, by Place of Residence and Age When Services
were Last Received

CHAPTER V: PREGNANCY, DELIVERY, AND INFANT HEALTH

- V.1 Respondents' Reaction to Most Recent Pregnancy, by Selected
Characteristics
- V.2 Spouse's/Partner's Reaction to Most Recent Pregnancy, by
Selected Characteristics
- V.3 Amount of Physical Work During Most Recent Pregnancy, by
Selected Characteristics
- V.4 Amount of Standing at Work During Most Recent Pregnancy, by
Selected Characteristics
- V.5 Amount of Stress at Work During Most Recent Pregnancy, by
Selected Characteristics
- V.6 Trimester When Prenatal Care Began During Most Recent
Pregnancy, by Selected Characteristics
- V.7 Number of Prenatal Care Visits during Most Recent Pregnancy,
by Selected Characteristics
- V.8 Who Provided the Most Prenatal Care during Most Recent
Pregnancy, by Selected Characteristics
- V.9 Where most of Prenatal Care Was Provided during Most Recent
Pregnancy, by Selected Characteristics
- V.10 Time Spent Travelling to the Place of Prenatal Care, by Means of
Transportation
- V.11 Waiting Time for Prenatal Care during Most Recent Pregnancy,
by Selected Characteristics
- V.12 Whether Respondent Received Ultrasound Examination during
Most Recent Pregnancy, by Selected Characteristics

- V.13 Degree to Which the Purpose of Ultrasound Examination Was Explained, by Selected Characteristics
- V.14 Whether Bed Rest was Prescribed during Most Recent Pregnancy and Length of Bed Rest, by Selected Characteristics
- V.15 Hospitalization during Most Recent Pregnancy and Length of Stay, by Selected Characteristics
- V.16 Percent of Women Hospitalized during Most Recent Pregnancy for Whom Various Conditions Were Diagnosed, by Parity
- V.17 Smoking Status of Women during Most Recent Pregnancy, by Selected Characteristics
- V.18 Percent Babies with Low Birthweight, by Parity
- V.19 Outcomes of Intended Pregnancies Ending since January 1988, by Selected Characteristics
- V.20 Assessment of Characteristics of Place of Delivery, by Parity
- V.21 Length of Delivery Relative to Expected Length, by Selected Characteristics
- V.22 Pain during Labor Relative to Expected Pain, by Selected Characteristics
- V.23 Percent of Respondents Who Thought They Received Enough Information on Pregnancy, by Selected Characteristics, by Parity
- V.24 Percent of Respondents Who Thought They Received Enough Information on Delivery, by Selected Characteristics, by Parity
- V.25 Who the Respondent Would Like to have Present at Next Delivery, by Selected Characteristics
- V.26 Whether the Respondent Would be Interested in Delivery at Home, by Selected Characteristics
- V.27 Most Important Problems during the First Week after Delivery, by Selected Characteristics

- V.28 Most Important Problems after Returning Home after Delivery, by Selected Characteristics
- V.29 Amount of Prenatal Counseling during the Most Recent Pregnancy, by Selected Characteristics
- V.30 Percent of Respondents' Children Born since January 1988 Who Were Breastfed, by Selected Characteristics
- V.31 Mean Duration of Breastfeeding for Children Born since January 1988, by Selected Characteristics

CHAPTER VI: YOUNG ADULT SEXUAL BEHAVIOR

- VI.1 Life Table Estimates of Percent of Women Who Have Ever Had Sexual Intercourse before Given Ages and Median Age at First Intercourse, according to Age at Interview
- VI.2 Percent of Women 15 to 24 Years of Age Who Have Ever Had Sexual Intercourse, by Selected Characteristics and Age at Interview
- VI.3 Relationship to First Sexual Partner and Length of Relationship at the Time of First Sexual Intercourse, according to Age at First Sexual Intercourse, Sexually Experienced Females, 15 to 24 Years Old
- VI.4 Contraceptive Method Used at First Sexual Intercourse, by Selected Characteristics, Sexually Experienced Females, 15 to 24 Years Old
- VI.5 Reason for not Using a Contraceptive Method at the Time of First Sexual Intercourse by Age at First Sexually Experienced Females, 15 to 24 Years Old
- VI.6 Relationship to Partner at Time of First Pregnancy, Whether She Married That Man, and Attitude of Partner Regarding the Pregnancy by Age at First Pregnancy, Women 15 to 24 Years Old with Any Reported Premarital Pregnancies
- VI.7 Attitude of Respondent's Family Regarding Her First Pregnancy, by Age at First Pregnancy, Women 15 to 24 Years Old with Any Reported Premarital Pregnancies

- VI.8 Age at Which Respondent's Parents First Discussed Family Planning Methods with Her, by Selected Characteristics, 15 to 24 Years Old Females

CHAPTER VII: REPRODUCTIVE HEALTH KNOWLEDGE AND ATTITUDES

- VII.1 Percent of Respondents Who Think That Women Should Be Able to Decide Whether to Have an Abortion for Any Pregnancy, by Selected Characteristics
- VII.2 Opinions on Whether Women Should Be Able to Have an Abortion under Given Circumstances, among Respondents Who Think There Should Be Restrictions on Whether Women Can Have Abortions
- VII.3 Percent of Respondents Who Think That Women Should Be Able to Decide Whether to Have an Abortion under Selected Circumstances, by Selected Characteristics
- VII.4 Percent of Respondents with Correct Knowledge of When a Woman Is the Most Likely to Conceive during Her Menstrual Cycle, by Selected Characteristics
- VII.5 Opinions about the Reliability of Oral Contraceptives, by Selected Characteristics
- VII.6 Opinions about the Safety of Oral Contraceptives, by Selected Characteristics
- VII.7 Opinions regarding Various Statements Concerning Oral Contraceptives
- VII.8 Opinions regarding Various Statements Concerning Women's Roles and Reproduction

CHAPTER VIII: WOMEN'S HEALTH

- VIII.1 Frequency of Receipt of Gynecologic Exams, by Region of Residence, Age, and Education
- VIII.2 Primary Reason for Receiving Gynecologic Exam Less Than Yearly

- VIII.3 Knowledge and Use of Breast Self-Examination (BSE) by
Selected Characteristics
- VIII.4 Current Cigarette Smoking Status, by Selected Characteristics
- VIII.5 Average Number of Cigarettes Smoked per Day, Selected
Characteristics Women Who Currently Smoke
- VIII.6 Average Weekly Alcohol Consumption, by Selected
Characteristics
- VIII.7 Percent of Respondents Who Are Substantially Overweight, by
Selected Characteristics

CHAPTER IX: TRANSMISSION OF HIV/AIDS

- IX.1 Percent of Respondents Who Think That the AIDS Virus Can Be
Transmitted in Various Ways, by Education of Respondent
- IX.2 Opinions about the Effectiveness of Condoms in Protecting
against HIV Infection, by Selected Characteristics
- IX.3 Respondents' Opinions about Their Own Risk of Contracting
AIDS, by Selected Characteristics

LIST OF FIGURES

CHAPTER II: RESPONDENT CHARACTERISTICS

- I.1 Education of Respondents by Age Group
- II.2 Religion of Respondents by Age Group
- II.3 Percent of Respondent Households With Selected Possessions/
Amenities
- II.4 Percent Ever Married Before Selected Ages

CHAPTER III: CHILDBEARING/ABORTION

- III.1 Total Fertility Rates, Selected Central and Eastern European
Countries
- III.2 Fertility Rates, 20-24 Year-Old Women, Selected European
Countries
- III.3 Fertility Rates, 30-34 Year-Old Women, by Selected European
Countries
- III.4 Number of Additional Children Desired According to Number of
Living Children, Fecund Women

CHAPTER IV: FAMILY PLANNING

- IV.1 Current Contraceptive Use Among Married Women
- IV.2 Current Contraceptive Use Among Married Women by Number of
Living Children
- IV.3 Importance of Selected Factors in Decision to Use Traditional
Contraception
- IV.4 Interest in Surgical Sterilization Among Women Wanting No More
Children by Number of Living Children

CHAPTER V: PREGNANCY, DELIVERY, AND INFANT HEALTH

- V.1 Percent of Respondents With No Prenatal Care During Their First Trimester, By Region
- V.2 Percent of Respondents With No Ultrasound Exam During Her Most Recent Pregnancy
- V.3 Selected Indicators and Aspects of Prenatal Care During Respondents' Most Recent Pregnancy
- V.4 Percent Babies With Low Birthweight by Region
- V.5 Most Important Problem for Respondent During the Week Following Her Most Recent Birth
- V.6 Most Important Problem for Respondent After Returning Home Following Delivery

CHAPTER VI: YOUNG ADULT SEXUAL BEHAVIOR

- VI.1 Percent of Women Who Ever Had Sexual Intercourse Before Given Age By Age at Interview, 15-24 Year-Old Respondents
- VI.2 Contraceptive Method Used at First Sexual Intercourse, By Age at First Sex And By Education, 15-24 Year-Old Respondents

CHAPTER VII: REPRODUCTIVE HEALTH KNOWLEDGE AND ATTITUDES

- VII.1 Percent of Respondents Who Think Women Should Be Able to Decide Whether To Have An Abortion For Any Pregnancy, By Education And By Religion
- VII.2 Opinions About The Reliability of Oral Contraceptives
- VII.3 Opinions About The Safety Of Oral Contraceptives

CHAPTER VIII: WOMEN'S HEALTH

- VIII.1 Cigarette Smoking Status By Education

SUMMARY

Introduction

From February to July of 1993, a nationally representative household survey of 15 to 44 year-old women, the Czech Republic Reproductive Health Survey (CRRHS), was conducted throughout the Czech Republic. This survey was undertaken in order to explore important issues related to reproduction and women's health in the Czech Republic. The survey was carried out at a time when major reforms in the Czech health care system were underway. It was felt that in this period of change and uncertainty it was important that policymakers and health care providers be aware of the current status of reproductive health and related topics in the population and the use of and need for various health services. The 1993 CRRHS was intended to serve several purposes, among them: updating basic information regarding such topics as family planning use and needs, use of maternal and child health services, and selected women's health issues; determining reproductive health needs for the country as a whole and for population subgroups; and more closely examining some reproductive health topics of special interest, for instance the reasons for high incidence of induced abortion and low prevalence of modern contraceptive use.

The 1993 CRRHS was a collaborative effort between several organizations, all of which contributed to its success. Among those organizations were: the United States Agency for International Development, which financed the survey, Factum (a Czech public opinion research firm), the Czech Statistical Office, The World Health Organization Collaborating Center for Perinatal Medicine at the Czech Institute for Maternal and Child Health Care, and the Division of Reproductive Health of the United States Centers for Disease Control and Prevention (CDC).

The CRRHS questionnaire covered a wide range of topics related to reproductive health in the Czech Republic. The subject areas included were: Pregnancy and childbearing history; knowledge and use of contraceptive methods; use of maternal and child health services; sexuality, contraception, and pregnancy among young adults; women's health and reproduction issues; knowledge about HIV/AIDS and its transmission; and social, economic, and demographic characteristics.

Methodology

The 1993 CRRHS was designed to collect information from a representative sample of 15 to 44 year-old women living in households in the Czech Republic. Selection of survey respondents was made using a three-stage cluster design. In the first stage of sampling, 332 census enumeration districts (EDs) were systematically selected from throughout the country. The second stage consisted of selection of a cluster of individual dwelling units within each sampled ED. Finally, in homes in which more than one eligible women

lived, one woman was randomly chosen as a respondent.

Data collection took place from February until June of 1993 and was done by intensively trained female interviewers from throughout the Czech Republic. Each interviewer was assigned to visit selected households in sample sectors (EDs) easily accessible from her home. Interviews were conducted at respondents' homes and generally lasted between 30 and 60 minutes. At the Czech Statistical Office questionnaires were entered onto computer data files and concurrently edited, using software developed at CDC.

A total of 12,747 dwelling units were visited in the 332 sample sectors. Women eligible for interview were identified as residing in 38.2% of these units. Of the 4,870 households identified as containing at least one 15-44 year-old women, interviews were completed in 4,497, for a response rate of 92.3%. Of the women selected, 5.3% refused to be interviewed and 1.6% were never found at home after repeated visits.

Respondent Characteristics

The residential characteristics, age, and marital status distributions of the sample appears to have been very representative of the 15-44 female population of the Czech Republic. Approximately two-thirds of respondents were married or living in a consensual union at the time of interview, while one-fourth had never been married. By ages 35-39 about one in ten women were currently divorced or separated. The typical age at marriage in the Czech Republic remains quite young, with the median age just 20.7 years. Marriages are highly concentrated between 18 and 25 years of age: only 6% of women were married before age 18, but only 12% were not married by age 25. It is evident that the age at marriage had not changed in recent years, being almost identical for all cohorts of respondents.

About four of every ten respondents had received a secondary school diploma. Educational attainment has been increasing, demonstrated by the fact that the percentage who attended only primary school fell from 24% to 7% in a 20-year period, while the percentage who obtained a secondary school diploma rose from 41% to over 50% in the 20-24 and 25-29 year-old cohorts. Almost two-thirds of respondents said they had no religious affiliation and 31% said they were Roman Catholic. The vast majority of respondents lived in homes that contained a bathroom or shower (98%) and a color television (89%). In addition, most homes had an automatic washing machine (75%) and an automobile (64%). Thirty-three percent of homes had a telephone.

Childbearing/Abortion

Vital statistics on births in the Czech Republic are virtually complete, so that measuring current fertility rates was not a major objective of the survey. However, the data collected in the survey provide much more detailed information on reproduction, as well as information on aspects of childbearing not covered by vital statistics. Each respondent

was asked to give a history of all the pregnancies she had ever had, including information about each pregnancy and its outcome.

Both childlessness and large families are rare in the Czech Republic. Only 3% of respondents had given birth to four or more children. Among women 35 to 44 years of age, the vast majority of whom have completed their childbearing, there is little variation in family size, with 57% having had two children. Few of the oldest respondents have had very small families (3% with no children and 12% with one child among 40-44 year-olds). For most women/couples childbearing starts soon after marriage (only 28% of women married less than five years were still childless) and stops at relatively early ages as well.

The total fertility rate (TFR) for the period leading up to the CRRHS (1990-1992), based on data from the survey was 1.87 births per woman, identical to the rate computed from Czech Republic vital statistics. Childbearing is heavily concentrated in the 20-24 and 25-29 year age groups, with very little childbearing taking place after ages 30-34. Compared to the rest of Europe, Czech women have unusually high fertility before ages 25-29 and unusually low fertility after ages 30-34. There was a clear inverse relationship between education and fertility. The total fertility rate was higher among Catholics (2.1) than among those with no religious affiliation (1.8).

Childbearing tends to start early in life, with the median age at first birth being 22.0 years. As with age at marriage, there is no indication that the age at first birth has been increasing. Within five years of the date of first marriage, only 7% of women had not had a live birth.

There is a strong preference for a two-child family among women throughout the population. Few women desired to remain childless or to have more than three children. Sixty-nine percent of fecund, married respondents desired to have no more children. Among those with two living children the figure was 86% and among those with at least three children it rose to 94%.

Planning Status of Pregnancies

After adjustments were made for the underreporting of abortions, we estimated that only 51% of recent pregnancies were planned, 16% were mistimed (i.e., occurred before they were planned), and 28% were unwanted (i.e., in excess of the number wanted), with the remainder unspecified. The proportion of pregnancies that were unwanted rose sharply after ages 30-34 and once women had at least two living children. Most mistimed pregnancies occurred relatively before ages 25-29. After correction for underreporting of abortion it appears that slightly more than half of mistimed pregnancies were aborted and that only about 4% of unwanted pregnancies resulted in a live birth.

Knowledge and Use of Contraception

Knowledge of pregnancy prevention methods was not an important obstacle to use of contraception. Knowledge of condoms and oral contraceptives was nearly universal and was only slightly lower for the IUD, withdrawal, and natural family planning. Even the least widely known method asked about, vasectomy, was familiar to two-thirds of respondents. Among women currently married or living with a partner, 92 percent had used contraception at some time during their lives.

Sixty-nine percent of women who were married or living with a man at the time of interview were currently using some method of family planning. Couples with two or three children were the most likely to be using contraception (75 and 78%, respectively), while only 29% of childless couples were using. Current contraceptive use increased with level of education, at least up to the level of secondary school completion. Women who were not affiliated with a religion were somewhat more likely to be using than women belonging to an organized religion, though there was virtually no difference in contraceptive prevalence between Catholics who regularly attended church and those who did not.

The most commonly used contraceptive methods were reported to be withdrawal (22% of married/in union women), condoms (17%), the IUD (15%), and oral contraceptives (8%), which together accounted for over 90% of all contraception. No other methods were currently employed by more than 3% of married women. About 3% of women had been contraceptively sterilized. No respondents reported that their partner had undergone a vasectomy.

Reasons for not Using Contraception

About two-thirds of women not using contraception at the time of interview gave reasons that related to either a lack of current sexual activity or their pregnancy status (i.e., unable to become pregnant, trying to become pregnant, or currently pregnant). Among the broad assortment of other reasons given, the most common was fear of health effects related to contraception. Religion and inability to obtain methods were rarely cited. Based on these results, it is estimated that as many as one-third of Czech women who were not using contraception were at some risk of unintended pregnancy due to non-use of contraception.

Not surprisingly, a lack of sexual activity was the principal reason for not using contraception among women who were not currently married. Among women married or in union, just over half of nonusers reported a reason related to pregnancy status or lack of sexual activity. Reasons such as fear of health effects, difficulty in getting pregnant, and postpartum/breastfeeding were much more common among these women than among those not in union.

Need for Family Planning Services

The concept of "women in need of family planning services" is useful in determining the extent to which services are reaching those who are at risk of becoming pregnant, but who do not desire to do so. If only women who are sexually active, fecund, not pregnant, not desiring to become pregnant and using no contraceptive method are considered to be in need, then 10% of Czech women of reproductive age are in need of family planning services. If users of withdrawal and natural methods are also included, the figure rises to 31%.

Use of Less Effective Methods

Even though overall contraceptive prevalence is high in the Czech Republic, there is considerable reliance on the use of methods of relatively low effectiveness, particularly withdrawal. This reliance on methods of low reliability is, no doubt, one of the key factors in bringing about high rates of unintended pregnancy and induced abortion. Of six factors about which users of less reliable methods were asked, only a fear that negative health effects associated with the use of modern methods (75%) was an important consideration in method selection. Religious factors were not a consideration at all for 97% of these women. Many users of these methods were overly confident in their reliability relative to modern methods. Almost half of these respondents thought that their method was at least as effective as highly effective modern methods.

Problems with Current Contraceptive Methods

Eighty-five percent of contraceptive users said they had had no major problems with their current method. This figure was fairly consistent between methods, ranging from 90% for NFP down to 80% for a combination of condoms and withdrawal. About half of those with concerns reported that reliability/quality was their major one. Consistent with this low level of reported concerns about current methods, 73% of current contraceptors stated that they would prefer to be using their current method rather than switching to another one. Among those desiring to switch from their current method, 40% preferred to use oral contraceptives, followed by the IUD (25%). Surprisingly, the third most preferred method (15% of those preferring another method) was implants, which were not yet available in the Czech Republic at the time of the survey.

Contraceptive Sterilization

Like most of the other formerly communist states of central and eastern Europe, the Czech Republic has low rates of surgical sterilization, both tubal ligation and, to an even greater extent, vasectomy. Only three percent of married women of childbearing age reported having undergone surgical sterilization, while no women reported their partners to have had a vasectomy, despite the fact that most women complete their desired childbearing

many years before becoming unable to bear children. The major exception was the relatively small group of women who have at least four living children, over one-fourth of whom had been sterilized. Only 8% of the respondents who desired no more children reported being interested in undergoing tubal ligation, in addition to the 3% already sterilized.

The reason most often given by respondents for not being interested in surgical sterilization was simply that they had not thought about undergoing the procedure. The second most commonly given reason was a fear of health risks associated with the procedure, followed by a fear that the woman/couple may decide to have another child and a fear of surgery.

Prenatal Care

Heavy physical exertion and emotional stress are thought by some to increase the risk of adverse pregnancy outcomes. Overall, 15% of women described themselves as having done much physical work and 27% said they did a large amount of standing during their jobs. The amount of both physical work and standing decreased sharply as educational level increased. Seventeen percent of women said they were under much stress on their jobs during pregnancy and 32% said they were under moderate stress.

The Czech Republic has an impressive record regarding prenatal care coverage. Less than 1% of women with a live birth since the beginning of 1988 received no prenatal care during their most recent pregnancy leading to a live birth. Ninety-four percent of those women began receiving prenatal care during their first trimester, while only 5% waited until after the first trimester. The percentage of women receiving early prenatal care ranged from 97% in South Bohemia down to 90% in North Bohemia and 91% in East Bohemia. Women who were less likely to begin care in the first trimester included those who: were less than 20 years of age or between 35 and 44 years; were not currently married; had only a primary education; were affiliated with a religion other than Catholicism; or had low household income.

Not only did women tend to begin prenatal visits early, but 91% reported making at least 10 prenatal visits during pregnancy, with only 1% reporting fewer than six visits. The proportion of women making at least 10 prenatal visits was high across all geographic and socioeconomic categories.

Ultrasound Examinations

Ninety-two percent of women who received prenatal care reported that they underwent an ultrasound examination during their last pregnancy leading to a live birth. The percentage having an ultrasound exam varied from 98% in South Bohemia and 97% in South Moravia down to 85% in North Moravia and 86% in West Bohemia. The procedure was more commonly done for the youngest (15-24 years) and oldest (40-44)

groups of women. However, one-fourth of women who had undergone ultrasound said that no one had explained the reason for doing this procedure and another one-fifth had only had the procedure explained to them slightly.

Hospitalization during Pregnancy

Hospitalization during pregnancy (other than for labor and delivery) is a common phenomenon in the Czech Republic. Twenty-nine percent of women giving birth since 1988 were hospitalized during some part of their pregnancy, with a mean length of stay of almost one month and a median of two weeks. Although older mothers were no more likely than others to be hospitalized during pregnancy, they tended to be hospitalized for the longest periods of time (mean = 57 days, median = 21 days). The conditions most commonly cited as being responsible for hospitalization during pregnancy were bleeding during the first half of pregnancy (39% of hospitalizations) and early or false labor (33%).

Smoking during Pregnancy

Sixty-eight percent of respondents with recent live births were not smokers when they became pregnant. An additional 21% reported that they stopped smoking upon learning they were pregnant, with the remaining 11 % continuing to smoke during pregnancy. About one-third of this group smoked at least 10 cigarettes per day. As educational level increased, the likelihood that a woman smoked during pregnancy decreased sharply, from 32% among women with only a primary school education, to 2% for those who attended university.

Birthweight

The incidence of low birthweight (LBW) (i.e., under 2,500 grams) among recent births was 5.8% among survey respondents. The incidence of LBW was slightly higher among primiparous women (7%) than among multiparous women (5%). Although this difference was not large, the LBW rate was lower among multiparous women in all but a few categories of survey respondents. The highest LBW rates were found in cities of at least 20,000 population, in North Moravia, among the oldest (40-44 year-old) and youngest (15-19 year-old), single women, and poorly educated women.

Labor and Delivery

The CRRHS questionnaire also inquired into respondents' opinions regarding services received for recent pregnancies and deliveries. Women were asked their impressions about the facility in which they most recently delivered in regard to: distance from home, physical facilities, crowdedness, attentiveness of staff, and competence of staff. Each of the five characteristics was rated as "good" by between one-half and three-quarters of respondents. The characteristic eliciting the most "poor" ratings (20%) was

crowdedness. The aspect of care about which respondents were the most satisfied was the competence of the staff.

Overall, four-fifths of respondents said that they received adequate information regarding their pregnancy. Multiparous women were somewhat more satisfied than primiparous women. The level of satisfaction with information provided about delivery was 68%, lower than for information about pregnancy. Among multiparae 72% were satisfied, compared with 61% of primiparae.

Problems in the Post Partum Period

Forty percent of respondents said they had no major problems during the first week following delivery. The problems most commonly mentioned related to the health and care of the baby (18%), the woman's own health (17%), and breastfeeding (11 %). Care of the child was mentioned most commonly by the youngest and oldest respondents and by women who had never been married. Breastfeeding concerns stood out as a problem among university educated and primiparous women.

Fifty-eight percent of respondents said they had no major problems after returning home from the hospital. The problems mentioned most frequently were her own health (10%), health of the child (8%), care of her child/children (7%), and home environment (5%). Health of the child was a major concern among the youngest women and single women. Care of the child was most commonly mentioned by university educated and primiparous women, but rarely by multiparous women.

Breastfeeding

Ninety-one percent of children born to respondents since the beginning of 1988 were breastfed for at least some period of time. The proportion breastfed was lowest for infants with mothers ages 15-19 (84%) or ages 40-44 (71 %). Despite the fact that relatively few infants were never breastfed, the mean duration of breastfeeding was short: 3.7 months for all children and 4.2 months for children who were reportedly breastfed.

Young Adult Reproductive Health

The survey estimate of the median age at first intercourse was 17.5 years. It appears that the age at which females begin having sexual relations has been decreasing, since the median age was 0.5 years less for those 15-19 years of age than for those five years older. Sexual intercourse before age 15 is still fairly rare for girls. By about age 21, only a small fraction of women were not yet sexually experienced. The proportions reporting that they had had sexual intercourse were 36% of 15-17 year-olds, 82% of 18-19 year-olds, and 98% of 20-24 year-olds. Bohemian women tend to become sexually active slightly earlier than Moravian women (39% experienced, as opposed to 31% among 15-19 year-olds). Likewise, women espousing no religion start sexual relations earlier than

Catholics (37% compared with 32%). Clearly premarital sexual intercourse has become the norm in the Czech Republic: Over 99% of sexually experienced women first had intercourse before they were married.

Forty-two percent of young adult respondents reportedly used no form of contraception the first time they had sexual intercourse. Among those who did use contraception, over 90% used either withdrawal or condoms. Of those females who did not use contraception at first intercourse, 44% said the reason was that they did not expect to have sex, while 27% said they did not think it was possible for them to become pregnant.

About one-half of 15-24 year-old respondents reportedly had ever been pregnant at the time of interview and about half of those said they were not married at the time they became pregnant. (Because of abortion underreporting, however, the actual percentages may be higher than indicated in the survey.)

Reproductive Health Knowledge and Attitudes

The rate of induced abortions in the Czech Republic is quite high, with official statistics for 1992 showing 48 abortions per 1000 15-44 year-old women and 78 abortions for every 100 live births. Abortion laws are permissive, although minor restrictions have been added in recent years and the cost of having an abortion performed has increased substantially.

An overwhelming majority of childbearing age women (85%) felt that women/couples should have the right to decide whether to have a pregnancy terminated by means of induced abortion for any reason they choose. Every segment of the population examined was similarly opposed to legal restrictions on a woman's obtaining an abortion. Even among Catholic women who attend mass regularly, seven of every ten felt there should be no limitations on access to abortion. Even among the 14% of respondents who thought that there should be some restrictions on access to abortion, an overwhelming majority felt that abortion was justified under at least some circumstances, particularly if a pregnancy endangered a woman's life or if there was risk of a fetal defect.

Only about half of respondents said that a women who took oral contraceptives correctly could be completely or almost sure that she would not become pregnant and one of every five women indicated that the reliability of oral contraceptives was quite low. About one-fifth of women felt that oral contraceptives were unsafe, while a slightly smaller percentage said they did not know if they were safe.

Fifty-three percent of women were aware that oral contraceptives make menstrual periods more regular, while 42% felt that oral contraceptives cause weight gain. Sizable minorities of respondents agreed with the statements that oral contraceptives: increase

the risk of getting cancer (25%); are bad for blood circulation (21%); can cause infertility (16%); and cause nervousness (13%). About half of respondents for each of these four statements said they did not know whether they were true.

Respondents were asked whether they agreed or disagreed with selected statements regarding women's roles and reproduction. Few women agreed with the statements that "Child care is women's work" and "A woman should be a virgin when she marries", while most agreed that it is all right for women not to have children if they so desire. These results indicate that most Czech women do not feel that women should be restricted to traditional roles.

Women's Health

It is recommended that women of childbearing age receive complete gynecologic examinations on a yearly basis; 60% of respondents reported that they go for gynecologic exams at least that often. At the other extreme, 15% of women said they never go for such exams. The group least likely to be examined was 15-19 year-olds, among whom only 30% had yearly exams and 59% never had them. Two-thirds of women who did not receive yearly exams said the reason for this was that they did not have any gynecologic problems.

It has been well established that cigarette smoking is an important risk factor for cancer, respiratory diseases, and cardiovascular disease, among other chronic conditions. Thirty percent of CRRHS respondents reported that they currently smoked. Another 12% had smoked in the past, but no longer did. Smoking prevalence was highest in North Bohemia and Prague and lowest in Moravia and adjacent East Bohemia. Rates also tended to be higher in large cities than elsewhere. Only 19% of smokers were light smokers, that is, they typically smoked fewer than five cigarettes per day. The largest proportion of smokers, 44%, smoked 10 to 19 cigarettes per day, while 16% smoked at least one pack per day.

Relatively few Czech women reported chronically consuming large amounts of alcoholic beverages. Only 3% of respondents said that they consume an average of more than seven alcoholic drinks per week. Almost eight of every ten women consumed either fewer than three or three to seven drinks per week, while one-fifth said they did not drink at all.

Based on reported heights and weights, 15% of respondents met the criterion for being substantially overweight. This percentage was lowest in Prague (10%) and tended to be highest in places with populations under 5,000 people (21%).

Knowledge of Transmission of HIV/AIDS

Even though, according to official statistics, the incidence of HIV infection was not yet

very high in the Czech Republic at the time of the CRRHS, a threat of increased spread of the infection existed. This was especially true in light of the increased numbers of people entering the Czech Republic from other countries and the newfound freedom of Czechs to travel outside their own country following the fall of the communist government. The CRRHS included a series of questions intended to measure the knowledge of Czech women regarding HIV/AIDS in general and the means by which HIV is transmitted to individuals.

All but five respondents (99.9%) reported that they had heard of AIDS. In general, there appeared to be a high level of awareness of those means by which HIV is known to be transmissible—sharing needles (98%), male homosexual relations (96%), and heterosexual relations (94%). Receiving a transfusion, which in many settings can lead to transmission, was felt to be a cause of AIDS by 89% of women. Almost half of women thought a person could be infected by giving blood or by going to a physician or dentist, behaviors that cannot transmit infection except under unusual circumstances. Among those methods asked about by which the virus is known not to be transmitted, the proportion who felt that HIV could be spread ranged from 28% for kissing on the mouth down to 3% for shaking hands.

Only about three-fourths of respondents felt that condoms provided excellent or good protection against transmission of HIV. Eighty-three percent of women felt they had no risk of becoming infected with the AIDS virus, 3% felt they had a low risk, and 5% felt they had a high risk. There was very little variation in these percentages across the population.

I. INTRODUCTION/SURVEY METHODOLOGY

In 1993, a national survey of reproductive health issues was conducted in the Czech Republic. The Czech Republic, with a population of 10.4 million people, lies in central Europe, covering an area of 78,864 square kilometers (30,590 square miles). Since 1918, the Czech Republic had constituted the western three-fifths of Czechoslovakia prior to the January 1993 breakup of that country. The country consists of two historical areas, Bohemia in the west, which is divided into six regions, and Moravia and Silesia in the east, divided into two regions.

At the time the survey was carried out, the crude birth rate (births per 1,000 population per year) was estimated to be about 12 per 1,000 and the crude death rate was about 11.1 per 1,000, with the annual rate of natural increase being about 0.1 percent (Population Reference Bureau 1994). According to the 1991 Census, 75% of the Czech population lived in places classified as urban, with 22% of people living in cities with at least 100,000 people (Czechoslovakia Federal Statistical Office 1991). Average life expectancy was estimated to be 77 years among females and 69 years among males (National Public Health and Hospital Institute 1994). Ethnically, the Czech Republic is highly homogeneous, with the overwhelming majority of residents (about 95%) identifying themselves as Czech (or Bohemian or Moravian). The largest numbers of non-Czech residents are Slovak, Romany (Gypsy), Polish, and Silesian. The Czech language is the primary language for all but a very small percentage of the entire population.

Survey Background

The 1993 Czech Republic Reproductive Health Survey (CRRHS) was a nationally representative household survey of women 15 to 44 years of age, undertaken in order to explore important issues related to reproduction and women's health in that country. This survey was carried out at a time when substantial reforms in the Czech health care system were taking place or were being considered (Albert 1992; Massaro et al 1994; NPHHI 1994; Raffel 1992).

The reforms taking place have largely grown out of the fall of communism and the resultant changeover to a free market economy, the rise of democracy, and the breakup of the Czechoslovak Federation. Among the changes underway at the time of the survey were: Reforming the administrative structure of the health care system from one in which most responsibility for health care delivery resided at the regional level to one in which districts, municipalities, and the national Ministry of Health take over those responsibilities; privatization of the health sector, including hospitals, polyclinics, physician services, and pharmacies; and, reforming health care financing by introducing a system of health insurance, rather than having health care funding completely centralized and financed from general tax revenues. In this period of change and uncertainty it is important that policymakers and health care providers be aware of the current status of reproductive health

throughout the population and the use of and need for various services related to family planning and maternal and child health.

The CRRHS was intended to serve several purposes. The last large, national survey covering a wide array of demographic and reproductive health topics was the 1977 Czechoslovakia Fertility Survey (World Fertility Survey 1978). Because no such surveys had been conducted in recent years on family planning and reproductive health in Czechoslovakia or the Czech Republic, this survey was the first opportunity in many years to update basic measures, such as contraceptive prevalence, unmet need for contraception, use of maternal and child health services, and many others. These measures constitute important information in determining a population's needs and problems. In recent years, at least two useful surveys have been carried out, but both were relatively limited in regard to both content or sample size (Kraus 1987; Uzel, Ketting, et al. 1992). The size and scope of the 1993 CRRHS allows highly detailed analysis with regard to many reproductive health questions for subgroups of the Czech population.

There are several issues in the area of reproductive health in the Czech Republic that were known to merit close examination prior to carrying out the 1993 CRRHS. Official statistics show that the rate of induced abortion and the ratio of abortions to live births has remained very high (although they appear to have declined recently). Reducing the number of abortions would be in the best interest of public health in the Czech Republic, regardless of one's views regarding the legality and morality of abortion. It appears that modern, highly reliable contraceptive methods are not as widely used as they are in most other developed countries. The failure to use such methods contributes to high rates of unintended pregnancy. Furthermore, pregnancy and unprotected intercourse among Czech women under 18 years of age appear to be quite widespread. In addition, relatively little data exist on the extent of certain behaviors that affect the health of women and their infants, such as smoking, alcohol consumption, and the prevalence of being overweight. The CRRHS was designed to provide information that would contribute to strategies for addressing these issues in the Czech Republic.

Besides information about health status and practices, the 1993 CRRHS measured attitudes and opinions about the quality and types of reproductive health services offered, restrictions on the availability of induced abortion, and problems associated with the use of modern contraceptives. Also, as AIDS emerges as an important public health problem, it is important to find out how well informed women are about the means of transmission and prevention of this disease.

Organizational Structure

The 1993 Czech Republic Reproductive Health Survey was a collaborative effort between several organizations and many individuals. It would not have been successful without the contributions of each and the cooperation that existed between them. Funding for the 1993 CRRHS was provided by the United States Agency for International Development, through a

contract with the World Health Organization's European Office. Fieldwork was conducted by Factum, a Czech public opinion research firm, which coordinated the recruitment and training of interviewers and all aspects of data collection. The Czech Statistical Office (CSO) was responsible for preparing the sampling frame and carrying out data entry and editing, as well as providing the physical facilities for the survey. The World Health Organization (WHO) Collaborating Center for Perinatal Medicine at the Czech Institute for Maternal and Child Health Care assisted with the design of the questionnaire and data analysis and was responsible for financial accounting. The Division of Reproductive Health of the United States Centers for Disease Control and Prevention (CDC) provided assistance in survey design, questionnaire development and survey implementation and played the leading role in analysis of the data and report preparation. CDC, CSO, and Factum in developing the survey methodology. All of the above-mentioned organizations collaborated on the development of the questionnaire, data analysis, and report preparation. In addition, many other individuals provided valuable assistance or information that was important in the development and implementation of the survey.

Questionnaire Content

The CRRHS questionnaire covered a wide range of topics in the area of reproductive health and related areas. The specific subject areas included were:

- ▶ Pregnancy and childbearing history
- ▶ Knowledge, past and present use of, and need for contraceptive methods
- ▶ Use of maternal and child health services
- ▶ Sexuality, contraception, and pregnancy among young adults (15-24 years)
- ▶ Women's health issues
- ▶ Knowledge and attitudes about selected issues in reproduction
- ▶ Knowledge about HIV/AIDS and its transmission
- ▶ Social, economic, and demographic characteristics

An English translation of the CRRHS questionnaire is included as Appendix A of this report. Most of the results in this report are presented according to geographic, social, economic, and demographic characteristics, making it possible to identify the population segments where health needs or problems tend to be most acute.

Sample Selection

The 993 CRRHS was designed to collect information from a nationally representative sample of women of reproductive age from throughout the Czech Republic. The universe from which women were selected consisted of all females between the ages of 15 and 44 years living in households in the Czech Republic at the time that the survey was carried out. Although some pregnancy and childbearing takes place outside the ages 15 to 44, because of

the relative rarity of these events at those ages, it was decided that it would be most efficient to limit the sample to women between 15 and 44 years of age.

Selection of survey respondents was made using a three-stage cluster sampling design. First, 166 pairs of basic census enumeration districts (ED) (332 districts in all) were systematically selected from throughout the country, with the probability of selection proportional to the population of each census enumeration district. These 332 ED constituted the primary sampling units (PSU) for the survey. The sample included ED in all but one of the Czech Republics 85 districts. The Czech Statistical Office then made listings available of all known dwelling units in each of the selected ED. These listings were taken from the 1991 Population and Housing Census. Because the selection of PSUs was done proportional to population size and no strata were oversampled, the sample was geographically self-weighting.

In the second stage of selection, individual dwelling units were sampled from each ED chosen in the first stage. Because of slightly different mean numbers of 15 to 44 year-old women per household according to size of community, the number of dwelling units selected in an ED depended on the population of the community in which the ED was located, as follows:

<u>Community Population</u>	<u>Selected Dwelling Units</u>
1-4,999	43
5,000-19,999	37
20,000+	39

One dwelling unit was randomly chosen from the each PSU. That unit and the 42, 36, or 38 units (depending on the community population) listed next constituted the sample of housing units to be visited. No substitution was made for dwellings that were unoccupied or where nobody was at home. By varying the number of units selected according to the average number of women per household we were able to hold fairly constant the average numbers of women living in selected ED across population size categories. If the selected ED had fewer households than were needed to constitute a sample sector, all dwelling units were sampled and another ED (the one with the next highest number, which was usually geographically contiguous) was added to the sample.

The third and final stage of sampling consisted of the selection of individual respondents. In households in which more than one woman between the ages of 15 and 44 lived, the household respondent was asked for a listing of all such females. One of those women was randomly selected by the interviewer to be the survey respondent. During analysis each record was weighted by the number of women of childbearing age in the household, to compensate for the fact that only one woman per household was selected with probability inverse to the number of women in the household.

Data Collection

Data collection for the CRRHS was done by about 150 female interviewers from throughout the Czech Republic. Each interviewer received intensive training in interviewing techniques, field procedures, and proper administration of the CRRHS questionnaire. Each interviewer attended one of three training courses held immediately before the beginning of field work, two in Prague and one in Brno in January and February of 1993. These sessions included practice interviews with community residents to ensure that interviewers were able to administer the questionnaire competently.

Data collection lasted from February until June of 1993. Each interviewer was assigned to visit selected households in from one to three sample sectors easily accessible from her home. Interviews were conducted at respondents' homes and generally lasted between 30 and 60 minutes. Completed questionnaires were sent by the interviewer to her supervisor, who reviewed each questionnaire and, if it was satisfactorily completed, forward it to the Czech Statistical Office.

Data Entry/Editing

At the Czech Statistical Office questionnaires were reviewed once more by coordinators and sent for data entry. Questionnaire data was entered onto computers by three clerks, using SURVEY, a data entry/editing package developed at CDC. The SURVEY package is used to edit data as they are entered, by identifying data items that are outside of their allowable range or inconsistent with other information on the questionnaire.

Response Rates

A total of 12,747 dwelling units were visited in the 332 sample sectors. Women eligible for interview were identified as residing in 38.2% of these units ([Table I.1](#)). In 52.2% of the households visited there were no 15-44 year-old women resident. Nobody was home at 4.0% of units and 3.0% of units were uninhabited. Residents refused to be interviewed at 2.7% of households. As the size of community increased, the percent of refusals increased slightly and the percent of households with no eligible females decreased.

Of me 4,870 households identified as containing at least one 15-44 year-old female, interviews were completed in 4,497, for a completion rate of 92.3%. Of those women selected, 5.3% refused to be interviewed and 1.6% were not found at home, even after repeated visits. The completion rate fell slightly as community size grew, from 93.8% in towns under 5,000 population to 91.3% for cities with at least 20,000 people.

In regard to residential characteristics, the sample appears to have been highly representative of the 15-44 year-old population of the Czech Republic. Of the Czech Republic's nine regions, only in Prague was the difference between the 1991 Census and CRRHS proportion of the female 15-44 year-old population as much as two percentage points ([Table I.2](#)).

Likewise, the census and survey distributions of size of place of residence were very similar.

TABLE I.1
Final Interview Status, by Size of Place
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Interview Status	Total	Size of Place		
		<5,000	5,000-19,999	20,000+
Household Selection				
Eligible Woman Identified	38.2	38.1	39.6	37.6
No Eligible Women in Household	51.9	54.8	52.5	49.7
Resident Not Home	4.0	2.0	2.6	6.1
Refusal	2.7	1.7	2.3	3.5
Vacant Household	3.0	3.0	3.0	2.8
Other	0.1	0.1	0.1	0.2
Total	100.0	100.0	100.0	100.0
Number of Households	12,747	4,114	2,647	5,986
Households with Eligible Women Identified	4,870	1,569	1,048	2,253
Individual Selection				
Completed Interview	92.3	93.8	92.3	91.3
Selected Woman Not Home	1.6	1.1	1.7	1.9
Selected Woman Refusal	5.3	4.3	5.4	5.9
Other	0.8	0.8	0.6	0.9
Total	100.0	100.0	100.0	100.0
Number of Women Interviewed	4,497	1,472	967	2,058

TABLE I.2
Region of Residence and Size of Place of Residence,
1993 Czech Republic Reproductive Health Survey Respondents
and Women 15-44 Years Old in the 1991 Czech Republic Census
(Percent Distributions)

Region/Size of Place	1991 Census	1993 CRRHS	
	Percent	Percent	No. of Women
Region			
Bohemia	61.0	60.2	(2,737)
Prague	11.9	9.9	(456)
Central Bohemia	10.6	10.3	(460)
South Bohemia	6.7	6.4	(295)
West Bohemia	8.4	9.2	(413)
North Bohemia	11.6	12.8	(582)
East Bohemia	11.8	11.6	(531)
Moravia	39.0	39.8	(1,760)
South Moravia	19.6	20.0	(873)
North Moravia	19.4	19.7	(887)
Size of Place			
Less than 5,000	33.1	33.6	(1,472)
5,000-19,999	19.0	21.0	(967)
20,000 +	48.0	45.4	(2,058)
Total	100.0	100.0	(4,497)

II. RESPONDENT CHARACTERISTICS

Not only did the residential distribution of survey respondents closely resemble the distribution for the nation as a whole as reported in the census, the age and marital status distributions of women were also similar in the 1991 Census and the CRRHS ([Table II.1](#)). The proportion of women within each five-year age group was within two percentage points in the two sources, indicating that the CRRHS was representative of the national population with regard to age. Marital status distributions were also similar, although survey respondents were slightly more likely to be currently married and less likely to be never married/in union than were women enumerated in the 1991 Census.

Distributions of marital status, educational attainment, and religious affiliation at the time of interview for the CRRHS respondents are displayed in [Table II.2](#). Approximately two-thirds of respondents were married or living in a consensual union (i.e., living with a man, but not legally married) at the time, while one-fourth had never been married. The proportion of women married or in union stops rising after ages 25-29 at about 85%. By ages 30-34 almost one-tenth women were currently divorced or separated.

About four of every ten respondents had received a secondary school diploma. About one-fifth of secondary school graduates had attended university. The CRRHS results indicate that educational attainment has been increasing. The percentage who attended only primary school fell from 24% to 7% in a 20-year period, while the percentage who obtained a secondary school diploma rose from 41 % to over 50% in the 20-24 and 25-29 year-old cohorts ([Figure II.1](#)). (We have ignored 15-19 year-olds in this part of the analysis data because many respondents of that age were still in school at the time of interview.)

Almost two-thirds of respondents said they had no religious affiliation. Thirty-one percent of respondents said they were Catholic. The percentage saying they had no religion decreased steadily as age increased ([Figure II.2](#)). Among those under age 30 about seven of every ten women reported they belonged to no religion.

[Table II.3](#) shows that the largest group of women (40%) lived in households where the total monthly income was between 3,000 and 7,000 Crowns (about US\$100-250). Relatively few respondent households had a monthly income below 3,000 Crowns or above 15,000 Crowns. Sixty-one percent of respondents said they were currently working, while another 8% were reportedly on maternity leave. Among those who were employed, two-thirds worked between 40 and 44 hours per week.

The CRRHS questionnaire included questions on whether respondents' homes contained selected possessions, listed in the top panel of [Table II.4](#) and in [Figure II.3](#). The vast majority of homes contained a bathroom or shower (98%) and a color television (89%). In addition, most homes had an automatic washing machine (75%) and an automobile

Figure II.1
Education of Respondents,
by Age Group

Percent Distribution, 1993 CRRHS




Figure II.2
Religion of Respondents,
by Age Group

Percent Distribution, 1993 CRRHS





Figure II.3
Percent of Respondent Households
with Selected Possessions/Amenities
1993 CRRHS



(64%). Telephones were present in 33% of respondent households.

[Table II.5](#) reveals that, in spite of the recency of the conversion to a free market economy, 43% of respondents lived in residences owned by her or her family and only one-third still lived in residences rented from the state. Ten percent of respondents lived in residences which they shared with another family. Most of the households in which respondents lived contained three or four people.


[Table II.6](#), which presents distributions of several characteristics, controlling for respondent education, excludes women under 20 years of age, since many of them had not yet finished their education. Not surprisingly, there was a strong correlation between the education of respondents and their partners ([Table II.6](#)), but, it was not unusual for women to be married to men with slightly less education than themselves. There was a distinct association between education and household income, but even among those who attended university there were relatively few women in the highest income category (11%). Since income increased with educational attainment, it is not surprising that the number of selected possessions (color television, automatic washer, automobile, VCR, telephone, personal computer) rose with education as well. The proportion of respondents with five or six of these items increased from 6% for women with primary education to 23% for those with any university education.

It is clear from the results shown in [Table II.7](#) and in [Figure II.4](#) that age at marriage in the Czech Republic is typically quite young, with the median age at marriage (based on life table estimates) just 20.7 years. (Though not strictly comparable, the mean age at first marriage for Europe as a whole is 25 years and for Eastern Europe is 22 years (Population Reference Bureau, 1994b)). It also appears that marriages are highly concentrated between 18 and 25 years of age: only 6% of women married before 18, but only 12% were not married by age 25. It also is evident that the age at marriage has not changed in recent years, being almost identical for 15-24, 25-34, and 35-44 year-old respondents. Age at marriage was similar in Bohemia and Moravia and varied little according to the size of the place of residence. Median age at marriage increased with educational attainment, but even among those women who attended university, it was only 23.5 years.

[Table II.8](#) shows the percentages of first marriages that were still intact at the time of interview, controlling for the number of years that had passed since the date of marriage. Overall, 79% of marriages were still intact, ranging from 91% of those that began less than five years before interview to 70% of those that began at least 20 years before interview. Several variables appear to be related to the survival of marriages, even when duration is held constant. Marriages in Bohemia were less likely to survive than those in Moravia at all durations. Marriage survival was inversely correlated with size of place, with only 73% of marriages of women living in large cities still intact. Women who attended only primary school were less likely to have intact first marriages, regardless of years since marriage. Women affiliated with religions other than Roman Catholicism

Figure II.4
Percent Ever Married before Selected Ages
(Life Table Estimates)

1993 CRRHS



were the most likely to experience marital dissolution. Catholics who regularly attended religious services were the most likely to be in intact first marriages. Women who married young were much more likely than other women to no longer be in their first marriage, especially if the marriage took place before age 18.

TABLE II.1
Comparison of Percent Distributions of Age and Marital Status
according to the 1991 Czech Census and
the 1993 Czech Republic Reproductive Health Survey

	1991 Census	1993 CRRHS		Difference
Age/Marital Status	Percent	Percent	No. Women	
Age				
15-19	18.3	19.7	(662)	+1.4
20-24	14.9	15.7	(756)	+0.8
25-29	14.9	15.4	(828)	+0.5
30-34	15.5	13.8	(720)	-1.7
35-39	18.1	17.6	(805)	-0.5
40-44	18.5	17.7	(726)	-0.8
Marital Status				
Never Married	26.6	20.6	(925)	-2.6*
Consensual Union	*	3.4	(152)	
Married	64.9	68.0	(3065)	+3.1
Divorced/Separated	7.4	7.1	(313)	-0.3
Widowed	1.1	0.9	(42)	-0.2
Total	100.0	100.0	(4497)	

*In the 1991 census, individuals living in a consensual union were classified as never married. Therefore, in calculating the difference between the CRRHS and the 1991 census women who were never married and in consensual union were combined.

TABLE II.2
Selected Sociodemographic Characteristics of Respondents, by Age Group
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Total	Age Group					
		15-19	20-24	25-29	30-34	35-39	40-44
Marital Status							
Married	63.7	6.4	58.3	83.3	81.6	82.9	82.3
Consensual Union	3.1	2.6	5.2	2.5	3.4	2.5	2.5
Divorced/Separated	6.5	0.2	3.4	6.6	9.0	10.5	10.0
Widowed	1.0	0.0	0.0	0.1	1.4	1.7	2.9
Single	25.7	90.8	33.1	7.5	4.7	2.4	2.2
Education							
Primary	18.0	22.2	6.6	10.1	14.1	26.9	24.2
Secondary No Diploma	40.7	64.3	40.2	34.5	36.6	29.2	34.8
Secondary Diploma	33.8	12.7	44.3	44.5	37.4	34.9	34.6
Any University	7.6	0.9	8.9	10.9	11.8	8.9	6.4
Religion							
None	64.5	72.9	68.9	70.8	63.4	57.6	53.3
Catholic	31.2	23.9	26.1	25.8	32.2	37.3	41.5
Czech Evangelical	2.0	1.6	1.2	1.4	1.9	2.5	3.0
Czech Hus Church	0.6	0.0	0.8	0.2	1.1	1.1	0.6
Other	1.5	1.4	2.5	1.6	1.2	0.9	1.4
Not Stated	0.3	0.2	0.5	0.1	0.3	0.5	0.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number of Respondents	(4497)	(662)	(756)	(828)	(720)	(805)	(726)

TABLE II.3
Percent Distributions of Monthly Household Income, Employment Status
of Respondents and Average Number of Hours Worked per Week
1993 Czech Republic Reproductive Health Survey

Characteristics	Percent
Monthly Household Income (Crowns)*	
0-3,000	4.0
3,001-7,000	39.8
7,001-10,000	28.7
10,001-15,000	14.4
15,001 +	4.3
Don't Know	6.8
Not Stated	1.9
Whether Currently Employed	
Employed	60.7
Employed, on Maternity Leave	8.3
Not Employed	31.0
Average Hours Worked**	
Less than 40	16.7
40-44	68.4
45-49	5.0
50 or More	9.9
Total	100.0
Number of Respondents	(4,497)

* At the time of the CRRHS, \$1.00(US) equalled approximately 28 Crowns.

**Women currently working only (N=2,761).

TABLE II.4
Percent of Respondent Households with Various Possessions/Amenities
and Percent Distribution of Number of Selected Possessions* in Households
1993 Czech Republic Reproductive Health Survey

	Percent
Possessions/Amenities	
Bathroom/Shower	98.6
Color Television	89.3
Automatic Washing Machine	74.6
Automobile	63.8
VCR	34.6
Telephone	33.2
Weekend House (Have access to)	22.8
Weekend House (Own)	17.3
Local Heating	15.9
Personal Computer	9.7
Number of Selected Possessions*	
0-1	13.6
2	19.4
3	28.9
4	23.3
5-6	18.7
Total	100.0
Number of Respondents	(4,497)

*The selected possessions are: color television, automatic washing machine, automobile, VCR, telephone, and personal computer.

TABLE II.5
Ownership of Residence, Number of Families Living in Residence
and Number of People Living in Residence
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Percent
Ownership of Residence	
Owned by Respondent's Family	43.2
Cooperative Ownership	16.8
Rented from Private Owner	6.3
Rented from State	33.7
Number of Families in Residence	
1 Family	89.8
2 or More Families	10.2
Number of People in Residence	
1-2	12.4
3	28.0
4	42.8
5	12.4
6 or More	4.5
Total	100.0
Number of Respondents	(4,497)

TABLE II.6
Husband's Education, Household Income, and Number of Selected Possessions,
by Respondent's Education, Women 20 Years and Older
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Total	Respondent's Education			
		Primary	Secondary, No Diploma	Secondary, Diploma	Any University
Partner's Education					
Primary	10.6	32.3	8.0	5.0	0.6
Secondary, No Diploma	52.0	54.6	70.6	42.5	11.6
Secondary, Diploma	26.2	10.8	18.5	39.6	29.3
Any University	11.3	2.3	2.9	13.0	58.5
Household Income*					
0-3,000	4.0	8.3	4.4	2.3	1.5
3,001-7,000	41.7	52.7	46.3	36.9	24.0
7,001-10,000	28.6	24.8	30.3	31.7	26.5
10,001-15,000	15.0	6.0	11.2	18.6	31.0
15,001 +	4.3	1.8	2.5	5.3	11.0
Don't Know	3.3	3.3	2.9	3.2	5.3
Not Stated	2.2	3.1	2.4	2.1	0.8
Possessions**					
0-1	13.6	29.1	14.6	8.6	6.5
2	19.5	23.7	22.8	16.1	15.5
3	28.9	25.5	29.7	29.7	24.5
4	23.3	15.7	23.1	26.2	30.1
5-6	14.7	6.0	10.5	19.5	23.0
Total	100.0	100.0	100.0	100.0	100.0
Number of Cases	(3,835)	(605)	(1,347)	(1,511)	(372)

* Monthly income in Czech Crowns. \$1.00(US) equalled approximately 28 Crowns

**The selected possessions are: color television, automatic washing machine, automobile, VCR, telephone, and personal computer.

TABLE II.7
Life Table Estimates of Percent of Women Ever in Union Before Ages 18, 20, and 25
and Median Age at First Union, by Selected Characteristics
1993 Czech Republic Reproductive Health Survey

Characteristics	Percent in Union by Age:			Median Age at First Union
	18	20	25	
Age at Interview				
15-24	6.1	38.3	--	20.7
25-34	6.4	38.2	88.8	20.7
35-44	6.1	35.3	87.4	20.8
Region				
Bohemia	6.4	37.8	87.7	20.7
Moravia	5.8	35.7	89.1	20.8
Size of Place				
Less than 5,000	6.2	39.2	90.0	20.6
5,000-19,999	5.7	37.2	88.1	20.8
20,000 +	6.3	35.2	87.1	20.8
Education				
Primary	14.7	53.4	91.6	19.8
Secondary, No Diploma	7.1	45.2	92.1	20.3
Secondary, Diploma	2.8	29.1	88.6	21.0
Any University	2.4	7.4	66.7	23.5
Total	6.2	37.0	88.3	20.7

TABLE II.8
Percent of Women Ever in Union Still in First Marriage/Union
by Selected Characteristics, by Years since First Marriage
1993 Czech Republic Reproductive Health Survey

Characteristics	Years since First Marriage				Total
	0-4	5-9	10-19	20+	
Total	90.7	84.6	74.3	69.6	78.9
Region					
Bohemia	89.6	81.9	73.0	66.0	76.8
Moravia	92.5	88.5	76.4	75.8	82.1
Size of Place					
<5,000	93.1	91.5	83.1	79.3	85.9
5,000-19,999	93.1	85.1	74.7	73.3	80.4
20,000+	88.1	79.2	67.9	59.1	72.9
Education					
Primary	84.4	81.9	72.2	64.3	72.1
Secondary, No Diploma	88.8	87.0	72.7	72.7	79.0
Secondary, Diploma	93.0	82.3	76.4	70.2	80.7
Any University	93.0	89.3	77.5	*	84.5
Religion					
None	89.5	84.5	73.2	68.4	78.7
Catholic, Attends	93.1	95.4	80.2	80.0	85.0
Catholic Does Not Attend	94.4	86.3	77.2	71.8	80.0
Other	89.8	60.0	63.7	60.0	68.8
Age at First Marriage					
<18	90.0	79.3	55.9	60.0	67.9
18-19	90.2	84.5	72.2	67.2	76.8
20-21	92.0	85.9	76.7	73.4	80.8
22+	90.4	84.7	77.8	80.9	82.4

*Fewer than 25 women

III. CHILDBEARING/ABORTION

One of the major reasons for conducting the 1993 CRRHS was to learn more about trends, differentials, and patterns of childbearing in the Czech Republic. Vital statistics on births in the Czech Republic are virtually complete, so that measuring current fertility rates was not a primary objective of the survey, unlike similar surveys carried out in developing countries. However, the data collected in the survey provide much more detailed information on reproduction, as well as information on various aspects of childbearing that are not recorded by the vital statistics system. Each respondent was asked to give a history of all the pregnancies she had ever had, including information on the duration, and the outcome of each pregnancy, as well as the sex and survival of each child born alive. Respondents were asked about the intendedness of each pregnancy ending after 1987 and breastfeeding of children born after 1987. Finally, women were asked about the numbers of children they would like to have and the number they planned to have when they first married. (The survey results on health and medical aspects respondents' pregnancies and deliveries, as well as maternal health are presented in Chapter 5, "Pregnancy, Delivery, and Infant Health").

Children Ever Born

[Table III.1](#) presents percentage distributions of the number of live births to respondents according to age and years since the start of first union at the time of interview. Only 3% of respondents had given birth to four or more children. The largest number of live births for any respondent was seven, reported by only three (0.1%) women. Among women 35-39 years and 40-44 years, the vast majority of whom had completed their childbearing, there was strikingly little variation in family size. Fifty-seven percent have had two children, by far the most common number of children for these cohorts of women. Few of the oldest respondents have had very small families (3% with no children and 12% with one child among 40-44 year-olds) and, likewise, few have had large families (5% with four or more children). Similar figures are seen for those women who were first married (or in union) at least 10 years before interview. Among those women never in union, 5% reported having any live births. It is evident, as will be demonstrated in later tables, that childbearing tends to start soon after marriage (only 28% of women married less than five years were still childless) and stop at relatively early ages as well.

[Table III.2](#) presents mean numbers of children ever born (parity) according to selected characteristics, controlling for respondents' ages at the time of interview. The fact that the mean parity for women at ages 35-39 and 40-44, 2.1 and 2.2, respectively, are higher than the total fertility rate at the time of the survey (1.9, as will be shown below) indicates that fertility rates have fallen somewhat in recent years.

Differences in mean numbers of live births tended to be small among most population

subgroups examined. There was virtually no difference between Bohemia and Moravia, while there was a very slight inverse correlation between size of place and mean parity. A stronger inverse correlation was noted between educational attainment and parity, especially at younger ages. However, unlike in most other developed countries, where an inverse correlation exists between income and parity, no such relationship was observed in the Czech Republic. This may be the result of a situation which is only now starting to change, whereby salaries and social status have not been highly associated. The mean parity of Roman Catholics tended to be slightly higher than for women with no religious affiliation.


Current Fertility

The total fertility rate (TFR) for the three-year period preceding the CRRHS (1990-1992), based on data from the survey was 1.87 births per woman, identical to the rate computed from Czech Republic vital statistics ([Table III.3](#)). This confirms that, as in all but a few European countries, fertility rates have fallen below the replacement level of 2.1 births per woman (Althaus 1992). It also lends great confidence in the quality of the live birth data reported by women in the CRRHS. [Figure III.1](#) shows TFRs for many countries in Europe in about 1991. The rate for the Czech Republic, although low by world standards, is not atypical for Europe. For the former Czechoslovakia, the TFR rose between the late 1960s and the mid-1970s from about 2.0 children per woman to a level of 2.5 children per woman. Fertility then declined steadily, but very gradually, from the middle of the 1970s until the time of the survey (United Nations 1986-1990). (For a description of the substantial decline in Czech fertility over the past two centuries, see Fialova et al. (1990)). Throughout the former Soviet bloc fertility fell until the mid 1960s, after which trends diverged, depending predominantly on government policies and abortion restrictions (Festy 1991).

[Table III.3](#) also shows that age-specific fertility rates computed from survey data closely match those from Czech vital statistics. [Table III.3](#) reveals that childbearing in the Czech Republic is very heavily concentrated in the 20-24 and 25-29 year age groups, with very little childbearing taking place after ages 30-34. [Figures III.2](#) and [III.3](#) demonstrate the extent to which childbearing starts and ends early in the Czech Republic. These figures portray age-specific fertility rates for almost all European countries (excluding former Soviet republics) for 1990, the midpoint of the time period for which fertility was measured in the CRRHS data. It is readily apparent that the Czech Republic has unusually high fertility at ages 20-24 and unusually low fertility at ages 30-34. This is a pattern of childbearing exhibited by most eastern European populations in recent decades. Despite the social and economic changes experienced recently and the Czech Republic's nearness to and exposure to western Europe, there was still no indication of any changes in this pattern at the time of the survey.


[Table III.4](#) presents TFRs for various population subgroups for the period 1990-1992. Fertility was somewhat lower in larger cities (over 20,000 population) than in smaller

Figure III.1
Total Fertility Rates,
Selected Central and Eastern European Countries
1991*




* Source of Data: Population Reference Bureau, "1992 World Population Data Sheet".

Figure III.2
Fertility Rates, 20-24 Year-Old Women,
Selected European Countries,
1990*



* Source of Data: United Nations Population Division, "1991 Demographic Yearbook".

Figure III.3
Fertility Rates, 30-34 Year-Old Women,
by Selected European Countries,
1990*



* Source of Data: United Nations Population Division, "1991 Demographic Yearbook".

places, 1.7 and 2.0 children per woman, respectively. There was a clearly inverse relationship between education and fertility. Fertility was especially low among women who had attended a university, 1.5 births per woman. The fertility rate was higher among Catholics (2.1) than among those with no religious affiliation (1.8).

Timing of First Births

It has already been noted that marriage tends to occur early in the Czech Republic and that, by European standards, rates of childbearing at young ages are very high. [Table III.5](#) reveals that, even though the percentage of women with births before their eighteenth birthday was quite low (3%), the percentage who have given birth before their twenty-fifth birthday was extremely high (79%). The median age at first birth, based on life table estimates, was 22.0 years, virtually the same as the mean age at first birth reported in the 1977 Czechoslovak Fertility Survey (World Fertility Survey, 1978). The timing of first births varied little according to region or size of place of residence. As with age at marriage, there was no indication that the age at first birth had been increasing, unlike most other developed countries. In fact, the median age at first birth was lower and the percentages giving birth before selected ages were slightly higher for the younger cohorts than for the oldest cohort. However, the median age at first birth increased with education and was particularly high among women who attended a university (25.4 years). By age 25, just under half of these women had given birth, a far lower percentage than for other women.

[Table III.6](#) focuses on the timing of first births relative to the beginning of marriage or cohabitation. Only 4% of respondents reported giving birth before being in union, but by the time they had been in union for just one year, 54% reportedly had borne a child. Within five years of the date of first union, only 7% of women had not had a live birth. The median duration at first birth was only 0.9 years (about 11 months). Clearly, the societal norm in the Czech Republic has remained having a child within a short time of marrying. There was no sign that couples are waiting longer to have their first child; the median duration at first birth is similar for all cohorts. The well educated waited only slightly longer to have a child than other women. Thus, the fact that well educated women tend to have their first birth later in life than other women is primarily a function of the fact that they tend to marry later, and not that they delay childbearing once they are married.

Desired Number of Children

All respondents were asked about the number of children they currently desired to have, while ever-married respondents were also asked the number they had intended when they first got married. [Table III.7](#) shows that women had a very strong preference for a two-child family at the time of marriage. Two-thirds of all women ever in union and three fourths of those who had formed an idea of how many children they wanted, said they had desired two children. Very few women said that they had wanted to remain

childless (1%) or have four or more children (3%). There was an overwhelming preference for two children in every subgroup of the population examined. There was relatively little variation in desired family size at marriage according to place of residence, age, years since marriage, or education. Religious Catholics (i.e., those attending mass at least monthly) were more likely to express a desire for a larger family, with about one-third saying they wanted at least three children.

[Table III.8](#), which consists of a cross-tabulation of the number of children planned at the time of marriage by the number currently planned, shows that the preference for the two-child family remains strong after marriage and that there is much changing of preferences and/or failure to achieve desired family size. Most women who revised their preferences reported they now wanted to have two children. None of the women who said they had wanted to remain childless at marriage still wished to do so when interviewed and almost half wanted to have two children. Forty percent of those originally wanting one child said they now desire to have two. One-third of those originally wanting three children wanted two at the time of interview.


According to [Table III.9](#), of all fecund respondents who were currently in union, 69% desired to have no more children. Among those with two living children the figure was 86% and among those with at least three children it rose to 94%. Somewhat surprisingly, except among childless women and those with at least three living children, the proportion of women wanting no more children declined as education increased. This relationship did not exist between desire for more children and household income, however. Catholics who attended church regularly were less likely than others to want to have no more children. Women using modern contraceptive methods were the most likely to want no more children. However, large percentages of women using no contraception also wanted no more children, an indication of the existence of unmet need for family planning.

The figures in [Table III.10](#) and [Figure III.4](#) reconfirm the popularity of the two-child family in the Czech Republic. Sixty-seven percent of women with no living children would have liked to have two children, 49% of those with one child wanted one more, and 86% of those with two children wanted no more. There appears to be considerable acceptance of the one-child family, since 30% of women with one child wanted no more, even though few childless women said they only wanted one. There is almost no desire to have large families, with very few women indicating they want to have three or more children. The majority of this small group consisted of women with no children yet, many of whom will, no doubt, reduce their expectations as they get older.

Planning Status of Pregnancies

For every pregnancy ending since January 1988, respondents were asked, first, whether they intended to get pregnant and, if the pregnancy was unintended, whether they had intended to have no more children or had intended to wait longer to have them. If the

Figure III.4
 Number of Additional Children Desired
 According to Number of Living Children
 Fecund Women, Percent Distribution, 1993 CRRHS



respondent wanted to have no more children, the pregnancy was termed "unwanted". If she wanted to wait longer to become pregnant, it was termed "mistimed". If she reported that she did want to become pregnant when she did, the pregnancy was considered "planned". (David et al. (1988) discuss some of the disadvantages encountered for many Czech children who were unplanned in their research on births to women who were denied abortions.)

[Table III.11](#) presents percentage distributions of planning status for all reported pregnancies according to selected characteristics. The figures presented in [Table III.11](#), however, almost certainly represent a substantial underestimate of unplanned, and particularly, unwanted pregnancies in the Czech Republic. As is discussed later in this chapter, we estimated that about half of induced abortions (and the pregnancies leading to them) were not reported by respondents. We can safely assume that all but a small proportion of these pregnancies were unplanned and that a large majority of them were unwanted. After adjustments were made for abortion underreporting we estimated that only about 51% of recent pregnancies were planned, 16% were mistimed, and 28% were unwanted, with the remainder of unspecified status.

Despite the underreporting of unplanned pregnancies, the results in [Table III.11](#) are useful for examining relative levels of unintendedness among various population subgroups. As size of place increased, so did the percentage of pregnancies that were reportedly unwanted. The proportion of pregnancies that were unwanted rose sharply from ages 25-29 and once women had at least two living children. Most mistimed pregnancies occurred relatively early: before ages 25-29, i.e., before many women/couples have attained their desired family size. The percent planned was lowest and the percent unwanted was highest among women with only a primary school education. There was relatively little difference in planning status according to religion, except that Catholics who regularly attended services reported somewhat fewer mistimed and unwanted pregnancies than other women did.

[Table III.12](#) shows distributions of pregnancy outcomes according to the reported planning status of pregnancies. Again because of underreported induced abortion, the figures in this table relating to the three categories of unplanned pregnancy (mistimed, unwanted, and unspecified) are not completely reliable. Of planned pregnancies, 82% resulted in a live birth. In addition, most of the 6% of pregnancies that were current would also end with a live birth. After correction for underreporting, it appears that slightly more than half of mistimed pregnancies were aborted. Unwanted births are relatively rare events in the Czech Republic. After adjusting for unreported abortions, we estimated that only about 4% of unwanted pregnancies resulted in a live birth.

Induced Abortions

Much of central and eastern Europe in recent decades has been characterized by very high incidence of induced abortion (Blayo 1991; David 1992; Ketting 1992). Abortion

has been widely used in most of these countries as a primary or backup method of family planning. [Table III.13](#) consists of a list of recent induced abortion indicators, based on official statistics, for most of the countries in that part of the world, taken from Blayo (1993). According to these statistics, except for Albania, all other countries recorded at least 38 percent of known pregnancies ending in induced abortion. The rates for the Czech Republic (1.56 lifetime abortions per woman and 46% of pregnancies ending in abortion, as well as 42 abortions per 1,000 15-44 year-old women and 84 abortions per 100 live births in 1991) were lower than, in many of the former Soviet bloc countries of eastern Europe, but were likewise much higher than the rates found in the rest of Europe. Since Czech medical statistics are thought to provide virtually complete counts of induced abortion, the CRRHS was not intended as a tool to measure the incidence of abortion. The completeness of Czech abortion statistics allowed us to estimate the completeness of abortion reporting in the CRRHS and to adjust the survey estimates.

As has been found in most surveys that have addressed the subject, it was apparent that the number of induced abortions was substantially underreported in the 1993 CRRHS. Based on comparisons with Czech medical statistics, it was estimated that CRRHS respondents only reported between 45 and 50 percent of the abortions they underwent.

Finally, women who reported having any abortions since the beginning of 1988 were asked to give the most important reason that they decided to have an abortion, rather than continue with the pregnancy. [Table III.14](#) shows that over one-third of abortions occurred primarily because women did not want to have any more children. Fifteen percent were because women wanted to wait longer to have another child. Fourteen percent were said to be the result of the inability to afford another child. Less commonly stated reasons were: childbearing was risky for the woman (10%), her partner wanted her to have the pregnancy terminated (5%), her relationship had ended (3%), and diagnosis or fear of a fetal defect (3%). It is unknown whether these percentages would change significantly if survey respondents had reported all induced abortions.

TABLE III.1
Total Number of Live Births, by Age and Years since Marriage*
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Age Group/ Years since 1st Marriage	Live Births					Total	No. of Women
	0	1	2	3	4 +		
Age Group							
15-19	94.3	5.5	0.2	0.0	0.0	100.0	(662)
20-24	47.3	36.8	14.4	1.2	0.4	100.0	(756)
25-29	12.0	32.9	46.2	8.2	0.7	100.0	(828)
30-34	6.5	18.8	55.1	15.1	4.6	100.0	(720)
35-39	4.5	14.7	56.7	18.1	6.0	100.0	(805)
40-44	3.0	11.5	56.8	23.4	5.2	100.0	(726)
Years since Marriage							
Never Married	95.4	4.4	0.2	0.0	0.0	100.0	(899)
Less than 5	27.9	53.6	17.3	1.2	0.0	100.0	(783)
5-9	5.7	29.2	54.7	9.3	1.1	100.0	(802)
10-14	3.6	14.9	60.8	15.9	4.8	100.0	(762)
15-19	2.0	12.1	60.0	19.7	6.2	100.0	(702)
20 +	0.5	10.1	55.3	27.4	6.7	100.0	(549)
Total	30.1	19.2	37.1	10.9	2.8	100.0	(4497)

*In this and all subsequent tables, the term "marriage" includes consensual unions and "married" respondents include those living in consensual unions.

TABLE III.2
Mean Number of Children Ever Born,
by Age and Selected Characteristics
1993 Czech Republic Reproductive Health Survey

Characteristics	Current Age						Total
	15-19	20-24	25-29	30-34	35-39	40-44	
Total	0.1	0.7	1.5	1.9	2.1	2.2	1.4
Region							
Bohemia	0.1	0.7	1.5	1.9	2.0	2.1	1.4
Moravia	0.1	0.7	1.6	2.0	2.2	2.3	1.4
Size of Place							
<5,000	0.1	0.8	1.7	2.0	2.2	2.3	1.5
5,000-19,999	0.0	0.7	1.5	2.0	2.1	2.1	1.4
20,000+	0.1	0.7	1.4	1.8	2.0	2.1	1.3
Marital Status							
Ever Married	0.5	1.0	1.6	2.0	2.1	2.2	1.8
Never Married	0.0	0.1	0.3	0.3	*	*	0.1
Education							
Primary	0.1	1.4	1.9	2.2	2.4	2.4	1.7
Sec, No Diploma	0.1	0.9	1.7	2.1	2.0	2.2	1.2
Sec, Diploma	0.1	0.6	1.4	1.8	1.9	2.1	1.4
Any Universty	*	0.1	1.0	1.5	1.9	1.7	1.2
Religion							
None	0.1	0.7	1.5	1.9	2.0	2.1	1.3
Catholic	0.1	0.7	1.6	2.1	2.3	2.3	1.6
Other	0.0	0.6	1.5	1.8	1.9	2.5	1.5
Household Income							
0-7000	0.1	0.9	1.6	2.0	2.1	2.2	1.5
7001-10000	0.0	0.6	1.5	2.0	2.1	2.2	1.5
10001+	0.0	0.4	1.4	1.9	2.1	2.2	1.4
Not Stated	0.0	0.3	1.4	1.6	2.1	2.2	0.7

*Fewer than 25 respondents in category

TABLE III.3
Age-Specific Fertility Rates and Total Fertility Rates (TFR)
1993 Czech Republic Reproductive Health Survey and
1991 Vital Statistics for the Czech Republic

Age	Age-Specific Fertility Rates	
	CRRHS *	Vital Statistics
15-19	49	45
20-24	176	174
25-29	92	105
30-34	41	38
35-39	11	11
40-44	4	2
Total Fertility Rate	1.87	1.87

* Calculated for the three-year period, 1990-1992.

TABLE III.4
Total Fertility Rates by Selected Characteristics
1993 Czech Republic Reproductive Health Survey

Characteristics	Total Fertility Rate*
Total	1.87
Region	
Bohemia	1.92
Moravia	1.79
Size of Place	
< 5,000	1.98
5,000-19,999	2.01
20,000 +	1.72
Education	
Primary Only	2.09
Secondary, No Diploma	1.88
Secondary, Diploma	1.84
Any University	1.47
Religion	
None	1.77
Catholic	2.11

*Calculated for three-year period, 1990-1992.

TABLE III.5
Life Table Estimates of Percent of Respondents with Any Live Births before Selected Ages
and Median Age at First Birth, according to Selected Characteristics
1993 Czech Republic Reproductive Health Survey

Characteristics	Percent with a Live Birth Before Age:			Median Age at First Birth
	18	20	25	
Total	3.3	22.4	78.6	22.0
Region				
Bohemia	3.3	22.7	77.3	21.9
Moravia	3.4	22.0	80.5	22.1
Size of Place				
Less than 5,000	3.6	23.6	81.9	21.7
5,000-19,999	3.2	23.1	78.7	22.0
20,000 +	3.2	21.3	76.0	22.2
Current Age				
15-24	3.3	24.3	NA	21.9
25-34	3.7	23.4	79.0	21.8
35-44	3.0	20.7	78.0	22.2
Education				
Primary	11.3	39.9	81.2	20.8
Secondary, No Diploma	2.8	29.5	86.4	21.2
Secondary, Diploma	1.2	13.6	76.3	22.5
Any University	0.0	2.2	47.3	25.4
Religion				
No Religion	3.1	22.1	78.6	22.0
Catholic, Attend	1.8	16.5	69.7	22.7
Catholic, Not Attend	4.7	25.0	80.5	21.8
Other	1.0	20.1	74.7	22.3

TABLE III.6
Life Table Estimates of Percent of Respondents with Any Live Births before Selected Marriage Durations*
and Median Duration at First Birth (in Years), according to Current Age and Education of Respondent
1993 Czech Republic Reproductive Health Survey

Characteristics	Years since Beginning of First Marriage					Median Duration
	Before Union	1	2	5	10	
Total	4.2	54.4	75.6	92.9	96.5	0.9
Current Age						
15-24	1.6	52.0	73.7	90.1	---	1.0
25-34	5.2	57.3	77.0	92.8	96.2	0.9
35-44	4.3	53.0	75.0	93.2	96.7	0.9
Education						
Primary Only	7.4	59.6	78.6	91.4	95.0	0.8
Secondary, No Diploma	3.7	60.0	80.1	93.9	97.1	0.8
Secondary, Diploma	3.3	49.4	72.4	93.6	97.1	1.0
Any University	3.7	42.0	63.7	89.6	94.4	1.4

*Years since beginning of first marriage or cohabitation

TABLE III.7
Number of Children Planned When Respondent First Married
according to Selected Characteristics, Women Ever Married
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Number of Children Planned						Total	(N)
	0	1	2	3	4+	Unsure		
Total	1.0	8.2	67.3	9.5	3.1	11.0	100.0	(3,598)
Region								
Bohemia	1.2	9.0	67.5	8.1	2.8	11.5	100.0	(2,206)
Moravia	0.6	6.9	67.0	11.6	3.7	10.2	100.0	(1,392)
Size of Place								
Less than 5,000	0.5	6.5	66.4	11.0	2.3	13.3	100.0	(1,181)
5,000-19,999	0.8	8.0	67.8	10.5	2.7	10.2	100.0	(784)
20,000+	1.4	9.5	67.7	7.8	4.0	9.7	100.0	(1,633)
Age								
15-19	0.0	8.9	67.3	9.9	1.0	12.9	100.0	(90)
20-24	1.7	9.0	73.6	6.6	1.7	7.3	100.0	(552)
25-29	1.2	8.1	70.0	9.7	2.1	9.0	100.0	(776)
30-34	1.3	9.0	61.7	10.0	3.0	15.1	100.0	(690)
35-39	0.5	9.1	63.8	10.5	4.2	12.0	100.0	(782)
40-44	0.6	6.1	68.8	9.6	4.2	10.8	100.0	(708)
Years since First Marr.								
Less than 5	1.8	10.5	69.9	7.9	2.0	7.9	100.0	(783)
5-9	1.2	8.9	68.0	8.1	3.0	10.8	100.0	(802)
10-14	1.4	7.1	66.5	10.4	1.9	12.7	100.0	(762)
15-19	0.1	8.1	65.5	11.1	3.7	11.5	100.0	(702)
20+	0.3	6.0	60.5	9.8	5.2	12.3	100.0	(549)
Education								
Primary	1.1	8.9	59.2	11.7	3.5	15.7	100.0	(597)
Secondary, No Dipl.	0.6	7.9	69.0	8.6	2.4	11.5	100.0	(1,299)
Secondary, Diploma	1.2	8.2	70.0	8.4	3.4	8.8	100.0	(1,390)
Any University	0.9	7.3	64.7	13.7	4.3	9.1	100.0	(312)
Religion								
None	1.0	9.9	68.3	7.9	2.5	10.4	100.0	(2,296)
Catholic, Attend	0.0	2.4	50.2	21.1	8.9	17.4	100.0	(181)
Catholic, Not Attend	0.9	5.7	68.8	9.8	3.1	11.5	100.0	(965)
Other	1.8	5.4	64.9	14.3	5.4	8.3	100.0	(144)

TABLE III.8
Number of Children Planned When Respondent First Married
by Total Number of Children Currently Planned, Fecund Married Respondents
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Children Planned Currently	Children Planned at First Marriage					
	0	1	2	3	4 +	Unsure
0	0.0	0.6	0.2	0.9	0.0	0.7
1	24.3	39.3	5.3	3.2	4.3	9.4
2	48.6	39.9	72.4	33.5	27.0	42.5
3	10.8	7.5	13.4	45.4	35.7	22.7
4 +	0.0	1.0	1.8	6.9	27.8	10.4
Not Sure	16.2	11.7	6.8	10.1	5.2	14.3
Total	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of Women</i>	<i>(36)</i>	<i>(282)</i>	<i>(2283)</i>	<i>(310)</i>	<i>(100)</i>	<i>(359)</i>

TABLE III.9
Percent of Respondents Reportedly Wanting to Have No More Children
according to Selected Characteristics, Fecund Married Women*, 15-44 Years of Age
1993 Czech Republic Reproductive Health Survey

Characteristics	Total	Living Children**			
		0	1	2	3+
Total	69.3	6.4	27.8	85.7	93.8
Region					
Bohemia	68.6	7.5	26.6	85.9	93.5
Moravia	70.3	4.6	29.7	85.4	94.2
Education					
Primary	84.5	***	41.1	93.4	93.7
Secondary, No Diploma	70.1	6.8	25.7	85.9	94.8
Secondary, Diploma	65.7	4.2	28.6	84.2	94.2
Any University	53.5	7.4	20.8	79.0	***
Religion					
No Religion	67.4	4.9	28.0	86.1	95.9
Catholic, Attend	73.0	***	25.0	67.6	84.1
Catholic, Not Attend	75.1	14.8	28.4	87.7	94.8
Other	68.2	***	***	83.1	87.9
Household Income (Crowns/Month)					
0-7,000	63.3	4.2	18.1	83.5	92.4
7,001-10,000	73.6	10.2	33.9	88.4	92.9
10,001 +	75.4	3.1	42.6	86.1	98.0
Current Contraception					
Modern Method	76.0	10.3	35.4	87.9	95.7
Traditional Method****	69.6	2.5	28.8	85.9	93.9
No Method	54.9	5.4	17.4	79.8	87.1

*Women who have been contraceptively sterilized are considered to want no more children.

**Women who were pregnant at the time of interview are classified as having one more living child than the actual number when interviewed.

***Fewer than 25 respondents in category

****Withdrawal and natural family planning

TABLE III. 10
Number of Additional Children Desired, according to Current Number of Living Children
Fecund Women 15 to 44 Years of Age
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Additional Children Desired	Total	Number of Living Children			
		0	1	2	3+
0	52.7	5.2	30.4	86.3	94.0
1	13.8	6.9	48.6	5.3	1.6
2	20.9	66.6	6.8	0.4	0.0
3+	4.1	13.1	1.1	0.1	0.9
Not Sure	8.5	8.1	13.1	7.9	3.5
Total	100.0	100.0	100.0	100.0	100.0
Number of Women	(4,258)	(1,030)	(931)	(1,788)	(509)

*Women who have been contraceptively sterilized are considered to want no more children.

**Women who were pregnant at the time of interview are classified as having one more living child than the actual number when interviewed.

TABLE III.11
Planning Status of Pregnancies Ending after 1987, All Women 15-44 Years of Age
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Planning Status					Total	No. of Women
	Planned	Unplanned			Unsure		
		Mistimed	Unwanted	Unsure			
Total	65.9	13.1	16.2	1.1	3.6	100.0	(2,591)
Region							
Bohemia	64.6	13.4	17.1	1.3	3.8	100.0	(1,604)
Moravia	68.2	12.7	14.9	1.0	3.3	100.0	(987)
Size of Place							
Less than 5,000	73.8	11.0	11.3	0.7	3.1	100.0	(837)
5,000-19,999	68.9	12.4	15.0	1.2	2.6	100.0	(555)
20,000+	59.3	14.9	20.1	1.4	4.3	100.0	(1,199)
Pregnancy Outcome							
Live Birth	83.9	10.0	2.2	0.5	3.5	100.0	(1692)
Miscarriage/Stillbirth	81.8	5.6	7.1	0.5	5.1	100.0	(187)
Induced Abortion	9.4	24.4	59.8	3.1	3.3	100.0	(588)
Current Pregnancy	79.4	11.5	3.8	1.5	3.8	100.0	(124)
Age at End of Pregnancy							
15-19	60.1	28.2	3.7	1.1	6.9	100.0	(318)
20-24	74.3	16.2	6.0	0.7	2.8	100.0	(1,105)
25-29	68.1	8.1	19.4	1.5	2.9	100.0	(676)
30-34	57.5	5.6	32.6	1.5	2.9	100.0	(322)
35-39	37.2	0.0	57.0	1.7	4.1	100.0	(141)
40-44	36.8	0.0	50.0	2.6	10.5	100.0	(29)
Living Children*							
0	77.0	16.3	1.5	1.0	4.3	100.0	(677)
1	79.7	13.1	4.1	1.0	2.2	100.0	(1,114)
2	39.5	11.7	43.4	2.1	3.3	100.0	(604)
3	33.9	7.6	51.5	0.0	7.0	100.0	(148)
4+	38.9	5.6	40.7	0.0	14.8	100.0	(48)
Education							
Primary	56.8	10.1	23.6	0.3	9.2	100.0	(325)
Secondary, No Diploma	66.0	13.1	16.7	1.1	3.1	100.0	(1,008)
Secondary, Diploma	69.9	12.7	13.7	1.1	2.6	100.0	(1,014)
Any University	61.6	19.2	14.8	2.4	2.0	100.0	(244)
Religion							
None	64.7	14.5	16.9	0.8	3.1	100.0	(1,744)
Catholic, Attend	78.3	7.0	10.1	0.8	3.9	100.0	(121)
Catholic, Not Attend	67.7	9.9	16.1	1.9	4.3	100.0	(615)
Other	61.4	15.8	14.0	2.6	6.1	100.0	(106)

*Number of living children at the time the pregnancy occurred.

TABLE III. 12
Outcomes of Reported Pregnancies Ending after 1987
according to Planning Status of Pregnancy
Women 15 to 44 Years of Age
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Outcome of Pregnancy	Total	Planning Status of Pregnancy			
		Planned	Mistimed	Unwanted	Unplanned (Unspecified)
Live Birth	64.3	81.8	48.9	8.8	25.8
Miscarriage/Stillbirth	7.3	9.1	3.1	3.2	3.2
Induced Abortion	23.6	3.4	43.8	86.8	64.5
Current Pregnancy	4.8	5.8	4.2	1.1	6.5
Total	100.0	100.0	100.0	100.0	100.0
<i>Number of Pregnancies</i>	<i>(2,591)</i>	<i>(1,739)</i>	<i>(339)</i>	<i>(395)</i>	<i>(30)</i>
					<i>(88)</i>

TABLE III.13
Mean Number of Induced Abortions per Woman
and Induced Abortions per 100 Known Conceptions,
Based on Recent Official Statistics
Selected Countries of Central and Eastern Europe

Country	Year	Mean Abortions per Woman	Abortions per 100 Conceptions
Albania	1990	0.64	17
Belarus	1989	3.32	62
Bulgaria	1989	1.79	50
CZECH REPUBLIC	1990	1.56	46
Hungary	1990	1.24	40
Poland*	1989	2.22	52
Romania	1990	5.70	73
Russia	1989	3.97	66
Slovakia	1990	1.23	38
Ukraine	1989	2.96	61

*Since that time, legal access to abortion has been severely restricted.

Source of data: Blayo, 1993.

TABLE III.14
Reasons for Choosing to Have an Induced Abortion
Abortions Taking Place to Respondents after 1987
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Reason for Abortion	Percent
Wanted No More Children	34.2
Wanted to Wait Longer for Child	15.3
Could Not Afford Another Child	14.1
Childbearing Risky for Respondent	9.5
Partner Wanted Her to Have Abortion	4.5
Relationship Had Ended	3.3
Fetal Defect	2.8
Parents Wanted Her to Have Abortion	1.4
Was Not Married	1.3
Other Reasons	12.7
Reason Not Stated	0.9
Total	100.0
<i>Number of Abortions</i>	<i>(588)</i>

IV. FAMILY PLANNING

One of the primary objectives of the 1994 CRRHS was to learn more about the use of both modern and traditional methods of preventing pregnancy. Many of the countries of central and eastern Europe are known to have relatively low rates of childbearing, despite modest levels of use of modern contraception, largely through widespread reliance on induced abortion (Blayo 1991; David 1992). The 1993 CRRHS was designed to provide detailed information on women's knowledge and use of contraceptive methods, factors involved in method selection, reasons for nonuse of modern methods by women/couples wishing to avoid pregnancy, unmet need for family planning services, interest in surgical sterilization, and related topics.

Knowledge of Methods for Preventing Pregnancy

Each respondent was asked whether she was familiar with each of nine contraceptive methods. Women were not asked about methods generally unavailable in the Czech Republic, such as Norplant and injectable contraceptives. It is clear that knowledge of pregnancy prevention methods is not an important obstacle to use of contraception. As shown in [Table IV.1](#), among women in union, reported knowledge of every method asked about was widespread. Knowledge was nearly universal for condoms and oral contraceptives and was only slightly lower for the IUD, withdrawal, and natural family planning (variously known as the fertile days method, periodic abstinence, or the rhythm method). Even the least well known method, vasectomy, was familiar to about two-thirds of respondents.

For each of the nine methods, knowledge tended to increase with respondents' level of education. Among women who had attended university, knowledge was over 90 percent for every method. There were no substantial differences in knowledge of contraception according to other socioeconomic, demographic, and residential characteristics examined, except that respondents under the age of 20 were considerably less likely than others to know about each of the nine methods.

Ever Use of Contraception

Among women currently married or living with a partner, 92 percent had used contraception at some time during their lives ([Table IV.2](#)). When the analysis is limited to modern methods (i.e., withdrawal and natural methods are excluded), this percentage drops to 79 percent. Among modern methods, over half of these women had had sex with a partner who used condoms and about one-third had ever used an IUD or oral contraceptives. Other modern methods were rarely used. Over 70 percent of women in union had ever employed withdrawal with a partner to avoid pregnancy. One-third reported ever using any of the natural methods.

The percentage of women in union ever using any family planning method was virtually uncorrelated with respondents' ages. The proportion ever using a modern method, however, increases to ages 25-29 and then levels off. Ever use of the IUD, oral contraceptives, and sterilization, methods requiring a physician, were most strongly correlated with age. There was a strong, direct correlation between respondent education and ever use of contraception, such that 96 percent of women with a secondary school diploma and 98 percent of those who attended university had ever used a method. The only methods for which ever use did not increase with education were the IUD and sterilization.


Current Use of Contraception

Sixty-nine percent of women who were married or living with a man were using a family planning method at the time of interview ([Table IV.3](#)), a contraceptive prevalence rate slightly below the levels found in most other developed countries. There was relatively little difference in contraceptive prevalence between regions of the Czech Republic. The one region to stand out was South Bohemia, where only 55% of women in union reportedly were using a contraceptive method, 14 percentage points below the national figure. Couples with two or three children were the most likely to be using contraception (75% and 78%, respectively), while only 29% of childless couples were contracepting. This is another indicator that most couples still want to have a child as soon as they marry. As with ever use, current contraceptive use increased with level of education, at least up to the level of secondary school completion. Women not affiliated with an organized religion were somewhat more likely to be using than women who belonging to a religion. Somewhat surprisingly, there was virtually no difference in contraceptive prevalence between Catholics who regularly attended church and those who did not (both 64%). Women affiliated with religions other than Catholicism (mainly Protestants of assorted denominations) were the least likely to report that they were practicing contraception (59%).

Overall, the most commonly used contraceptive methods were reported to be withdrawal (19% of all respondents), condoms (15%), and the IUD (12%), which together accounted for just over four-fifths of all contraception ([Table IV.4](#)). Oral contraceptives were the fourth most common method, used by about 8% of all women. No other methods were currently employed by more than 2% of women. About 2% of women had been contraceptively sterilized. No respondents reported that their partner had undergone a vasectomy.

Because most contraceptors were women currently in union, the mix of methods used among them is similar to the population as a whole ([Figure IV.1](#)). Although only 46% of previously married women were using contraception, those who were using were more likely to employ an IUD or oral contraceptives than were currently married users. Among women who had never been married, 38% were using some method of pregnancy prevention. However, within this group over three-fourths of users relied on

Figure IV.1
**Current Contraceptive Use
 among Married Women**
 Percent Distribution, 1993 CRRHS



condoms, withdrawal, or both.

Withdrawal was the most commonly used method regardless of the number of living children, except among those with four or more children, among whom contraceptive sterilization was most widely used (27%) ([Table IV.5](#)). Relative reliance on modern methods grew as number of children increased ([Figure IV.2](#)). Condom use was slightly less common than withdrawal regardless of the number of children. IUD use was most prevalent among women with at least two children, among whom it was the second most widely used method. Oral contraceptive use was highest among women with between one and three living children. The relationships between age and contraceptive use ([Table IV.6](#)) appear to be, to a large extent, a function of the number of living children, rather than just age itself. The percentage of couples using withdrawal or condoms is relatively constant across all ages. The IUD, a method used primarily to limit childbearing in the Czech Republic, was relatively flat from ages 25-29 to 40-44. Oral contraceptive use is highest at ages 20 to 34, after which it declined sharply.

The only strong relationship apparent between contraceptive use and education was that condom use increased with educational level ([Table IV.7](#)). The highest proportion of sterilized women (5%) was among those with no secondary school education. The use of natural family planning methods appears to increase somewhat with increasing education.


Most of the difference in contraceptive prevalence between women espousing no religion and others was accounted for by much higher use of oral contraceptives among the former ([Table IV.8](#)). The level of use of withdrawal and condoms was almost indistinguishable between groups. Not surprisingly, the use of natural methods (NFP) was most widespread among Catholic women who regularly attended church, but even among this group only 5% of all women and 7% of contraceptors were utilizing NFP.

[Table IV.9](#), which is restricted to respondents who were fecund, in union, were not pregnant and did not want to become pregnant at the time of interview, compares contraceptive use between those who wanted no more children (limiters) and those who wanted to delay their next pregnancy (spacers). Overall contraceptive prevalence in these two groups was virtually identical (82% and 81%, respectively). Limiters were considerably more likely to be using an IUD (20% versus 9%) or female sterilization (4% versus 0%). Reported levels of use of oral contraceptives, withdrawal, and condoms were slightly higher among spacers than limiters.

Source of Contraception

There is no national family planning program in the Czech Republic, nor does the government distribute contraceptive methods. Oral contraceptives must be prescribed by a gynecologist, while IUDs must also be prescribed and inserted by gynecologists. These methods have, until recently, been available almost free of charge. With the

Figure IV.2
**Current Contraceptive Use among Married Women,
 by Number of Living Children**
 1993 CRRHS



changes underway in health care financing, the costs of these methods to consumers are likely to increase (Visser, Uzel et al. 1993). Condoms are available from pharmacies and other commercial outlets.

[Table IV.10](#) displays distributions of reported sources of oral contraceptives and IUDs according to the respondents' region of residence. Sixty-one percent of pill users obtained them from regional or district physicians, with 17% getting their supplies from pharmacies (information was not available on the physicians from whom these women received their prescriptions), 11% using polyclinic physicians, and 8% using private physicians. The percentage using regional or district physicians ranged from 80% in Prague to 41% in Central Bohemia.

Most women currently using an IUD had the device inserted at either a regional or district clinic (47%) or at a hospital (35%). Only 2% obtained their IUD from a private physician, a much lower percentage than for oral contraceptives. The proportion using regional/district physicians ranged from 70% in Prague down to only 25% in South Moravia. South Moravia had the highest proportion of IUD insertions at hospitals (56%) and Prague had the lowest (8%).

Reasons for not Using Contraception

Respondents who were not using a family planning method at the time of interview were asked to state their most important reason for not employing a method. About two-thirds of these women gave reasons that related to either a lack of current sexual activity (38%) or their pregnancy status (i.e., subfecund, trying to become pregnant, or currently pregnant) ([Table IV. 11](#)). Among the broad assortment of other reasons given, the most common was fear of health effects related to contraception (7%). About 11 % of women said they were not using because they thought it was difficult for them to get pregnant, because they were having sex only sporadically, or because they were postpartum or breastfeeding, indicating they felt that contraception was not very necessary. Religion and inability to obtain methods were rarely mentioned; even among Roman Catholics, fewer than 1 % said that religion was their major reason for non-use. Based on the results in this table, it appears that almost one-third of Czech women who were not using contraception may have been at some risk of unintended pregnancy due to non-use of contraception.

Not surprisingly, a lack of sexual activity was the principal reason for not using among those not currently married. Among those in union, just over half of nonusers reported a reason related to pregnancy status or sexual activity. Reasons such as fear of health effects, difficulty in getting pregnant, and postpartum/breastfeeding were much more common among these women than among those not in union.

Contraceptive Failure/Discontinuation

The 1993 CRRHS contained a contraceptive use/pregnancy calendar which tracked women's contraceptive histories from the beginning of 1988 to the date of interview. This allowed estimates to be made of yearly failure and discontinuation rates for each commonly used contraceptive method in the Czech Republic. Because of substantial underreporting of induced abortions in the survey (See Chapter III) we assume that failure rates and, to a lesser degree, discontinuation rates calculated from the reported histories of respondents are too low. It appears that reporting of pregnancies not resulting in abortion was virtually complete, but that about half of the 40-45 percent of pregnancies ending in abortion were omitted from the pregnancy histories. Therefore, we estimate that overall contraceptive failures are underestimated by about 20 to 25 percent. There is no obvious reason to expect a correlation between the method used at the time of failure and whether the failure was reported, so we assume that the failure rate for each method is on the order of 25 percent higher than presented in [Table IV.12](#).

The 12 month failure rates presented in [Table IV.12](#), especially when adjusted for underreporting of abortions, are very much in line with rates typically reported for these methods, both worldwide (Trussell and Kost 1987) and in eastern Europe (Romania Institute for Mother and Child Care 1995). After adjusting for underreporting, we estimate pregnancy rates within one year of about 3% of pill users, 4% of IUD users, 7% of condom users, 15% of withdrawal users, and 26% of NFP users. [Table IV.12](#) also displays unadjusted rates after 24 and 36 months.

[Table IV.13](#) presents 12, 24, and 36 month contraceptive discontinuation rates. These figures represent the probability that a couple beginning use of a particular method would no longer be using that method after a given duration, regardless of the reason for stopping. These rates should only be slightly affected by any underreporting of abortion, since discontinuation occurs for many reasons other than unintended pregnancy. Of the five methods examined, discontinuation of the IUD was far lower than for the others, with only 13% stopping use in the first 12 months. Even after three years, three-fourths of IUD users continued using. Rates of discontinuation for the other four methods in common use were extremely similar to each other.

Although the percentages in [Table IV. 14](#) cannot be used to estimate rates of discontinuation for specific reasons, they do give a good idea of the relative contributions of various reasons to method termination. Overall, pregnancy accounted for about one-fourth of reported terminations (even more if unreported abortions are taken into account), followed by stopping in order to become pregnant (20%) and inconvenience of the method (16%). However, the relative importance of the reasons for discontinuation varied greatly for different methods. Among oral contraceptive users, side effects accounted for 27% of discontinuation, followed by physician recommendations (20%), health concerns (14%), and a desire to become pregnant (13%). Among IUD users, physician recommendations were the overwhelming reason

for stopping (42%), followed by side effects (23%), and pregnancy (19%). Reasons for condom discontinuation were more evenly spread between method inconvenience (26%), desire for pregnancy (23%), and method failure (18%). Most withdrawal users terminated because of pregnancy (32%), desire for pregnancy (22%), or inconvenience of the method (15%).

Need for Family Planning Services

The concept of "women in need of family planning services" is one that is useful in determining the extent to which such services are reaching those who are at risk of becoming pregnant, but who would prefer not to. In [Table IV.15](#) two different definitions are used to estimate the proportion of women of childbearing age in need of family planning services. The first definition (A) considers women in need to be those who are in a sexual union, are not currently pregnant, are fecund, are not using any form of contraception, and do not want to become pregnant. Definition (B) also counts as in need women who are using withdrawal or natural family planning methods, because of the lower effectiveness of these methods (See [Table IV. 12](#)). According to these two definitions, 10% and 31%, respectively, of Czech women of reproductive age were in need of family planning services at the time of interview.


When definition A was used, women in need increased with age and number of living children and decreased as education and income (except for the lowest category) increased. The proportion in need was lowest among those with no religion. When definition B was used, most of the differentials noted disappear, indicating that most differences could be attributed to greater use of withdrawal in certain groups. Using this definition, we see that there were no appreciable differences in women in need according to place of residence, education, religion, or income. Even across age and number of children categories almost all differences disappeared, except that childless women were less likely than others to be in need.

Use of Less Effective Methods

Even though overall contraceptive prevalence is high in the Czech Republic, we have seen that there is considerable reliance on the use of methods of relatively lower effectiveness, particularly withdrawal. This reliance on less effective methods is, no doubt, one of the key factors in bringing about high rates of induced abortion in the Czech Republic. If all contraceptors were to use highly effective methods there would be relatively low levels of unmet need for family planning services.

Women who were avoiding pregnancy through the use of withdrawal (including withdrawal in combination with condoms) or natural family planning (NFP) methods were asked a series of questions regarding their method choice. Each woman was asked how important each of six factors was in the couple's selection of their current contraceptive method. The results of these questions, displayed in [Table IV.16](#) and

Figure IV.3
 Importance of Selected Factors in Decision
 to Use Traditional Contraception
 Percent Distribution, 1993 CRRHS



[Figure IV.3](#), demonstrate that only a fear that negative health effects are associated with the use of modern methods was an important consideration in the selection of withdrawal and NFP. Forty-four percent said that fear of health effects was very important and an additional 31% said it was somewhat important in making their decision. Among the other five factors, partner's preference was easily the most important, considered very important by about one in every ten instances. Cost, availability, and knowledge of other methods were each very important to 3-4% of women, while religious factors were not a consideration at all for 97% of these women.

Health considerations were important in choosing a contraceptive method for a large proportion of less effective method users in all age and education categories ([Table IV.17](#)). However, they were somewhat less important among women under 25 years of age. Cost, availability, and knowledge of methods were less likely to be reported as important factors as both age and education increased, so that few older or well educated women said they were at all important. The only group among which more than 5% said religion was important were those using NFP.

As shown in [Table IV. 18](#), almost half of users of less effective methods thought that their method was at least as effective as more modern methods, such as oral contraceptives and the IUD. Only one woman in three felt that her method was less effective. With increasing age there was a decreasing proportion of women who recognized the poorer reliability of their current method, As might be expected, the percentage recognizing the lower effectiveness of withdrawal and NFP increased sharply with education, from 18% to 41 %. There was little difference between the opinions of withdrawal and NFP users.

Problems with Current Contraceptive Methods

Based on responses to a question asked of all contraceptive users regarding the most significant problems or concerns they had with their current contraceptive method, it appears that most women are relatively happy with their method. Eighty-five percent said they had no major problems. This figure was fairly consistent between methods, ranging from 90% for NFP down to 80% for a combination of condoms and withdrawal ([Table IV.19](#)). Overall, about half of those with concerns reported that reliability/ quality was their major one. Small numbers of women reported health concerns, inconvenience, and side effects as problems. Reliability was especially a concern of withdrawal user and, to a lesser extent, among NFP and condom users. Health concerns were rare except among oral contraceptive (8%) and IUD (5%) users.

Consistent with this low level of reported concerns about current methods, 73% of current contraceptors stated that they would prefer to continue using their current method rather than switching to another one ([Table IV.20](#)). This percentage was fairly constant across users of most methods, except for IUD users, 90% of whom were satisfied with their method, and users of a combination of condoms and withdrawal, among whom only 54% preferred their current method(s).

Among those desiring to switch from their current method, the greatest number preferred to use oral contraceptives (40%), followed by the IUD (25%). Surprisingly, the third most preferred method (15% of those preferring another method) was implants, which were not yet available in the Czech Republic at the time of the survey. Of IUD users preferring to switch methods, 41% favored implants and an additional 25% preferred sterilization, either female or male. Few women wanted to switch from their current method to condoms, withdrawal, or natural methods.

Contraceptive Sterilization

Unlike many other countries of the world with high rates of contraceptive prevalence, the formerly communist states of central and eastern Europe have very low rates of surgical sterilization, both tubal ligation and, to an even greater extent, vasectomy. As was demonstrated earlier in this chapter, the Czech Republic is not an exception; sterilization is not widely practiced there. A major factor in the low prevalence of surgical sterilization in the Czech Republic is a law that prohibits the procedure for women with fewer than four children. Only three percent of married women of childbearing age report having undergone surgical sterilization, while no women report their partners to have undergone a vasectomy. Despite the facts that most women complete their desired childbearing many years before becoming unable to bear children and that contraception is widely practiced, relatively few women have been willing and able to become surgically sterilized. The major exception is the relatively small group of women who have at least four living children, over one-fourth of whom had been sterilized. The CRRHS questionnaire included several questions designed to learn more about the reasons for the infrequency of tubal ligation and to gauge the interest in this procedure among Czech women.


The results displayed in [Table IV.21](#) and [Figure IV.4](#), when viewed in conjunction with the small proportion of sterilized women, demonstrates a lack of interest in the procedure. Among the 2,345 respondents who desired no more children, only 8% said they were interested in undergoing tubal ligation, in addition to the 3% already sterilized. Another 7% were not sure whether they would consider it. Four out of every five women said they would not consider the procedure, despite reporting a desire to terminate childbearing. The low level of interest exists throughout the surveyed population (except, as mentioned among those with four or more children), with the proportion either already sterilized or interested in sterilization not exceeding 15% for any geographic, age, educational, or religious category examined.

Respondents reported a variety of reasons for not being interested in surgical sterilization ([Table IV.22](#)). The leading response, given by about one of every three women claiming not to be interested, was simply that they had not thought about undergoing the procedure. This is a strong indication that the low levels of sterilization stem not so much from a conscious decision not to be sterilized as it is from the fact that it has not been made available or discussed by health care providers as an option and that the

Figure IV.4

Interest in Surgical Sterilization among Women Wanting No More Children, by Number of Living Children

Percent Distribution, 1993 CRRHS



population knows little about the procedure. The second most commonly given response was a fear of health risks associated with the procedure (20%). This was followed by the fear that the woman/couple may decide to have another child (12%) and a fear of surgery (11 %). No other reason was mentioned by more than 4% of respondents wanting no more children. It is interesting to note that despite the opposition of the Catholic church to the procedure, only 1 % of women reported religious grounds for their lack of interest. There were only minor differences in the reasons given for women of different educational attainment. As education increased women were more likely to say they were worried about health risks or might want another child and less likely to cite a fear of surgery or objections of their husband.

There were several noteworthy differences between the characteristics of surgically sterilized women and the survey sample as a whole ([Table IV.23](#)). Of course, sterilized women tended to be older (two-thirds between ages 35 and 44) and have more children (97% with at least two) than the general population of 15-44 year-old women. In addition, though, about equal numbers of sterilized women lived in Bohemia and Moravia. One-third of sterilized women never went beyond primary school, even though this was true of only 18% of all respondents. Many sterilizations take place relatively early in women's lives: 45% of sterilized respondents had the procedure before the age of 30.

Use of Infertility Services

In trying to have the number of children they desire, couples often have need of services that help them to increase the likelihood of conceiving and bearing a living child, not just services that will help them to prevent conception and childbirth. The CRRHS included a module on the use of infertility services by respondents and their partners. As shown in [Table IV.24](#), 13% of women who had ever been in union had used infertility services at some point in their life, two-thirds of those before 1988. There was little difference in the proportion who had used these services according to respondent characteristics. The university educated respondents appear slightly less likely than others to have used infertility services.

[Table IV.25](#) displays the percent of those seeking infertility services from a physician or clinic who received specific types of treatment or advice. The most common service rendered was providing women/couples with advice on timing intercourse so as to maximize the probability of conception (54%). Forty percent of women and 37 percent of their partners had various tests performed on them. Thirty six percent had drugs prescribed to help them with their infertility problem, 21% were advised to try bed rest, the same percentage who reported that they underwent some type of surgical procedure. It appears that respondents in Moravia were more likely to undergo medical interventions than those in Bohemia. Among women seeking help for infertility, Moravian women were more likely to report having drugs prescribed, having surgery, or undergoing in vitro fertilization. As women become older it appears they are more

likely to receive medical interventions for their problem. Testing of both women and partner, surgery, the use of drugs, and especially in vitro fertilization are most common among those seeking treatment at ages 35 to 44.

TABLE IV.1
Percentage of Currently Married Respondents
Familiar with Various Contraceptive Methods, by Education
1993 Czech Republic Reproductive Health Survey

Contraceptive Method	Total	Education			
		Primary	Secondary No Diploma	Secondary Diploma	Any University
Condoms	99.3	97.8	99.4	99.7	100.0
Oral Contraceptives	98.6	96.1	98.5	99.8	100.0
IUD	96.6	90.9	96.0	99.6	99.8
Withdrawal	96.2	88.8	96.2	99.5	99.3
Natural Family Planning	94.2	83.6	93.9	99.0	99.8
Contraceptive Foam/Jelly	89.1	76.1	87.6	95.8	98.3
Female Sterilization	82.2	66.6	76.1	94.4	97.6
Diaphragm	71.4	49.3	64.6	86.2	93.9
Vasectomy	66.9	44.6	57.0	85.1	92.4
<i>Number of Respondents</i>	<i>(4,497)</i>	<i>(738)</i>	<i>(1,779)</i>	<i>(1,601)</i>	<i>(379)</i>

TABLE IV.2
Percentage of Currently Married Respondents Who Ever Used
Various Contraceptive Methods, by Education
1993 Czech Republic Reproductive Health Survey

Contraceptive Method	Age							Education			
	Total	15-19	20-24	25-29	30-34	35-39	40-44	Primary	Sec., No Diploma	Sec., Diploma	Any Univ.
Any Method	92.4	87.5	91.1	92.9	92.9	91.9	93.7	83.8	91.2	96.2	98.0
Any Modern Method*	79.4	58.3	71.1	80.5	81.5	80.3	83.7	68.6	76.5	85.6	85.7
Condoms	55.6	46.9	58.1	60.9	58.0	50.6	53.6	36.7	49.9	65.9	72.4
IUD	35.8	7.3	9.9	26.2	41.1	44.7	52.0	40.9	34.9	36.3	27.6
Oral Contraceptives	35.0	12.5	28.5	35.2	34.6	39.8	37.3	29.0	34.1	37.6	39.5
Vaginal Methods	5.3	8.3	4.8	6.4	4.6	5.7	4.3	2.4	5.3	5.7	9.3
Female Sterilization	2.7	0.0	0.0	2.0	3.0	4.1	3.7	4.8	2.5	1.9	2.3
Withdrawal	71.2	66.7	76.2	75.6	72.6	66.9	68.0	52.4	70.7	78.0	81.4
Natural Methods	33.4	15.6	30.2	37.8	35.8	31.8	33.6	21.2	25.2	40.8	60.8
No. of Respondents	(3,217)	(85)	(515)	(711)	(613)	(683)	(610)	(514)	(1,170)	(1,246)	(287)

*Modern methods are considered to be all methods listed except for withdrawal and natural methods.

TABLE IV.3
Percent of Currently Married Women Currently Using Any Contraceptive Method or
Any Modern Method, by Selected Characteristics
1993 Czech Republic Reproductive Health Survey

Characteristics	Percent Using:		Number of Respondents
	Any Contraception	Modern Contraception	
All Women	68.9	42.9	(3,217)
Region			
Bohemia, Total	69.0	41.8	(1,950)
Prague	69.0	41.3	(301)
Central Bohemia	76.8	47.5	(338)
South Bohemia	54.9	37.0	(221)
West Bohemia	66.4	44.0	(289)
North Bohemia	69.1	38.1	(406)
East Bohemia	72.1	50.7	(395)
Moravia, Total	68.9	44.5	(1,267)
South Moravia	68.1	46.3	(642)
North Moravia	69.8	47.3	(625)
Age			
15-19	51.0	27.1	(85)
20-24	59.6	33.7	(515)
25-29	71.2	46.4	(711)
30-34	78.3	49.5	(613)
35-39	71.1	43.4	(683)
40-44	65.4	42.2	(610)
Living Children			
0	28.5	14.6	(267)
1	64.0	36.9	(802)
2	75.1	47.0	(1,670)
3	77.8	51.3	(389)
4+	68.1	50.9	(89)
Education			
Primary	56.9	34.9	(514)
Secondary, No Diploma	65.1	37.9	(1,170)
Secondary, Diploma	76.7	50.8	(1,246)
Any University	75.4	44.5	(287)
Religion			
None	72.2	45.0	(2,034)
Catholic, attend mass	63.7	34.7	(166)
Catholic, not attend mass	64.0	41.0	(886)
Other	59.2	35.9	(122)

TABLE IV.4
Current Contraceptive Use, by Marital Status
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Current Use and Method*	Total	Marital Status		
		Married/ In Union	Previously Married	Never Married
Currently Using	59.1	68.9	45.5	37.5
Withdrawal only	18.6	22.2	10.9	11.5
Condoms only	15.4	16.7	7.2	14.4
IUD	11.6	15.3	16.8	0.3
Oral Contraceptives	7.7	8.0	7.4	6.8
Condoms + Withdrawal	2.2	2.0	0.5	3.2
Tubal Ligation	1.9	2.7	1.5	0.0
Natural Methods	1.5	1.9	1.2	0.6
Vaginal Methods	0.3	0.2	0.0	0.6
Not Using	40.9	31.1	54.5	62.5
Total	100.0	100.0	100.0	100.0
<i>Number of Respondents</i>	<i>(4,497)</i>	<i>(3,217)</i>	<i>(355)</i>	<i>(925)</i>

*Women reporting current use of more than one method were considered to be users of the more effective method, except for those reporting use of condoms and withdrawal, which is listed as a separate category in this table.

TABLE IV.5
Current Contraceptive Use, by Number of Living Children,
Women Married or in Union
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Current Use and Method*	Total	Living Children				
		0	1	2	3	4+
Currently Using	68.9	28.5	64.0	75.1	77.8	68.1
Withdrawal only	22.2	11.4	22.8	23.8	23.8	11.5
Condoms only	16.7	8.5	17.8	18.0	16.4	7.1
IUD	15.3	1.1	9.7	18.3	21.8	16.8
Oral Contraceptives	8.1	4.3	9.0	8.5	8.0	3.5
Tubal Ligation	2.7	0.4	0.2	2.1	4.9	26.5
Condoms + Withdrawal	2.0	1.4	2.3	2.2	1.2	0.0
Natural FP	1.9	1.1	2.0	2.1	1.4	2.7
Vaginal Methods	0.2	0.4	0.0	0.3	0.2	0.0
Not Using	31.1	71.5	36.0	24.9	22.2	31.9
Total	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of Respondents</i>	<i>(3,217)</i>	<i>(267)</i>	<i>(802)</i>	<i>(1,670)</i>	<i>(389)</i>	<i>(89)</i>

*Women reporting current use of more than one method were considered to be users of the more effective method, except for those reporting use of condoms and withdrawal, which is listed as a separate category in this table.

TABLE IV.6
Current Contraceptive Use, by Age
Women Married or in Union,
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Current Use and Method*	Age						
	Total	15-19	20-24	25-29	30-34	35-39	40-44
Currently Using	68.9	51.0	59.6	72.8	78.3	71.2	65.4
Withdrawal only	22.2	20.8	22.2	20.0	24.6	24.3	20.3
Condoms only	16.7	15.6	15.5	16.7	17.4	15.8	17.9
IUD	15.3	5.2	7.3	15.3	17.9	17.4	17.9
Oral Contraceptives	8.1	6.3	10.6	12.5	10.8	6.0	2.6
Tubal Ligation	2.7	0.0	0.0	2.0	3.0	4.1	3.7
Condoms + Withdrawal	2.0	3.1	2.6	3.1	2.2	1.4	1.0
Natural FP	1.9	0.0	1.1	2.4	2.1	2.2	1.9
Vaginal Methods	0.2	0.0	0.4	0.0	0.3	0.1	0.3
Not Using	31.1	49.0	40.4	27.2	21.7	28.8	34.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>Number of Respondents</i>	<i>(3,217)</i>	<i>(85)</i>	<i>(515)</i>	<i>(711)</i>	<i>(613)</i>	<i>(683)</i>	<i>(610)</i>

*Women reporting current use of more than one method were considered to be users of the more effective method, except for those reporting use of condoms and withdrawal, which is listed as a separate category in this table.

TABLE IV.7
Current Contraceptive Use, by Education of Respondent,
Women Married or in Union
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Current Use and Method*	Total	Education			
		Primary	Secondary No Diploma	Secondary Diploma	Any University
Currently Using	68.9	56.9	65.1	76.7	75.4
Withdrawal only	22.2	19.4	24.3	21.0	24.3
Condoms only	16.7	10.6	13.3	20.9	24.6
IUD	15.3	15.2	13.8	17.8	11.0
Oral Contraceptives	8.1	4.2	8.2	10.1	6.3
Tubal Ligation	2.7	4.8	2.5	1.9	2.3
Condoms + Withdrawal	2.0	0.8	2.0	2.5	2.3
Natural FP	1.9	1.8	0.9	2.5	4.3
Vaginal Methods	0.2	0.2	0.2	0.2	0.3
Not Using	31.1	43.1	34.9	23.3	24.6
Total	100.0	100.0	100.0	100.0	100.0
<i>Number of Respondents</i>	<i>(3,217)</i>	<i>(514)</i>	<i>(1,170)</i>	<i>(1,246)</i>	<i>(287)</i>

*Women reporting current use of more than one method were considered to be users of the more effective method, except for those reporting use of condoms and withdrawal, which is listed as a separate category in this table.

TABLE IV.8
Current Contraceptive Use, by Religion,
Women Married or in Union
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Current Use and Method*	Total	Religion			
		None	Catholic, Attends	Catholic, Not Attend	Other
Currently Using	68.9	72.9	63.7	64.0	59.2
Withdrawal only	22.2	23.2	22.3	19.8	19.7
Condoms only	16.7	17.0	15.0	16.2	17.6
IUD	15.3	14.6	13.0	18.1	9.9
Oral Contraceptives	8.1	10.4	1.6	4.6	5.6
Tubal Ligation	2.7	2.8	4.7	2.1	2.8
Condoms + Withdrawal	2.0	2.3	2.1	1.4	2.1
Natural FP	1.9	1.7	4.7	1.9	1.4
Vaginal Methods	0.2	0.3	0.5	0.0	0.0
Not Using	31.1	27.1	36.3	36.0	40.8
Total	100.0	100.0	100.0	100.0	100.0
<i>Number of Respondents**</i>	<i>(3,217)</i>	<i>(2,034)</i>	<i>(166)</i>	<i>(886)</i>	<i>(122)</i>

*Women reporting current use of more than one method were considered to be users of the more effective method, except for those reporting use of condoms and withdrawal, which is listed as a separate category in this table.

**Nine women did not state their religion.

TABLE IV.9
Current Contraceptive Use, by Whether Respondent Wants Any More Children
Fecund, Nonpregnant Women Currently in Union
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Current Use and Method*	Total	Want No More Children	Want More Children
Currently Using	81.5	81.6	80.8
Withdrawal only	26.4	25.4	31.1
Condoms only	19.8	19.3	22.0
IUD	18.1	20.4	8.7
Oral Contraceptives	9.2	8.5	12.4
Condoms + Withdrawal	2.1	1.8	3.6
Tubal Ligation	3.4	4.2	0.0
Natural Methods	2.0	1.9	2.6
Vaginal Methods	0.3	0.2	0.6
Not Using	18.5	18.4	19.2
Total	100.0	100.0	100.0
<i>Number of Respondents</i>	<i>(2,475)</i>	<i>(1,972)</i>	<i>(503)</i>

*Women reporting current use of more than one method were considered to be users of the more effective method, except for those reporting use of condoms and withdrawal, which is listed as a separate category in this table.

TABLE IV.10
Source of Oral Contraceptives and IUDs by Region
for Current Users of Those Methods
1993 Czech Republic Reproductive Health Survey

Method/ Source	Total	Region							
		Prague	Central Bohemia	South Bohemia	West Bohemia	North Bohemia	East Bohemia	South Moravia	North Moravia
Oral Contraceptives									
Reg./Dist. Physician	61.4	80.0	41.0	70.4	60.0	60.6	58.5	57.3	67.6
Pharmacy	17.1	12.5	28.2	0.0	14.3	15.2	13.9	22.7	20.6
Polyclinic Physician	11.1	2.5	20.5	7.4	11.4	10.6	16.9	12.0	5.9
Private Physician	7.5	5.0	2.6	22.2	8.6	10.6	9.2	6.7	1.5
Hospital	2.9	0.0	7.7	0.0	5.7	3.0	1.5	1.3	4.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number of Users	(375)	(38)	(35)	(25)	(34)	(62)	(60)	(59)	(62)
IUD									
Reg./Dist. Physician	47.3	70.0	55.2	68.4	45.1	55.2	34.8	25.2	54.8
Hospital	34.9	7.5	20.9	10.5	33.3	28.4	44.9	56.3	34.4
Polyclinic Physician	15.7	17.5	22.4	21.1	19.6	10.4	17.4	17.0	10.2
Private Physician	2.1	5.0	1.5	0.0	2.0	6.0	2.9	1.5	0.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number of Users	(557)	(37)	(59)	(33)	(48)	(62)	(62)	(114)	(142)

TABLE IV.11
Primary Reason for Not Using Contraception, by Marital Status
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Reason Not Using Contraception	Total	Marital Status		
		Married/ In Union	Previously Married	Never Married
Reasons Related to Pregnancy, Fecundity, and Sexual Activity	68.4	53.4	77.3	85.8
Not Sexually Active	37.7	2.2	54.6	79.0
Subfecund	12.9	20.9	17.7	1.4
Pregnant	9.1	16.2	0.5	2.1
Trying to Become Pregnant	8.7	14.1	4.5	2.7
Other Reasons	31.6	46.6	22.7	14.2
Fear Health Effects	7.2	11.8	4.5	2.0
Difficult to Get Pregnant	3.9	6.9	3.2	0.1
Occasional Sex Only	3.6	2.0	6.4	5.0
Postpartum/Breastfeeding	3.1	6.0	0.0	0.2
Previous Side Effects	2.6	4.4	2.7	0.2
Partner Opposes Use	1.9	3.2	0.9	0.3
Haven't Bothered	2.2	1.9	0.9	3.0
Access, Cost, Etc.	1.4	1.6	2.3	0.8
Religion	0.5	0.4	0.0	0.6
Other Reasons	1.6	2.6	0.0	0.7
Not Sure	3.7	5.4	1.8	2.0
Total	100.0	100.0	100.0	100.0
<i>Number of Respondents</i>	<i>(1,736)</i>	<i>(978)</i>	<i>(190)</i>	<i>(568)</i>

TABLE IV.12
Contraceptive Failure Rates, in Percents, for Selected Methods
after 12, 24, and 36 Months for Segments of Use Beginning after 1987
1993 Czech Republic Reproductive Health Survey

Contraceptive Method	Failure Rates after:		
	12 Months	24 Months	36 Months
Oral Contraceptives	2.4	5.4	9.1
IUD	3.1	5.7	6.6
Condoms	5.6	11.7	17.4
Withdrawal	11.9	23.1	32.1
Natural Family Planning	20.8	34.8	40.7

TABLE IV.13
Contraceptive Discontinuation Rates, in Percents, for Selected Methods
after 12, 24, and 36 Months for Segments of Use Beginning after 1987
1993 Czech Republic Reproductive Health Survey

Contraceptive Method	Discontinuation Rates after:		
	12 Months	24 Months	36 Months
Oral Contraceptives	39.5	57.0	66.8
IUD	13.0	20.2	26.4
Condoms	32.7	51.4	62.2
Withdrawal	36.0	53.6	67.3
Natural Family Planning	40.2	59.8	70.8

TABLE IV.14
Most Important Reason Reported for Discontinuing Use
of Contraceptive Method according to Method Used,
All Reported Segments of Use Beginning and Ending after 1987
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Reason for Discontinuation	Contraceptive Method				
	All Methods*	Oral Contracep.	IUD	Condoms	Withdrawal
Pregnancy	23.6	6.8	19.2	17.7	32.4
Desired Pregnancy	20.1	12.7	8.0	22.8	21.8
Method Inconvenient	15.7	4.8	0.8	25.5	15.3
Physician's Recommendation	8.3	19.7	42.4	5.2	3.1
Relationship End/Infreq. Sex	8.0	6.5	0.8	11.4	7.6
Side Effects	6.0	27.9	23.2	0.4	0.5
Partner Objected	4.7	0.9	0.8	5.9	6.1
Health Concerns	2.9	13.5	2.4	1.0	0.4
Switched to Better Method	2.5	0.0	0.0	1.3	4.4
Availability of Method	0.8	3.9	0.0	0.6	0.0
Other	4.6	2.5	2.4	4.3	6.0
Does Not Remember	2.7	0.9	0.0	3.9	2.6
Total	100.0	100.0	100.0	100.0	100.0
<i>Number of Segments of Use</i>	<i>(2,043)</i>	<i>(319)</i>	<i>(117)</i>	<i>(608)</i>	<i>(847)</i>

*Includes use of methods not listed separately.

TABLE IV.15
Percent of Women in Need of Family Planning Services,
according to Two Definitions, by Selected Characteristics
1993 Czech Republic Reproductive Health Survey

Characteristics	Definition A*	Definition B**	Number of Respondents
Total	9.8	30.6	(4,497)
Residence			
Bohemia	10.2	32.2	(2,737)
Moravia	9.1	28.2	(1,760)
Size of Place			
Less than 5,000	12.7	32.2	(1,472)
5,000-19,999	9.6	28.4	(967)
20,000 +	7.7	30.5	(2,058)
Age			
15-19	2.1	16.0	(662)
20-24	9.7	33.3	(756)
25-29	8.9	31.3	(828)
30-34	10.0	34.5	(720)
35-39	13.3	36.7	(805)
40-44	15.6	34.9	(726)
Living Children			
0	1.7	15.7	(1,165)
1	10.5	34.0	(982)
2	14.8	39.4	(1,832)
3	11.7	35.4	(419)
4 or More	18.3	31.8	(99)
Education			
Primary only	14.2	29.6	(738)
Secondary, No Diploma	10.6	31.7	(1,779)
Secondary, Diploma	7.1	29.9	(1,601)
Any University	6.4	30.1	(379)
Religion			
None	8.2	29.9	(2,913)
Catholic, Attends Services	9.9	28.9	(256)
Catholic, Does Not Attend	13.4	32.6	(1,134)
Other	12.8	31.5	(180)
Monthly Household Income (Crown)			
Less than 3,000	5.5	22.1	(188)
3,000-6,999	12.6	31.8	(1,885)
7,000-9,999	9.9	32.4	(1,278)
10,000-14,999	6.3	29.4	(615)
15,000 or More	7.3	39.7	(188)
Don't Know	3.0	16.1	(253)
Not Stated	13.6	37.9	(90)

*Women in a sexual union, not pregnant, fecund, and not desiring pregnancy and not using any method of contraception.

**The same as definition one plus couples using withdrawal or natural family planning methods.

TABLE IV.16
Importance of Selected Factors in Couples' Decisions
to Use Traditional Methods of Contraception
Current Users of Traditional Methods
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Factor	Importance of Factors				Total
	Very	Somewhat	Not at All	Not Sure	
Fear of Health Effects	43.8	31.0	22.0	3.2	100.0
Partner Preference	10.7	13.7	69.6	5.9	100.0
Cost of Methods	4.0	8.4	85.0	2.6	100.0
Method Availability	3.9	8.2	85.0	2.9	100.0
Knowledge of Methods	3.0	8.5	85.3	3.1	100.0
Religion	1.6	1.4	96.7	0.4	100.0

Number of Respondents = 819

TABLE IV.17
Percent of Current Users of Withdrawal or Natural Family Planning
Who Reported that Selected Factors Were Very Important or Somewhat Important
in The Selection of Their Contraceptive Method, by Age, Education, and Method
1993 Czech Republic Reproductive Health Survey

Characteristics	Factors Involved						No. of Users
	Health Effects	Partner Preference	Cost of Methods	Method Availabil.	Knowledge of Methods	Religion	
Total	74.8	24.4	12.4	12.1	11.5	3.0	(819)
Age							
15-19	63.1	34.2	21.6	18.9	26.1	0.0	(67)
20-24	69.5	25.6	17.1	14.6	15.9	2.4	(148)
25-29	76.0	20.9	10.4	10.1	9.5	3.8	(156)
30-34	81.7	20.7	14.0	11.0	9.2	4.9	(161)
35-39	78.8	25.4	10.4	13.0	9.3	1.6	(160)
40-44	75.4	22.8	3.5	7.0	4.6	4.1	(127)
Education							
Primary	71.6	25.5	17.7	19.2	17.0	2.1	(107)
Sec No Dipl	70.1	27.4	12.4	12.4	13.7	3.0	(337)
Sec Dipl	80.9	21.3	11.3	10.1	9.0	2.9	(297)
Any Univ.	77.5	17.5	7.5	6.3	2.5	3.8	(78)
Method							
Natural FP	74.1	27.2	14.8	9.9	6.2	8.6	(73)
Withdrawal	74.9	24.2	12.2	12.3	12.1	2.4	(746)

TABLE IV.18
Opinions of Current Users of Withdrawal or Natural Family Planning
regarding the Effectiveness of Their Current Method
Relative to Modern Methods, by Age, Education, and Method
1993 Czech Republic Reproductive Health Survey

Characteristics	Relative Effectiveness of Current Method				Total	No. of Users
	More Effective	Equally Effective	Less Effective	Do Not Know		
Total	12.1	36.8	33.7	14.7	100.0	(819)
Age						
15-19	5.4	25.2	45.1	24.3	100.0	(67)
20-24	10.4	32.9	41.5	15.2	100.0	(148)
25-29	15.2	34.8	38.6	11.4	100.0	(156)
30-34	7.9	39.6	31.1	21.3	100.0	(161)
35-39	14.5	46.6	25.4	13.5	100.0	(160)
40-44	16.4	36.3	26.3	21.1	100.0	(127)
Education						
Primary	17.0	37.6	18.4	27.0	100.0	(107)
Sec No Diploma	12.9	34.8	35.0	17.3	100.0	(337)
Sec Diploma	9.8	38.2	36.7	15.3	100.0	(297)
Any University	8.8	40.0	41.3	10.0	100.0	(78)
Method						
Natural FP	11.1	30.9	37.0	21.0	100.0	(73)
Withdrawal	12.2	37.4	33.4	17.1	100.0	(746)

TABLE IV.19
Most Important Problem or Concern with Current Contraceptive Method
according to Method Used
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Major Problem/ Concern	Current Contraceptive Method					
	All Methods	Orals	IUD	Condoms	Natural Methods	Withdrawal Condoms + Withdrawal
No Problems	84.5	83.4	87.2	85.3	90.2	79.7
Reliability/Quality	8.0	1.4	2.2	8.6	7.3	14.4
Health Concerns	2.9	5.3	8.2	1.1	0.0	0.9
Inconvenience	2.5	2.2	0.2	4.3	2.4	4.2
Side Effects	1.7	7.2	2.2	0.1	0.0	0.9
Other	0.4	0.5	0.0	0.6	0.0	0.0
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number of Users	(2,670)	(376)	(557)	(696)	(74)	(101)

TABLE IV.20
Preference for Current or Other Method, Current Contraception Users
and Method Preferred among Women Preferring Another Method
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Preferred Method	All Methods*	Current Contraceptive Method					
		Oral Contraceptives	IUD	Condoms	Natural FP	Withdrawal	Condoms + Withdrawal
Current Method	73.2	78.8	90.2	66.3	73.2	68.1	54.2
Other Method	26.8	21.2	9.8	37.7	26.8	31.9	45.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number of Users	(2,670)	(378)	(557)	(696)	(74)	(854)	(101)
Pref. Method of Women Wanting to Switch							
Orals	39.8	---	21.3	47.3	**	40.6	66.7
IUD	24.8	37.5	---	24.4	**	28.8	13.0
Implants	14.5	28.4	41.0	8.2	**	12.2	9.3
Female Steriliz.	6.7	5.7	13.1	7.2	**	6.3	3.7
Condoms	3.1	5.7	0.0	---	**	6.3	0.0
Vasectomy	2.4	4.6	11.5	2.5	**	0.6	0.0
Other	4.5	6.8	6.6	5.7	**	1.9	3.7
Not Sure	4.6	11.4	6.6	4.7	**	3.4	3.7
Total	100.0	100.0	100.0	100.0	**	100.0	100.0
Number of Users	(708)	(77)	(60)	(231)	(20)	(274)	(44)

*Includes 12 women using methods not listed individually.

**Fewer than 25 users of natural family planning who would prefer to switch to another method.

TABLE IV.21
Interest in Surgical Sterilization
among Fecund Women Wanting No More Children, by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Interest in Sterilization				Total	Number of Respondents
	Already Sterilized	Interested	Not Interested	Not Sure		
Total	3.8	8.2	81.3	6.8	100.0	(2,345)
Region						
Bohemia	3.2	8.2	82.6	5.9	100.0	(1,415)
Moravia	4.6	8.1	79.2	8.1	100.0	(930)
Age						
15-24	0.0	13.2	80.5	6.3	100.0	(156)
25-34	3.7	11.0	76.2	9.2	100.0	(910)
35-44	4.3	6.0	84.3	5.4	100.0	(1,279)
Living Children						
0-1	0.7	5.0	91.0	3.3	100.0	(382)
2	2.4	8.9	81.0	7.8	100.0	(1,509)
3	6.0	8.1	79.1	6.8	100.0	(371)
4 or More	29.8	9.6	55.8	4.8	100.0	(83)
Education						
Primary	6.3	8.4	79.9	5.4	100.0	(451)
Sec, No Dipl.	3.5	9.8	78.2	8.6	100.0	(859)
Sec, Diploma	2.6	6.8	84.0	6.5	100.0	(868)
Any Univ.	3.9	6.2	87.1	2.8	100.0	(167)
Religion						
None	3.8	8.6	80.3	7.3	100.0	(1,470)
Cath, Attends	7.4	3.7	83.8	5.2	100.0	(107)
Cath, Not Att	3.2	8.3	82.7	5.8	100.0	(670)
Other	3.6	6.4	82.7	7.3	100.0	(92)

TABLE IV.22
Primary Reason for Lack of Interest in Surgical Sterilization
among Fecund Women Wanting No More Children, by Education
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Reason Not Interested	Total	Respondent Education			
		Primary	Secondary, No Diploma	Secondary, Diploma	Any University
Haven't Thought About	34.2	35.2	35.2	34.3	25.2
Health Risk	20.1	15.9	19.6	22.3	23.2
Might Want Another Child	12.3	7.2	12.0	14.2	18.1
Fear of Surgery	10.8	14.6	12.6	7.7	7.7
Cultural/Social Reasons	4.1	2.7	4.8	4.1	5.2
Husband Objects	3.5	5.4	3.3	3.1	1.3
Don't Know Enough About	3.1	4.3	3.9	1.9	1.9
Too Old/Subfecund	1.7	2.9	1.6	1.5	0.6
Little Sexual Activity	1.4	2.7	1.3	0.9	0.6
Curr. Method Satisfactory	1.3	0.7	0.9	1.6	3.2
Religion	1.0	1.1	0.7	1.2	0.6
Other	3.4	3.4	1.6	3.5	10.3
Don't Know	3.2	4.0	2.6	3.6	1.9
Total	100.0	100.0	100.0	100.0	100.0
<i>Number of Respondents</i>	<i>(1,887)</i>	<i>(354)</i>	<i>(662)</i>	<i>(726)</i>	<i>(145)</i>

TABLE IV.23
Profile of Respondents Who Have Been Surgically Sterilized Compared to All Respondents
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Sterilized Respondents	All Respondents
Residence		
Bohemia	51.0	61.0
Moravia	49.0	39.0
Size of Place		
Less than 5,000	37.3	33.1
5,000-19,999	23.5	19.0
20,000+	39.2	48.0
Current Age		
Under 30	14.7	48.0
30-34	18.6	15.5
35-39	34.3	18.1
40-44	32.4	18.5
Living Children		
0-1	3.0	49.3
2	39.2	37.1
3	27.5	10.9
4 or More	30.4	2.8
Education		
Primary only	34.3	18.0
Secondary, No Diploma	33.3	40.7
Secondary, Diploma	25.5	33.8
Any University	6.9	7.6
Religion		
None	61.8	64.5
Catholic, Attends Services	9.8	6.0
Catholic, Does Not Attend	24.5	25.2
Other	3.9	4.1
Not Stated	0.0	0.3
Age at Operation		
20-24	11.8	--
25-29	33.3	--
30-34	23.5	--
35-39	28.4	--
40-44	2.9	--
Year of Operation		
Before 1985	30.4	--
1985-1989	34.3	--
1990-1991	20.6	--
1992-1993	14.7	--
Number of Respondents	(89)	(4,497)

TABLE IV.24
Use of Infertility Services according to
When Services Were Last Received and Selected Characteristics,
Women Ever in Union
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Last Use of Infertility Services			Total	Number of Respondents
	Never Used	Before 1988	1988 or Later		
Total	87.1	8.4	4.5	100.0	(3,572)
Residence					
Bohemia	87.2	8.4	4.4	100.0	(2,189)
Moravia	87.0	8.4	4.7	100.0	(1,383)
Size of Place					
Less than 5,000	88.8	7.8	3.4	100.0	(1,177)
5,000-19,999	86.7	8.0	9.1	100.0	(783)
20,000 +	86.1	9.1	4.9	100.0	(1,612)
Current Age					
15-19	88.8	0.0	11.2	100.0	(87)
20-24	91.5	0.9	7.6	100.0	(542)
25-29	86.1	4.3	9.6	100.0	(767)
30-34	85.2	10.5	4.4	100.0	(688)
* 35-39	86.2	11.6	2.2	100.0	(782)
40-44	87.5	12.5	0.0	100.0	(706)
Education					
Primary only	88.4	8.5	3.1	100.0	(594)
Secondary, No Diploma	88.3	7.1	4.7	100.0	(1,292)
Secondary, Diploma	86.3	8.9	4.8	100.0	(1,377)
Any University	82.8	12.0	5.2	100.0	(309)
Religion					
None	87.3	8.0	4.7	100.0	(2,241)
Catholic	87.0	8.8	4.2	100.0	(1,141)
Other	85.8	7.1	3.4	100.0	(178)

TABLE IV.25

Percent of Respondents Receiving Selected Treatment/Counseling
among Those Who Visited a Physician or Clinic for Infertility Problems
according to Place of Residence and Age When Services Last Received
1993 Czech Republic Reproductive Health Survey

Treatment Received	Total	Region of Residence		Age When Last Received Treatment			
		Bohemia	Moravia	Under 25	25-34	35-44	
Advice on Timing Intercourse	54.1	56.0	51.2	54.7	53.2	52.5	
Tests on Her	40.3	38.3	43.4	35.6	44.4	57.4	
Tests on Partner	36.6	39.2	32.7	30.4	41.3	60.7	
Prescribed Drugs	36.2	29.6	46.3	35.0	34.9	45.9	
Prescribed Bed Rest	21.3	20.6	22.4	22.2	21.4	16.4	
Surgery	20.7	19.6	22.4	17.6	24.6	29.5	
Advice on Timing Contraception	12.6	13.2	11.7	14.9	10.3	4.9	
Advice to Reduce Stress	5.8	5.8	5.9	6.7	4.8	3.3	
Advice to Exercise/Go to Spa	4.3	4.5	3.9	4.0	4.0	6.6	
Artificial Insemination	3.7	3.5	3.9	2.4	7.1	3.3	
In Vitro Fertilization	1.7	1.3	2.4	0.9	0.8	8.2	
Natural Medicines	1.4	0.3	2.9	1.8	0.8	0.0	
Other	5.0	6.1	3.4	4.6	5.6	6.6	
<i>Number of Respondents</i>	<i>(478)</i>	<i>(290)</i>	<i>(188)</i>	<i>(301)</i>	<i>(120)</i>	<i>(57)</i>	

V. PREGNANCY, DELIVERY, AND INFANT HEALTH

The 1993 CRRHS included a module, administered to all respondents who had given birth since the beginning of 1988, on maternal and child health issues related to her most recent pregnancy and live birth. The topics covered in this module included: reactions to learning she was pregnant; stress and physical activity during pregnancy; prenatal counselling; amount, timing, and source of prenatal care; use of ultrasound; complications/hospitalization during pregnancy; use of cigarettes and alcohol during pregnancy; birth weight; type of delivery; information on the delivery (including type of delivery, analgesia, and quality of care issues); and problems encountered after the child's birth.

Unlike reproductive health surveys in developing countries, the 1993 CRRHS did not have as its objectives the measurement and detailed analysis of infant morbidity and mortality. The national medical statistics system compiles detailed and virtually complete information on infant deaths in the Czech Republic and collects information on other aspects of infant morbidity and mortality as well. Infant mortality in the Czech Republic in 1993 was reported to be about 9 deaths per 1,000 live births. This level was lower than the rate reported for any other former communist country in Europe. However, the rate was still slightly higher than for most western European countries.

Even though it was not necessary for the survey to measure infant and early childhood mortality, it was decided that it was important to collect information on factors known to be related to the health of infants, most of which are not routinely included in the medical statistics system. These factors include such things as breastfeeding, cigarette smoking during pregnancy, and use of prenatal services. These data can be used to determine the prevalence of selected risk factors and to learn in which segments of the population is the health of infants most at risk from these factors.

Reaction to Pregnancy

Among those women who had a live birth since the beginning of 1988, about two-thirds said they were immediately happy about the pregnancy that resulted in their most recent birth ([Table V.1](#)). About one-fourth said they easily accepted the pregnancy, 8% eventually accepted it, and no women said they never accepted the pregnancy. Since we know there are many unintended pregnancies in the Czech Republic, we do not know the degree to which these results indicate that Czech women easily adjust to unintended pregnancies or that most truly unintended pregnancies are aborted. Poorly educated women were the least likely to be immediately happy about their pregnancy.

The distribution of reactions to pregnancy among spouses/partners, according to respondents, was barely distinguishable to that among respondents themselves ([Table V.2](#)). A small percentage reportedly never accepted the pregnancy. As with

respondents, poorly educated men were the least likely to be immediately happy.

Physical Exertion/Stress during Pregnancy

Heavy physical exertion and emotional stress are thought by some to increase the risk of adverse pregnancy outcomes. CRRHS respondents were asked about the extent of physical exertion, standing at their jobs, and on-the-job stress during their most recent pregnancy. [Tables V.3](#), [V.4](#), and [V.5](#) display distributions of the degree of job-related heavy physical work, standing, and stress, respectively, during respondents' most recent pregnancy leading to a live birth. Overall, 15% of women described themselves as having done much physical work and 27% said they did a large amount of standing during their jobs. Not surprisingly, the amount of both physical exertion and standing decreased sharply as educational level increased. Physical exertion and standing were both least common among women in their twenties.

Seventeen percent of women said they were under a large amount of stress on their jobs during pregnancy and 32% said they were under moderate stress. The proportion under great stress was much higher among university educated women (41%) than among others (11%-18%).


Prenatal Care

As seen in [Table V.6](#), the Czech Republic has an impressive record regarding prenatal care coverage, which has become almost universal. Fewer than 1% of women with a live birth since the beginning of 1988 received no prenatal care during their most recent pregnancy leading to a live birth. Ninety-four percent of those women began receiving prenatal care during their first trimester, while only 5% waited until after the first trimester. (Of this 5%, most started receiving care in the fourth month of pregnancy.)

The percentage of women not receiving early prenatal care, seen in [Figure V.1](#), ranged from 3% in South Bohemia to 10% in North Bohemia and 9% in East Bohemia (both with 2% receiving no prenatal care). Receipt of prenatal care appears to be unrelated to the size of community. The groups of women who were less likely to begin care in the first trimester included those who: were less than 20 years of age or between 35 and 44 years, were not currently married, had only a primary education, were affiliated with a religion other than Catholicism, and had low household income. However, for only a few groups did the proportion receiving early prenatal care drop substantially below 90%: those belonging to "other" religions (83%) and those with only a primary education (87%).

Not only did women tend to begin prenatal care early, but 91% reported making at least 10 prenatal visits during pregnancy, with only 1% reporting fewer than six visits ([Table V.7](#)). The proportion of women making at least 10 prenatal visits was high across all geographic and socioeconomic categories, but was especially high in Prague (97%) and

Figure V.1
 Percent of Respondents With No Prenatal Care
 During Their First Trimester, by Region
 1993 CRRHS



East Bohemia (95%).

Almost two-thirds of those who made prenatal visits, reported that they received their care primarily from physicians, while one-third received care from both physicians and nurses/midwives. Only 3% said that nurse/midwives were their primary providers ([Table V.8](#)). Women from communities with fewer than 5,000 people were less likely than others to receive care primarily from physicians. Within regions the percentages primarily seeing physicians ranged from 70% in Prague to 54% in North Bohemia. Women at ages 15-19 and 40-44, as well as women with only a primary education, were much less likely than others to have received care principally from physicians.

Sixty percent of women received their prenatal care primarily at a district clinic, with the remainder going to a combination of district clinics and hospitals, or mainly a hospital ([Table V.9](#)). Women were less likely to primarily utilize district clinics if they lived in larger cities (54%), lived in Prague (37%), or were 40-44 years old (45%).

A large majority of women lived within 30 minutes of their usual source of prenatal care, regardless of their usual means of transportation, with a mean time to source of care of about 22 minutes ([Table V.10](#)). Only 1% of women required more than one hour to reach their care provider.


Percentage distributions of usual waiting time to receive prenatal care during the most recent pregnancy ending in a live birth are displayed in [Table V.11](#). Overall, 44% typically waited for over 30 minutes, while 15% waited for over one hour. Waiting time appears to be longer in medium sized places than in larger or smaller towns and cities. Of the regions in the Czech Republic, waiting time was the longest in West Bohemia and Central Bohemia, where over half of women typically waited more than 30 minutes for care. The shortest waiting times were in Prague. Interestingly, as women's level of education increased their reported waiting time tended to decrease.

Ultrasound Examinations

Ninety-two percent of women who received prenatal care reported that they underwent an ultrasound examination during their last pregnancy leading to a live birth ([Table V.12](#)). The percentage having no ultrasound exam varied from 2% in South Bohemia and 3% in South Moravia up to 15% in North Moravia and 14% in West Bohemia ([Figure V.2](#)). These differences indicate possible differences in providers' practices or regional availability of ultrasound. The procedure was more commonly done for the youngest (15-24 years) and oldest (40-44) groups of women. The proportion receiving no ultrasound exam also decreased with education, from 11% for women with primary education only to 3% for those who attended university.

Despite the high ultrasound coverage, there was some concern that women were not being adequately informed of the purpose of these examinations. The CRRHS found that

Figure V.2
**Percent of Respondents with No Ultrasound Exam
 During Her Most Recent Pregnancy**
 1993 CRRHS



one-fourth of women who had undergone ultrasound reported that no one had explained the reason for doing this procedure and another one-fifth had only had the procedure explained to them slightly ([Table V.13](#)). Women in small towns were the most likely to say the procedure had been explained well. Among the regions of the country, women received the least explanation in Prague and in East Bohemia. Multiparous women were more likely than primiparous women to report having had the procedure explained to them, which may indicate that previous experience affects women's perceptions of how well procedures are explained. Women at ages 35-39 and 40-44, as well as well educated women, were much more likely than others to have had the procedure's purpose explained.

Bed Rest/Hospitalization during Pregnancy


[Table V.14](#) reveals that about three of every ten women had bed rest prescribed for part of their most recent pregnancy resulting in a live birth. The percentage varied from 38% in North Bohemia to 23% in West Bohemia, with little variation in prescribed length of bed rest. On the other hand, while there was little difference by age in the percent for whom bed rest was prescribed, older women tended to have considerably longer bed rest prescribed. The likelihood of bed rest increased slightly with educational level.

Hospitalization during pregnancy (other than for labor and delivery) is a common practice in the Czech Republic. Twenty-nine percent of women giving birth since 1988 were hospitalized during some part of their pregnancy, with a mean length of stay of almost one month and a median of two weeks ([Table V.15](#)). (It should be kept in mind that these figures exclude pregnancies not resulting in a live birth.) The percentage hospitalized is relatively high, but is not surprising in a system where there has been little economic pressure not to hospitalize individuals and where hospitalization provides an easy means of observing possible complications. There was relatively little geographic difference in the proportion hospitalized. Although older mothers were no more likely than others to be hospitalized during pregnancy, they tended to be hospitalized for the longest periods of time (mean = 57 days, median = 21 days). Primiparous women were slightly more likely to have been hospitalized than multiparous women.

As shown in [Table V.16](#), the conditions most commonly cited as being responsible for hospitalization during pregnancy were bleeding during the first half of pregnancy (39% of hospitalizations) and early or false labor (33%). Much less common were excessive swelling or edema (14%), hypertension related to the pregnancy (11%), and bleeding during the second half of pregnancy (11 %). Among those who were hospitalized, primiparous women were more likely than multiparous women to report being admitted with hypertension related to pregnancy or with early labor. Otherwise, conditions leading to hospitalization were similar in the two groups.

[Figure V.3](#) displays indicators of several key aspects of prenatal care from the CRRHS.

Figure V.3
Selected Indicators and Aspects of Prenatal Care
During Respondents' Most Recent Pregnancy



Smoking during Pregnancy

Cigarette smoking during pregnancy has been demonstrated to be a risk factor for adverse pregnancy outcomes, particularly for low birthweight. Sixty-eight percent of respondents with recent live births were not smokers when they became pregnant ([Table V.17](#)). An additional 21% reported that they stopped smoking upon finding out they were pregnant, with the remaining 11% continuing to smoke during pregnancy. About one-third of this group smoked at least 10 cigarettes per day. Women in Bohemia (and particularly women in North Bohemia and Prague) were much more likely than women in Moravia to smoke during pregnancy (14% and 8%, respectively). As educational level increased, the likelihood that a woman smoked during pregnancy decreased sharply, from 32% among women with only a primary school education, to 2% for those who attended university.

Birthweight/Pregnancy Outcomes


The incidence of low birthweight (LBW) (i.e., under 2,500 grams) among recent births was 5.8% among survey respondents ([Table V.18](#)). This is the same as the official rate of 5.8% for 1991 derived from national statistics, evidence that the survey data are of high quality in this regard. As is usually the case, the incidence of LBW was higher among primiparous women (7%) than among multiparous women (5%). Although this difference was not large, the LBW rate was lower among multiparous women in all but a few categories of survey respondents. The highest LBW rates were found in cities of at least 20,000 population, in North Moravia, among the oldest (40-44 year-old) and youngest (15-19 year-old), single, and poorly educated women. The incidence of low birthweight exceeded 6% only in North Moravia and Prague ([Figure V.4](#)).

By excluding unintended pregnancies from our tabulations, we were able to examine the distributions of pregnancy outcomes for just those pregnancies that women planned to have. This allows us to examine pregnancy outcomes from a physiological perspective (i.e., without having to worry about how unintended pregnancies end). Eighty-seven percent of all intended pregnancies since the beginning of 1988 resulted in live births ([Table V.19](#)). The only noteworthy differences observed according to place of residence, age, or education were lower proportions of live births and higher proportions of miscarriages among 15-19 year-olds and 35-44 year-olds. The induced abortion proportion was also higher among 40-44 year-olds, but it is possible this is a result of slight misreporting of intendedness of pregnancies among these women.

Labor and Delivery

The CRRHS questionnaire also inquired into respondents' opinions regarding services received for recent pregnancies and deliveries. Women were asked their impressions about the facility in which they most recently delivered in regard to: distance from home, physical facilities, crowdedness, attentiveness of staff, and competence of staff

Figure V.4
Percent Babies with Low Birthweight, by Region
1993 CRRHS



([Table V.20](#)). Only small differences were noted according to parity. Each of the five characteristics was rated as "good" by between one-half and three-quarters of respondents. The characteristic eliciting the most "poor" ratings (20%) was crowdedness. About one in ten women were unhappy about the distance from home, the physical facilities, and the attentiveness of the staff. The aspect of care about which respondents were clearly the most satisfied was the competence of the staff. The one area where satisfaction could most easily be improved is attentiveness of staff. The results indicate a need for a better and differentiated approach of the professional staff toward patients.

[Table V.21](#) reveals that for 41% of respondents labor and delivery lasted less time than they expected, while for 27% it lasted longer than expected. Multiparous women were more likely than primiparous women to have labor last shorter than they expected, probably a result of typically longer time of labor for first deliveries.

About one-third of women experienced more pain during labor and delivery than they anticipated ([Table V.22](#)). Even among women who had delivered previously, 31% experienced more pain than expected. There was a slight decrease in those reporting more pain than expected as education increased. The fact that for so many women the labor and delivery experience was very different than they expected indicates a need for better preparation and education of women during the prenatal period.

Overall, four-fifths of respondents said that they received adequate information regarding their pregnancy ([Table V.23](#)). Not surprisingly, multiparous women (who had previously gone through a full-term pregnancy) were somewhat more satisfied. Women from small communities tended to be more satisfied than women from larger cities, and older women were more likely to be satisfied than younger ones. There was no clear relationship between education and whether women thought they received enough information.

The level of satisfaction with information provided about delivery was 68%, lower than for information about pregnancy ([Table V.24](#)). Among multiparae 72% were satisfied, compared with 61% of primiparae. As with pregnancy information, satisfaction tended to decrease with increasing size of place and increase with age. Satisfaction with information also was lowest for the most educated women, regardless of parity.

Respondents who had a live birth since the beginning of 1987 were asked who they would like to be present and if they would be interested in home delivery, should they have another child. Although no figures are currently available, it is still relatively rare for a husband/partner to be in attendance at a woman's labor and delivery. Thirty-nine percent of women said they would like to have their partner present at their next delivery, while 59% preferred to have only the professional staff present ([Table V.25](#)). There was a very sizable difference in preferences according to respondents' ages, whereby the percentage wanting only staff members in attendance rose from 33% for 15-19 year-olds to 83% for 40-44 year-olds. Poorly educated women were somewhat more

likely than others to want only staff members present.

Relatively few women, regardless of characteristics, were interested in having a baby at home, rather than in a health facility ([Table V.26](#)). Eighty-eight percent of respondents were definitely not interested in home delivery, while 4% showed interest. There was no strong relationship apparent between interest in home delivery and the personal characteristics examined.

Problems in the Post Partum Period

When asked what their major problem was during the first week following delivery, 40% of respondents said they had no major problems ([Table V.27](#) and [Figure V.5](#)). The problems most commonly mentioned related to the health and care of the baby (18%), the woman's own health (17%), and breastfeeding (11%). Those reporting they had no major problems were more likely to live in small communities, be between the ages of 35 and 44, not to have attended university, and to have had more than one child. Care of the child was mentioned most commonly by 15-19 and 40-44 year-old women and by women who had never been married. Breastfeeding concerns stood out as a problem among university educated and primiparous women, but were rarely mentioned by 35-44 year-old women.

When asked about their most important problem after returning home from the hospital, 58% of respondents said they had no major problems ([Table V.28](#) and [Figure V.6](#)). The problems mentioned most frequently were her own health (10%), health of the child (8%), care of her child/children (7%), and home environment (5%). Women from small communities, 40-44 year-olds, and women who were multiparous were the most likely not to mention any problems. Their own health was most often mentioned by women from larger communities and women 40-44 years of age. Health of the child was a major concern among the youngest women and single women. Care of the child was most commonly mentioned by university educated and primiparous women, but rarely by multiparous women.

Prenatal Counselling

Overall, 24% of respondents attended prenatal counselling sessions before their last live birth and more than half of those attended fewer than four sessions ([Table V.29](#)). Attendance was higher in Moravia than in Bohemia, and was especially high in South Moravia. Women with only a primary school education were less likely than others to attend prenatal counselling. Primiparae were more likely to attend counselling and attended more sessions than multiparae.

Breastfeeding

Breastfeeding has long been known to have two important effects related to reproductive

Figure V.5
Most Important Problem for Respondent
During the Week Following Her Most Recent Birth
Percent Distribution, 1993 CRRHS





Figure V.6
Most Important Problem for Respondent
after Returning Home Following Delivery

Percent Distribution, 1993 CRRHS



health. First, by increasing exposure to maternal antibodies, by decreasing exposure to infection, and by improving nutrition, breastfeeding reduces morbidity and mortality among infants. Second, by delaying the return of ovulation, breastfeeding decreases the probability of conception, especially in the absence of use of effective means of contraception. Even though the magnitude of these effects are much less in developed countries like the Czech Republic than they tend to be in developing countries, the benefits of breastfeeding can still be substantial.

Ninety-one percent of children born to respondents since the beginning of 1988 were breastfed for at least some period of time ([Table V.30](#)). The proportion of infants breastfed was highest for those whose mothers: were ages 25-29 at the time of interview (94%); had a secondary school diploma (94%) or had attended university (92%); or were Catholic (93%). The proportion breastfed was lowest for infants with mothers ages 15-19 (84%) or ages 40-44 (71 %).

Despite the fact that relatively few infants were never breastfed, the mean duration of breastfeeding was short: 3.7 months for all children and 4.2 months for children who were reportedly breastfed ([Table V.31](#)). Although the survey data do not allow estimation of the duration of postpartum amenorrhea, it is unlikely that the short duration of breastfeeding causes more than a negligible reduction in rates of conception for the overall population. Mean durations of breastfeeding were longer among children of women with at least a secondary diploma than women without a diploma and were slightly longer among women ages 25-44 at the time of interview.

TABLE V.1
 Respondents' Reaction to Most Recent Pregnancy*, by Selected Characteristics
 1993 Czech Republic Reproductive Health Survey
 (Percent Distributions)

Characteristics	Respondents' Reaction to Pregnancy				Total	No. of Women
	Happy Immed.	Accepted Easily	Accepted Eventually	Other		
Total	66.0	25.3	8.1	0.6	100.0	(1382)
Region						
Bohemia	65.2	26.5	7.4	0.9	100.0	(855)
Moravia	67.2	23.4	9.2	0.2	100.0	(527)
Size of Place						
< 5000	67.1	25.7	6.8	0.4	100.0	(460)
5000-19999	65.7	26.6	6.7	1.0	100.0	(300)
20000 +	65.3	24.3	9.8	0.6	100.0	(622)
Age						
15-19	62.3	24.6	11.5	1.6	100.0	(51)
20-24	62.8	31.1	5.7	0.5	100.0	(418)
25-29	70.8	21.1	7.7	0.4	100.0	(533)
30-34	65.6	25.7	8.3	0.4	100.0	(249)
35-44	59.9	22.5	15.5	2.1	100.0	(131)
Education						
Primary	56.4	27.3	15.7	0.6	100.0	(162)
Secondary No Diploma	66.7	26.3	6.7	0.4	100.0	(531)
Secondary Diploma	68.5	23.7	7.3	0.5	100.0	(563)
Any University	64.6	25.2	7.9	2.4	100.0	(126)

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

TABLE V.2
Husband's or Partner's Reaction to Most Recent Pregnancy*, by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Husband's Reaction to Pregnancy					Total	No. of Women
	Happy Immed.	Accepted Easily	Accepted Eventually	Never Accepted	Don't Know**		
Total	64.9	23.5	8.1	2.4	1.1	100.0	(1382)
Region							
Bohemia	64.4	24.1	8.8	1.9	0.8	100.0	(855)
Moravia	65.6	22.6	7.1	3.1	1.6	100.0	(527)
Size of Place							
< 5000	67.9	23.8	5.5	1.5	1.2	100.0	(460)
5000-19999	64.1	25.0	7.4	2.9	0.6	100.0	(300)
20000 +	63.0	22.6	10.4	2.8	1.2	100.0	(622)
Age							
15-19	55.7	18.0	11.5	11.5	3.3	100.0	(51)
20-24	65.8	23.4	6.1	3.9	0.9	100.0	(418)
25-29	68.6	24.3	5.4	0.7	0.9	100.0	(533)
30-34	60.9	25.3	11.1	1.6	1.2	100.0	(249)
35-44	59.2	20.4	17.6	1.4	1.4	100.0	(131)
Education							
Primary	59.3	19.2	15.1	5.8	0.6	100.0	(162)
Secondary No Diploma	63.6	24.5	6.8	3.6	1.5	100.0	(531)
Secondary Diploma	67.3	23.5	7.6	0.7	0.9	100.0	(563)
Any University	66.9	25.2	6.3	0.0	1.6	100.0	(126)

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

**Includes two partners who never knew about the pregnancy.

TABLE V.3
Physical Work During Most Recent Pregnancy*, by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Amount of Physical Work				Total	No. of Women
	Much	Moderate	Little or None	No job		
Total	15.2	28.6	28.2	28.0	100.0	(1382)
Region						
Bohemia	15.3	29.8	28.7	26.1	100.0	(855)
Moravia	14.9	26.8	27.4	31.0	100.0	(527)
Size of Place						
< 5000	15.0	33.1	23.6	28.3	100.0	(460)
5000-19999	14.7	26.6	35.6	23.1	100.0	(300)
20000 +	15.5	26.3	28.0	30.2	100.0	(622)
Age						
15-19	19.7	26.2	21.3	32.8	100.0	(51)
20-24	16.1	29.7	23.8	30.4	100.0	(418)
25-29	12.9	27.1	31.0	29.0	100.0	(533)
30-34	13.4	30.4	29.2	26.9	100.0	(249)
35-44	21.8	28.9	32.4	16.9	100.0	(131)
Education						
Primary	22.7	30.2	12.2	34.9	100.0	(162)
Secondary No Diploma	21.1	34.6	16.0	28.3	100.0	(531)
Secondary Diploma	9.2	24.7	40.3	25.8	100.0	(563)
Any University	6.3	18.1	48.0	27.6	100.0	(126)

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

TABLE V.4
Amount of Standing during Work during Most Recent Pregnancy
by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Amount of Standing during Work				Total	No. of Women
	Much	Moderate	Little	No job		
Total	26.6	18.9	26.5	28.0	100.0	(1382)
Region						
Bohemia	27.4	19.8	26.7	26.1	100.0	(855)
Moravia	25.4	17.4	26.3	31.0	100.0	(527)
Size of Place						
< 5000	25.9	21.3	24.5	28.3	100.0	(460)
5000-19999	26.9	18.9	31.1	23.1	100.0	(300)
20000 +	26.9	17.0	25.9	30.2	100.0	(622)
Age						
15-19	34.4	13.1	19.7	32.8	100.0	(51)
20-24	26.1	18.8	24.7	30.4	100.0	(418)
25-29	24.3	19.3	27.5	29.0	100.0	(533)
30-34	26.5	19.4	27.3	26.9	100.0	(249)
35-44	33.8	19.0	30.3	16.9	100.0	(131)
Education						
Primary	32.6	16.9	15.7	34.9	100.0	(162)
Secondary No Diploma	37.7	15.7	18.4	28.3	100.0	(531)
Secondary Diploma	16.8	21.8	35.6	25.8	100.0	(563)
Any University	15.0	22.0	35.4	27.6	100.0	(126)

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

TABLE V.5
Amount of Stress at Work During Most Recent Pregnancy* by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Mental Stress During Work				Total	No. of Women
	Much	Moderate	Little	No Job		
Total	16.8	31.5	23.7	28.0	100.0	(1382)
Region						
Bohemia	16.4	33.0	24.5	26.1	100.0	(855)
Moravia	17.6	29.2	22.3	31.0	100.0	(527)
Size of Place						
< 5000	16.5	28.3	27.0	28.3	100.0	(460)
5000-19999	15.4	34.6	26.9	23.1	100.0	(300)
20000 +	17.8	32.4	19.7	30.2	100.0	(622)
Age						
15-19	11.5	31.1	24.6	32.8	100.0	(51)
20-24	13.8	30.8	24.9	30.4	100.0	(418)
25-29	16.8	31.8	22.4	29.0	100.0	(533)
30-34	19.8	33.6	19.8	26.9	100.0	(249)
35-44	23.2	28.9	31.0	16.9	100.0	(131)
Education						
Primary	12.2	22.1	30.8	34.9	100.0	(162)
Secondary No Diploma	11.4	31.5	28.8	28.3	100.0	(531)
Secondary Diploma	18.2	36.5	19.6	25.8	100.0	(563)
Any University	40.9	21.3	10.2	27.6	100.0	(126)

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

TABLE V.6
Trimester When Prenatal Care Began During Most Recent Pregnancy* by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Time of First Prenatal Care Visit			Total	No. of Women
	1 st Trimester	2 nd -3 rd Trimester	No Care		
Total	94.1	5.2	0.6	100.0	(1382)
Region					
Bohemia	93.5	5.6	0.9	100.0	(855)
Moravia	95.1	4.7	0.2	100.0	(527)
Prague	96.2	3.0	0.8	100.0	(128)
Central Bohemia	93.5	6.5	0.0	100.0	(162)
South Bohemia	97.1	2.9	0.0	100.0	(101)
West Bohemia	96.2	3.8	0.0	100.0	(125)
North Bohemia	90.2	7.6	2.2	100.0	(182)
East Bohemia	90.7	7.5	1.9	100.0	(157)
South Moravia	96.0	4.0	0.0	100.0	(258)
North Moravia	94.3	5.4	0.4	100.0	(269)
Size of Place					
< 5000	93.7	5.7	0.6	100.0	(460)
5000-19999	94.9	4.5	0.6	100.0	(300)
20000 +	94.1	5.3	0.6	100.0	(622)
Age					
15-19	91.8	8.2	0.0	100.0	(51)
20-24	93.7	5.4	0.9	100.0	(418)
25-29	96.3	3.0	0.7	100.0	(533)
30-34	93.7	5.9	0.4	100.0	(249)
35-44	89.4	10.6	0.0	100.0	(131)

TABLE V.6
Trimester When Prenatal Care Began During Most Recent Pregnancy* by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Time of First Prenatal Care Visit			Total	No. of Women
	1 st Trimester	2 nd -3 rd Trimester	No Care		
Marital Status					
Currently Married	94.5	4.9	0.6	100.0	(1274)
Previously Married	91.2	8.8	0.0	100.0	(64)
Single	89.1	9.1	1.8	100.0	(44)
Education					
Primary	87.2	11.6	1.2	100.0	(162)
Secondary No Diploma	94.2	4.9	0.9	100.0	(531)
Secondary Diploma	95.8	3.8	0.3	100.0	(563)
Any University	95.3	4.7	0.0	100.0	(126)
Religion					
None	94.7	4.6	0.6	100.0	(920)
Catholic	94.4	5.4	0.2	100.0	(396)
Other	83.3	13.6	3.0	100.0	(62)
Household Income					
0-3000	90.2	8.5	1.2	100.0	(80)
3001 - 7000	94.2	5.1	0.6	100.0	(759)
7001 - 10000	94.8	4.6	0.6	100.0	(322)
10001 +	94.9	5.1	0.0	100.0	(167)
Not Stated	91.9	6.5	1.6	100.0	(54)

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

(cont.)

TABLE V.7
 Number of Prenatal Care Visits During Most Recent Pregnancy*, by Selected Characteristics
 1993 Czech Republic Reproductive Health Survey
 (Percent Distributions)

Characteristics	Number of Prenatal Visits					Total	No. of Women
	0	1-5	6-9	10 +	Do Not Rememb.		
Total	0.6	0.7	5.4	91.4	1.8	100.0	(1382)
Region							
Bohemia	0.9	0.6	4.7	92.6	1.2	100.0	(855)
Moravia	0.2	0.9	6.7	89.5	2.7	100.0	(527)
Prague	0.8	0.0	1.5	97.0	0.8	100.0	(128)
Central Bohemia	0.0	0.0	7.7	92.3	0.0	100.0	(162)
South Bohemia	0.0	1.0	10.7	87.4	1.0	100.0	(101)
West Bohemia	0.0	1.5	6.1	89.4	3.0	100.0	(125)
North Bohemia	2.2	0.5	1.6	92.9	2.7	100.0	(182)
East Bohemia	1.9	0.6	2.5	95.0	0.0	100.0	(157)
South Moravia	0.0	0.4	8.8	87.9	2.9	100.0	(258)
North Moravia	0.4	1.4	4.6	91.1	2.5	100.0	(269)
Size of place							
< 5000	0.6	0.6	5.9	91.4	1.5	100.0	(460)
5000-19999	0.6	0.3	3.5	93.3	2.2	100.0	(300)
20000 +	0.6	0.9	6.0	90.6	1.9	100.0	(622)
Age Groups							
15-19	0.0	3.3	8.2	86.9	1.6	100.0	(51)
20-24	0.9	0.7	6.3	90.5	1.6	100.0	(418)
25-29	0.7	0.7	4.9	92.0	1.7	100.0	(533)
30-34	0.4	0.4	5.1	92.1	2.0	100.0	(249)
35-44	0.0	0.0	4.2	93.0	2.8	100.0	(131)

TABLE V.7
Number of Prenatal Care Visits During Most Recent Pregnancy*, by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Number of Prenatal Visits					Total	No. of Women
	0	1-5	6-9	10 +	Do Not Rememb.		
Marital Status							
Currently Married	0.6	0.8	5.5	91.5	1.6	100.0	(1274)
Previously Married	0.0	0.0	2.9	91.2	5.9	100.0	(64)
Single	1.8	0.0	7.3	89.1	1.8	100.0	(44)
Education							
Primary	1.2	1.2	6.4	89.0	2.3	100.0	(162)
Sec, No Diploma	0.9	0.9	5.0	91.9	1.3	100.0	(531)
Secondary, Diploma	0.3	0.5	5.9	91.2	2.1	100.0	(563)
Any University	0.0	0.0	3.9	93.7	2.4	100.0	(126)
Religion							
No Religion	0.6	0.7	5.6	91.7	1.4	100.0	(920)
Roman Catholic	0.2	0.7	5.4	91.5	2.2	100.0	(396)
Other	3.0	0.0	4.5	87.9	4.5	100.0	(62)
Household Income							
0-3000	1.2	0.0	8.5	86.6	3.7	100.0	(80)
3001-7000	0.6	0.9	6.3	90.5	1.7	100.0	(759)
7001-10000	0.6	0.9	4.3	93.3	0.9	100.0	(322)
10001 +	0.0	0.0	3.4	93.8	2.8	100.0	(167)
Not Stated	1.6	0.0	3.2	91.9	3.2	100.0	(54)

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

TABLE V.8
Principal Prenatal Care Provider during Most Recent Pregnancy* by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Prenatal Care Provider			Total	No. of Women
	Physician	Midwife/ Nurse	Both		
Total	62.6	2.7	34.6	100.0	(1374)
Region					
Prague	70.2	3.8	26.0	100.0	(127)
Central Bohemia	67.9	0.6	31.5	100.0	(162)
South Bohemia	57.3	1.9	40.8	100.0	(101)
West Bohemia	60.6	4.5	34.8	100.0	(125)
North Bohemia	54.4	3.9	41.7	100.0	(178)
East Bohemia	67.1	1.9	31.0	100.0	(155)
South Moravia	63.2	0.7	36.0	100.0	(258)
North Moravia	60.9	4.7	34.4	100.0	(268)
Size of Place					
< 5000	56.7	2.8	40.6	100.0	(457)
5000-19999	64.2	2.6	33.2	100.0	(299)
20000 +	66.2	2.8	31.0	100.0	(618)
Age					
15-19	42.6	3.3	54.1	100.0	(51)
20-24	64.5	2.5	33.0	100.0	(415)
25-29	65.7	2.4	31.8	100.0	(529)
30-34	55.2	2.8	42.1	100.0	(248)
35-44	66.9	4.2	28.9	100.0	(131)
Education					
Primary	48.8	2.9	48.2	100.0	(160)
Secondary No Diploma	62.2	3.1	34.7	100.0	(527)
Secondary Diploma	66.5	2.3	31.3	100.0	(561)
Any University	65.4	3.1	31.5	100.0	(126)

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

TABLE V.9
Principal Source of Prenatal Care during Most Recent Pregnancy* by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Place of Prenatal Care			Total	No. of Women
	District Office	District + Hospital	Hospital		
Total	59.9	34.6	5.6	100.0	(1374)
Region					
Prague	37.4	56.5	6.1	100.0	(127)
Central Bohemia	61.9	32.7	5.4	100.0	(162)
South Bohemia	58.3	38.8	2.9	100.0	(101)
West Bohemia	68.9	24.2	6.8	100.0	(125)
North Bohemia	61.7	33.3	5.0	100.0	(178)
East Bohemia	57.6	35.4	7.0	100.0	(155)
South Moravia	60.7	33.8	5.5	100.0	(258)
North Moravia	64.9	29.7	5.4	100.0	(268)
Size of Place					
< 5000	65.2	29.9	4.9	100.0	(457)
5000-19999	63.2	32.3	4.5	100.0	(299)
20000 +	54.4	39.1	6.5	100.0	(618)
Age					
15-19	59.0	39.3	1.6	100.0	(51)
20-24	58.4	36.8	4.8	100.0	(415)
25-29	63.5	31.3	5.3	100.0	(529)
30-34	57.1	36.9	6.0	100.0	(248)
35-44	56.3	33.8	9.9	100.0	(131)
Education					
Primary	69.4	25.9	4.7	100.0	(160)
Secondary No Diploma	62.7	34.5	2.7	100.0	(527)
Secondary Diploma	54.7	36.1	9.2	100.0	(561)
Any University	58.3	39.4	2.4	100.0	(126)

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

TABLE V.10
Percent Distribution of Time Typically Spent Travelling
and Mean Time Spent Travelling to Usual Source of Prenatal Care*,
by Means of Transportation
1993 Czech Republic Reproductive Health Survey

Means of Transport	Time Typically Spent Travelling				Mean Time (minutes)	Number of Women
	≤ 30 Minutes	31-60 Minutes	> 60 Minutes	Total		
Walk	97.0	2.8	0.1	100.0	15.1	(676)
Bus/Metro	76.1	20.9	2.9	100.0	30.1	(533)
Private Vehicle	89.6	9.0	1.4	100.0	24.5	(140)
Total	88.2	10.5	1.3	100.0	21.8	(1361)**

* Limited to the most recent pregnancy leading to a live birth since December 1987.

**Includes 12 women who used other means of transportation.

TABLE V.11
Waiting Time for Prenatal Care during Most Recent Pregnancy* by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Usual Waiting Time at Source of Prenatal Care					Total	No. of Women
	≤15 Min	16-30 Min	31-60 Min	> 60 Min	Do Not Remem.		
Total	20.8	34.6	29.0	14.8	0.8	100.0	(1365)
Region							
Prague	30.5	31.3	23.7	13.7	0.8	100.0	(127)
Central Bohemia	17.4	31.7	33.5	16.8	0.6	100.0	(161)
South Bohemia	15.5	41.7	31.1	10.7	1.0	100.0	(101)
West Bohemia	15.3	32.1	32.8	19.8	0.0	100.0	(124)
North Bohemia	22.3	31.8	30.7	14.5	0.6	100.0	(177)
East Bohemia	21.5	31.0	30.4	16.5	0.6	100.0	(155)
South Moravia	25.0	36.6	22.0	14.9	1.5	100.0	(254)
North Moravia	17.3	38.3	31.0	12.3	1.1	100.0	(266)
Size of Place							
< 5000	19.2	36.3	26.9	16.2	1.3	100.0	(454)
5000-19999	18.0	29.2	33.4	18.4	1.0	100.0	(294)
20000 +	23.2	35.9	28.4	12.0	0.5	100.0	(617)
Age							
15-19	9.8	41.0	34.4	14.8	0.0	100.0	(51)
20-24	18.6	36.1	29.0	15.6	0.7	100.0	(413)
25-29	22.7	33.6	28.7	14.2	0.8	100.0	(527)
30-34	21.3	32.5	30.5	14.9	0.8	100.0	(245)
35-44	24.3	34.3	25.0	14.3	2.1	100.0	(129)
Education							
Primary	17.9	35.7	32.7	13.1	0.6	100.0	(158)
Secondary No Diploma	16.8	34.1	34.5	13.5	1.1	100.0	(525)
Secondary Diploma	23.3	35.4	24.0	16.8	0.5	100.0	(556)
Any University	30.7	31.5	22.8	13.4	1.6	100.0	(126)

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

TABLE V.12
 Ultrasound Examination during Most Recent Pregnancy* by Selected Characteristics
 1993 Czech Republic Reproductive Health Survey
 (Percent Distribution)

Characteristics	Whether Had Ultrasound Examination			Total	No. of Women
	Yes	No	Do Not Remember		
Total	91.6	8.0	0.4	100.0	(1374)
Region					
Prague	94.7	4.6	0.8	100.0	(127)
Central Bohemia	95.2	3.0	1.8	100.0	(162)
South Bohemia	98.1	1.9	0.0	100.0	(101)
West Bohemia	86.4	13.6	0.0	100.0	(125)
North Bohemia	88.9	10.6	0.6	100.0	(178)
East Bohemia	90.5	9.5	0.0	100.0	(155)
South Moravia	97.1	2.9	0.0	100.0	(258)
North Moravia	85.3	14.7	0.0	100.0	(268)
Size of Place					
< 5000	91.7	8.3	0.0	100.0	(457)
5000-19999	89.0	11.0	0.0	100.0	(299)
20000 +	92.8	6.4	0.8	100.0	(618)
Age					
15-19	96.7	1.6	1.6	100.0	(51)
20-24	94.3	5.7	0.0	100.0	(415)
25-29	90.2	9.4	0.4	100.0	(529)
30-34	90.1	9.5	0.4	100.0	(248)
35-39	87.6	11.5	0.9	100.0	(105)
40-44	96.6	3.4	0.0	100.0	(26)
Education					
Primary	88.8	10.0	1.2	100.0	(160)
Secondary No Diploma	91.1	8.5	0.4	100.0	(527)
Secondary Diploma	91.8	8.0	0.2	100.0	(561)
Any University	96.9	3.1	0.0	100.0	(126)

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

TABLE V.13
Degree to Which the Purpose of Ultrasound Examination* Was Explained
by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Degree of Explanation about Ultrasound				Total	No. of Women
	Well	A little	Not at All	Do Not Remember		
Total	54.7	18.3	25.4	1.6	100.0	(1258)
Region						
Prague	48.4	20.2	29.8	1.6	100.0	(120)
Central Bohemia	59.4	15.6	23.1	1.9	100.0	(154)
South Bohemia	66.3	10.9	20.8	2.0	100.0	(99)
West Bohemia	56.1	25.4	18.4	0.0	100.0	(108)
North Bohemia	51.9	20.6	25.0	2.5	100.0	(158)
East Bohemia	41.3	25.2	32.2	1.4	100.0	(141)
South Moravia	59.5	14.0	24.2	2.3	100.0	(250)
North Moravia	53.8	18.1	27.3	0.8	100.0	(228)
Size of Place						
< 5000	58.8	19.2	20.6	1.4	100.0	(419)
5000-19999	54.0	17.8	27.2	1.1	100.0	(266)
20000 +	52.0	18.0	28.0	2.0	100.0	(573)
Age						
15-19	47.5	23.7	27.1	1.7	100.0	(49)
20-24	51.2	19.9	26.7	2.2	100.0	(391)
25-29	56.4	18.8	24.4	0.4	100.0	(477)
30-34	51.5	17.6	29.1	1.8	100.0	(224)
35-39	66.7	10.1	19.2	4.0	100.0	(92)
40-44	75.0	10.7	10.7	3.6	100.0	(25)
Education						
Primary	49.0	22.5	23.8	4.6	100.0	(141)
Secondary No Diploma	53.5	20.2	24.8	1.6	100.0	(479)
Secondary Diploma	57.1	15.9	26.3	0.8	100.0	(516)
Any University	56.1	16.3	26.0	1.6	100.0	(122)
Parity						
Primiparae	51.5	20.7	25.6	2.2	100.0	(521)
Multiparae	57.0	16.6	25.2	1.2	100.0	(737)

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

TABLE V.14
Prescribed Bedrest during Most Recent Pregnancy* and Length of Bedrest
by Selected Characteristics
1993 Czech Republic Reproductive Health Survey

Characteristics	% with Prescribed Bedrest	Length of Bedrest (in weeks)		No. of Women
		Mean	Median	
Total	29.1	6.8	4.0	(1382)
Region				
Prague	29.5	5.5	4.0	(128)
Central Bohemia	23.8	6.2	4.0	(162)
South Bohemia	25.2	5.9	3.5	(101)
West Bohemia	22.7	6.8	3.0	(125)
North Bohemia	38.0	5.7	3.0	(182)
East Bohemia	27.3	7.8	4.5	(157)
South Moravia	32.7	7.9	5.0	(258)
North Moravia	28.2	7.0	4.0	(269)
Size of Place				
< 5000	29.5	6.2	4.0	(460)
5000-19999	27.9	6.7	4.0	(300)
20000 +	29.4	7.2	4.0	(622)
Age				
15-19	29.5	5.2	3.0	(51)
20-24	27.9	5.9	3.0	(418)
25-29	30.5	6.7	4.0	(533)
30-34	29.2	7.4	4.0	(249)
35-39	27.4	8.2	4.0	(105)
40-44	27.6	11.9	12.0	(26)
Education				
Primary	25.0	5.4	3.0	(162)
Secondary No Diploma	25.8	7.1	4.0	(531)
Secondary Diploma	32.0	6.8	4.0	(563)
Any University	36.2	7.0	4.0	(126)

*Limited to ultrasound during the most recent pregnancy resulting in a live birth after December 1987

TABLE V.15
Hospitalization during Most Recent Pregnancy* and Length of Stay
by Selected Characteristics
1993 Czech Republic Reproductive Health Survey

Characteristics	% Hospitalized	Length of Stay (in days)		No. of Women
		Mean	Median	
Total	28.9	27.0	14.0	(1382)
Region				
Prague	28.8	27.1	14.0	(128)
Central Bohemia	30.4	20.4	14.0	(162)
South Bohemia	31.1	26.7	14.0	(101)
West Bohemia	22.7	19.8	14.0	(125)
North Bohemia	31.5	27.6	14.0	(182)
East Bohemia	29.2	28.0	15.0	(157)
South Moravia	27.9	25.6	14.0	(258)
North Moravia	29.3	34.3	14.5	(269)
Size of Place				
< 5000	28.9	29.4	17.5	(460)
5000-19999	30.1	23.1	14.0	(300)
20000 +	28.3	27.2	14.0	(622)
Age				
15-19	34.4	22.3	14.0	(51)
20-24	27.9	22.5	14.0	(418)
25-29	30.7	26.7	18.0	(533)
30-34	26.1	31.7	14.0	(249)
35-39	29.2	32.5	14.0	(105)
40-44	24.1	57.1	21.0	(26)
Education				
Primary	22.7	21.3	14.0	(162)
Secondary No Diploma	28.1	26.4	14.0	(531)
Secondary Diploma	31.3	28.8	18.0	(563)
Any University	29.9	27.0	14.0	(126)
Parity				
Primiparae	31.6	25.4	14.0	(561)
Multiparae	27.0	28.4	14.0	(821)

*Limited to the most recent pregnancy leading to a live birth after December 1987.

TABLE V.16
Percent of Women Hospitalized during Most Recent Pregnancy*
for Whom Various Conditions Were Diagnosed, by Parity
1993 Czech Republic Reproductive Health Survey

Reason for Hospitalization	Percent of Hospitalized Women		
	Total	Primiparae	Multiparae
Bleeding in First Half of Pregnancy	39.1	38.5	39.7
Early/Threatened/False Labor	32.9	35.3	30.8
Swelling/Edema	13.8	14.4	13.2
Hypertension Related to Pregnancy	11.1	16.6	6.6
Bleeding in Second Half of Pregnancy	10.6	11.2	10.1
Improper Fetal Position	7.0	7.0	7.1
Renal/Urinary Tract Problem	4.1	4.3	4.0
Previous Caesarean Section	3.6	1.1	5.7
Hypertension before Pregnancy	2.4	3.2	1.8
Diabetes	1.7	2.1	1.3
Low Weight Gain/IUGR	1.7	2.7	0.9
Twins	1.2	0.5	1.8
Other Conditions	17.6	18.2	17.2
<i>Number of Hospitalized Women</i>	<i>(397)</i>	<i>(174)</i>	<i>(223)</i>

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

NOTE: Columns add to more than 100%, since some women had more than one condition.

TABLE V.17
Smoking Status of Women during Most Recent Pregnancy* by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Smoking Status During Pregnancy						No. of Women
	Did Not Smoke Before	Stopped Smoking in Pregnancy	Smoked < 5/Day	Smoked 5-9/Day	Smoked 10 + /Day	Smoked DR How Much	
Total	67.6	20.9	4.5	4.4	2.3	0.3	100.0 (1382)
Region							
Bohemia	65.2	20.8	5.1	5.5	3.3	0.1	100.0 (855)
Moravia	71.4	21.0	3.6	2.7	0.7	0.5	100.0 (527)
Size of Place							
< 5000	71.3	20.0	2.7	4.4	1.3	0.2	100.0 (460)
5000-19999	74.4	18.3	3.2	1.9	1.3	1.0	100.0 (300)
20000 +	61.6	22.8	6.5	5.6	3.6	0.0	100.0 (622)
Age							
15-19	67.2	21.3	3.3	3.3	4.9	0.0	100.0 (51)
20-24	63.3	25.6	4.3	4.8	1.8	0.2	100.0 (418)
25-29	68.0	20.0	4.9	4.3	2.6	0.2	100.0 (533)
30-34	74.3	15.8	4.7	2.8	2.0	0.4	100.0 (249)
35-39	71.7	18.6	3.5	5.3	0.0	0.9	100.0 (105)
40-44	51.7	17.2	6.9	13.8	10.3	0.0	100.0 (26)

TABLE V.17
Smoking Status of Women during Most Recent Pregnancy* by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Smoking Status During Pregnancy						No. of Women
	Did Not Smoke Before	Stopped Smoking in Pregnancy	Smoked <5/Day	Smoked 5-9/Day	Smoked 10 +/Day	Smoked DR How Much	
Marital Status							
Currently Married	68.8	20.5	4.2	4.1	2.1	0.3	100.0 (1274)
Previously Married	61.8	23.5	4.4	5.9	4.4	0.0	100.0 (64)
Single	45.5	27.3	12.7	9.1	5.5	0.0	100.0 (44)
Education							
Primary	44.2	23.8	6.4	14.0	9.9	1.7	100.0 (162)
Secondary No Diploma	61.8	24.7	5.4	5.6	2.3	0.2	100.0 (531)
Secondary Diploma	76.0	18.5	3.8	1.4	0.3	0.0	100.0 (563)
Any University	86.6	11.0	1.6	0.0	0.8	0.0	100.0 (126)
Parity							
Primiparae	68.1	22.1	4.2	3.7	1.7	0.2	100.0 (561)
Multiparae	67.3	20.0	4.8	4.9	2.7	0.4	100.0 (821)

* Limited to the most recent pregnancy resulting in a live birth after December 1987

TABLE V.18
Percent Babies*with Low Birthweight (under 2,500 grams) by Parity
1993 Czech Republic Reproductive Health Survey

Characteristics	TOTAL		PRIMIPARAE		MULTIPARAE	
	% LBW	No. of Babies	% LBW	No. of Babies	% LBW	No. of Babies
Total	5.8	(1382)	7.1	(561)	4.9	(821)
Region						
Bohemia	5.1	(855)	6.8	(368)	3.8	(487)
Moravia	6.9	(527)	7.7	(193)	6.4	(334)
Prague	6.8	(128)	12.5	(61)	1.5	(67)
Central Bohemia	3.0	(162)	3.8	(74)	2.2	(88)
South Bohemia	5.8	(101)	10.5	(37)	3.1	(64)
West Bohemia	4.5	(125)	3.1	(59)	5.9	(66)
North Bohemia	6.0	(182)	8.6	(70)	4.4	(112)
East Bohemia	5.0	(157)	4.4	(67)	5.4	(90)
South Moravia	4.4	(258)	6.5	(99)	3.0	(159)
North Moravia	9.3	(269)	8.9	(94)	9.5	(175)
Size of Place						
< 5000	5.3	(460)	6.7	(171)	4.4	(289)
5000-19999	2.9	(300)	2.2	(127)	3.4	(173)
20000 +	7.6	(622)	9.7	(263)	6.0	(359)

TABLE V.18
Percent Babies* with Low Birthweight (under 2,500 grams) by Parity
1993 Czech Republic Reproductive Health Survey

Characteristics	TOTAL		PRIMIPARAE		MULTIPARAE	
	% LBW	No. of Babies	% LBW	No. of Babies	% LBW	No. of Babies
Age						
15-19	14.8	(51)	15.3	(49)	**	(2)
20-24	6.3	(418)	6.2	(287)	6.7	(131)
25-29	4.3	(533)	5.1	(174)	3.9	(359)
30-34	4.7	(249)	8.1	(37)	4.2	(212)
35-39	4.4	(105)	**	(12)	3.0	(93)
40-44	20.7	(26)	**	(2)	**	(24)
Marital Status						
Currently Married	5.5	(1274)	6.5	(490)	4.9	(784)
Previously Married	2.9	(64)	3.1	(30)	2.8	(34)
Single	16.4	(44)	15.4	(41)	**	(3)
Education						
Primary	8.1	(162)	8.9	(41)	7.9	(121)
Secondary No Diploma	6.3	(531)	8.3	(215)	4.9	(316)
Secondary Diploma	5.0	(563)	6.3	(244)	4.0	(319)
Any University	3.9	(126)	4.8	(61)	3.1	(65)

* Limited to the most recent pregnancy resulting in a live birth after December 1987

** Fewer than 25 live births

TABLE V.19
Outcomes of Intended Pregnancies Ending since January 1988, by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Outcome of Pregnancy				Number of Pregnancies
	Live Birth	Stillbirth	Spontaneous Abortion	Induced Abortion	
Total	86.8	0.5	9.1	3.6	100.0 (1,638)
Region					
Bohemia	86.3	0.4	8.9	4.4	100.0 (998)
Moravia	87.7	0.8	9.3	2.3	100.0 (640)
Age					
15-19	84.1	2.3	13.6	0.0	100.0 (34)
20-24	89.3	0.4	7.2	3.1	100.0 (492)
25-29	87.2	0.7	9.2	2.8	100.0 (681)
30-34	86.6	0.4	7.8	5.3	100.0 (281)
35-39	80.3	0.0	15.9	3.8	100.0 (121)
40-44	72.7	0.0	12.1	15.2	100.0 (29)
Education					
Primary Only	87.3	1.1	9.5	2.1	100.0 (183)
Secondary, No Diploma	87.8	0.5	7.0	4.8	100.0 (633)
Secondary, Diploma	85.8	0.6	10.7	2.9	100.0 (678)
Any University	86.9	0.0	10.3	2.8	100.0 (144)

TABLE V.20
Assessment of Characteristics of Place of Delivery* by Parity
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	PRIMIPARAE					MULTIPARAE				
	Opinion about the place of delivery					Opinion about the place of delivery				
	Good	Fair	Poor	Don't Know	Total	Good	Fair	Poor	Don't Know	Total
Distance from home	69.9	21.5	8.4	0.2	100.0	65.7	23.6	10.6	0.1	100.0
Physical facilities	55.1	32.6	11.1	1.2	100.0	57.0	32.4	9.4	1.2	100.0
Crowdedness	53.9	26.5	19.6	0.0	100.0	52.6	26.4	20.7	0.2	100.0
Helpfulness/Attentiveness of staff	59.8	29.4	10.8	0.0	100.0	65.0	26.5	8.3	0.1	100.0
Competence of staff	72.8	21.5	5.1	0.7	100.0	76.5	19.3	3.5	0.7	100.0

Number of Respondents = 561

Number of Respondents = 821

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

TABLE V.21
Length of Delivery* Relative to Expected Length by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Length of Delivery				Total	No. of Women
	Shorter Than Expected	As Long as Expected	Longer Than Expected	Planned Cesarean Section		
Total	41.1	27.2	27.1	4.7	100.0	(1382)
Region						
Bohemia	39.2	30.8	25.8	4.2	100.0	(855)
Moravia	44.0	21.4	29.2	5.4	100.0	(527)
Age						
15-19	44.3	29.5	23.0	3.3	100.0	(51)
20-24	41.5	24.7	29.7	4.1	100.0	(418)
25-29	40.4	28.8	26.7	4.1	100.0	(533)
30-34	44.3	25.3	23.7	6.7	100.0	(249)
35-44	35.2	31.0	28.2	5.6	100.0	(131)
Education						
Primary	41.3	23.8	31.4	3.5	100.0	(162)
Secondary No Diploma	40.4	26.7	27.6	5.4	100.0	(531)
Secondary Diploma	41.0	29.1	26.6	3.3	100.0	(563)
Any University	44.1	25.2	21.3	9.4	100.0	(126)
Parity						
Primiparae	34.5	28.5	31.8	5.2	100.0	(561)
Multiparae	45.7	26.2	23.8	4.3	100.0	(821)

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

TABLE V.22
Pain during Labor* Relative to Expected Pain by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Pain during Labor				Total	No. of Women
	Less Than Expected	As Much as Expected	More Than Expected	Planned Cesarean Section		
Total	20.5	40.5	34.3	4.7	100.0	(1382)
Region						
Bohemia	20.8	41.5	33.5	4.2	100.0	(855)
Moravia	20.1	38.9	35.5	5.4	100.0	(527)
Age						
15-19	18.0	37.7	41.0	3.3	100.0	(51)
20-24	20.2	35.8	39.9	4.1	100.0	(418)
25-29	19.3	43.9	32.7	4.1	100.0	(533)
30-34	24.5	43.1	25.7	6.7	100.0	(249)
35-44	20.4	38.7	35.2	5.6	100.0	(131)
Education						
Primary	23.3	33.7	39.5	3.5	100.0	(162)
Secondary No Diploma	20.5	39.6	34.4	5.4	100.0	(531)
Secondary Diploma	19.7	43.4	33.6	3.3	100.0	(563)
Any University	20.5	40.2	29.9	9.4	100.0	(126)
Parity						
Primiparae	18.8	36.5	39.5	5.2	100.0	(561)
Multiparae	21.8	43.3	30.6	4.3	100.0	(821)

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

TABLE V.23
Percent of Respondents Who Thought They Received Enough Information on Pregnancy*
by Selected Characteristics by Parity
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	TOTAL		PRIMIPARAE		MULTIPARAE	
	%	No. of Women	%	No. of Women	%	No. of Women
Total	79.9	(1382)	75.7	(561)	82.9	(821)
Region						
Bohemia	81.0	(855)	77.3	(368)	83.9	(487)
Moravia	78.1	(527)	72.7	(193)	81.3	(334)
Size of Place						
< 5000	84.0	(460)	78.8	(171)	87.1	(289)
5000-19999	80.4	(300)	78.4	(127)	82.0	(173)
20000 +	76.6	(622)	72.4	(263)	79.8	(359)
Age						
15-19	78.7	(51)	78.0	(49)	**	(2)
20-24	78.0	(418)	77.2	(287)	79.9	(131)
25-29	77.6	(533)	70.3	(174)	81.1	(359)
30-34	81.4	(249)	78.4	(37)	81.9	(212)
35-44	92.3	(131)	**	(14)	92.2	(117)
Education						
Primary	78.5	(162)	71.1	(41)	81.1	(121)
Secondary No Diploma	83.1	(531)	78.6	(215)	86.2	(316)
Secondary Diploma	78.7	(563)	76.2	(244)	80.7	(319)
Any University	73.2	(126)	66.1	(61)	80.0	(65)

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

**Fewer than 25 respondents.

TABLE V.24
Percent of Respondents Who Thought They Received Enough Information on Delivery*
by Selected Characteristics by Parity
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	TOTAL		PRIMIPARAE		MULTIPARAE	
	%	No. of Women	%	No. of Women	%	No. of Women
Total	67.5	(1382)	60.6	(561)	72.3	(821)
Region						
Bohemia	66.3	(855)	59.5	(368)	71.4	(487)
Moravia	69.4	(527)	62.7	(193)	73.5	(334)
Size of Place						
< 5000	74.9	(460)	65.4	(171)	80.7	(289)
5000-19999	63.1	(300)	56.0	(127)	68.5	(173)
20000 +	64.1	(622)	59.9	(263)	67.3	(359)
Age						
15-19	55.7	(51)	57.6	(49)	**	(2)
20-24	53.0	(418)	61.2	(287)	67.2	(131)
25-29	67.7	(533)	61.7	(174)	70.6	(359)
30-34	68.8	(249)	51.4	(37)	71.8	(212)
35-44	83.1	(131)	**	(14)	84.4	(117)
Education						
Primary	72.1	(162)	68.9	(41)	73.2	(121)
Secondary No Diploma	69.7	(531)	61.6	(215)	75.5	(316)
Secondary Diploma	67.1	(563)	61.3	(244)	71.7	(319)
Any University	52.8	(126)	48.4	(61)	56.9	(65)

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

**Fewer than 25 respondents.

TABLE V.25
 Person Desired to Be Present at the Next Delivery by Selected Characteristics
 Women with a Live Birth after December 1987
 1993 Czech Republic Reproductive Health Survey
 (Percent Distribution)

Characteristics	Person Desired to Be Present			Total	No. of Women
	Husband/ Partner	Mother/ Other Person	Professional Staff Only		
Total	38.8	2.6	58.6	100.0	(1382)
Region					
Bohemia	39.1	2.5	58.4	100.0	(855)
Moravia	38.4	2.7	58.9	100.0	(527)
Size of Place					
< 5000	35.4	1.7	62.9	100.0	(460)
5000-19999	39.7	3.8	56.4	100.0	(300)
20000 +	40.9	2.6	56.5	100.0	(622)
Age					
15-19	62.3	4.9	32.8	100.0	(51)
20-24	44.2	2.7	53.1	100.0	(418)
25-29	38.1	2.7	59.3	100.0	(533)
30-34	34.8	2.0	63.2	100.0	(249)
35-44	21.8	2.1	76.1	100.0	(131)
Education					
Primary	28.5	2.9	68.6	100.0	(162)
Secondary No Diploma	38.4	2.5	59.1	100.0	(531)
Secondary Diploma	41.9	2.7	55.4	100.0	(563)
Any University	40.9	1.6	57.5	100.0	(126)
Parity					
Primiparae	43.8	3.2	53.0	100.0	(561)
Multiparae	35.4	2.2	62.5	100.0	(821)

TABLE V.26
Interest in Delivery at Home for Future Deliveries by Selected Characteristics
Women with a Live Birth after December 1987
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Interest in Home Delivery				Total	No. of Women
	Yes	Possibly	No	Not sure		
Total	4.5	3.2	87.8	4.5	100.0	(1382)
Region						
Bohemia	4.4	3.4	88.1	4.1	100.0	(855)
Moravia	4.5	2.9	87.5	5.1	100.0	(527)
Size of Place						
< 5000	3.6	2.3	90.9	3.2	100.0	(460)
5000-19999	4.5	3.5	87.8	4.2	100.0	(300)
20000 +	5.1	3.7	85.6	5.6	100.0	(622)
Age						
15-19	8.2	3.3	80.3	8.2	100.0	(51)
20-24	4.5	4.5	85.5	5.4	100.0	(418)
25-29	5.2	2.8	87.9	4.1	100.0	(533)
30-34	2.0	2.8	92.5	2.8	100.0	(249)
35-44	4.2	1.4	90.1	4.2	100.0	(131)
Education						
Primary	5.8	2.3	90.7	1.2	100.0	(162)
Secondary No Diploma	5.9	2.7	86.8	4.5	100.0	(531)
Secondary Diploma	3.1	4.0	87.9	5.0	100.0	(563)
Any University	2.4	3.1	88.2	6.3	100.0	(126)
Parity						
Primiparae	4.4	3.7	87.7	4.2	100.0	(561)
Multiparae	4.5	2.9	88.0	4.6	100.0	(821)

TABLE V.27
Most Important Problem during the First Week after Delivery* by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Principal Problem during Week after Delivery										No. of Women
	No Problem	Own Health	Health/ Care of the Child	Breast-feeding	Hospital Environ.	Visiting/ Contact with Family	Prof. Staff	Other	Do Not Rem.	Total	
Total	40.0	17.8	17.0	11.0	5.0	4.3	2.5	1.0	1.3	100.0	(1382)
Region											
Bohemia	40.2	17.6	16.8	12.4	5.1	3.5	2.7	0.1	1.5	100.0	(855)
Moravia	39.7	18.1	17.4	8.9	4.7	5.4	2.2	2.5	1.1	100.0	(527)
Size of Place											
< 5000	49.2	16.5	13.9	9.3	5.1	3.4	1.3	0.6	0.8	100.0	(460)
5000-19999	40.1	19.2	17.6	11.9	2.9	4.8	2.2	0.3	1.0	100.0	(300)
20000 +	33.3	18.1	19.0	11.9	5.9	4.6	3.6	1.7	1.9	100.0	(622)
Age											
15-19	36.1	13.1	31.1	11.5	6.6	1.6	0.0	0.0	0.0	100.0	(51)
20-24	37.6	20.4	15.2	12.5	4.5	3.9	2.9	1.2	1.8	100.0	(418)
25-29	41.3	15.9	16.6	11.8	4.5	5.0	3.2	0.8	0.9	100.0	(533)
30-34	38.3	19.0	13.4	10.7	7.5	5.5	1.6	2.0	2.0	100.0	(249)
35-44	47.2	16.9	24.7	4.2	2.8	1.4	1.4	0.7	0.7	100.0	(131)

TABLE V.27
Most Important Problem during the First Week after Delivery* by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Principal Problem during Week after Delivery												
Characteristics	No Problem	Own Health	Health/ Care of the Child	Breast-feeding	Hospital Environ.	Visiting/ Contact		Prof. Staff	Other	Do Not Rem.	Total	No. of Women
						Family						
Marital Status												
Currently Married	40.8	17.6	16.6	11.1	5.0	4.2		2.4	1.1	1.4	100.0	(1274)
Previously Married	26.5	26.5	16.2	8.8	7.4	8.8		4.4	0.0	1.5	100.0	(64)
Single	38.2	12.7	29.1	12.7	1.8	0.0		3.6	1.8	0.0	100.0	(44)
Education												
Primary	40.7	20.3	15.7	8.1	5.2	2.9		2.9	0.6	3.5	100.0	(162)
Secondary No Diploma	45.2	17.3	16.0	10.8	2.5	4.0		1.4	1.1	1.6	100.0	(531)
Secondary Diploma	38.1	17.0	19.0	10.2	6.4	4.7		2.9	1.0	0.7	100.0	(563)
Any University	25.2	20.5	14.2	19.7	8.7	5.5		4.7	1.6	0.0	100.0	(126)
Parity												
Primiparae	32.1	19.6	20.1	14.4	4.7	4.1		2.9	1.0	1.2	100.0	(561)
Multiparae	45.6	16.5	14.9	8.7	5.1	4.4		2.3	1.1	1.4	100.0	(821)

* Limited to the most recent live birth after December 1987.

TABLE V.28
Most Important Problem after Returning Home Following Delivery* by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Principal Problem after Returning Home								Do Not Rem.	Total	No. of Women
	No Problem	Own Health	Health of the Child	Care of the Child	Home Environ.	Family	Economic	Other			
Total	58.1	10.0	8.4	7.1	4.5	3.4	2.2	4.6	1.7	100.0	(1382)
Region											
Bohemia	56.8	10.3	8.4	8.0	4.2	3.6	2.6	4.2	1.8	100.0	(855)
Moravia	60.1	9.4	8.5	5.8	5.1	2.9	1.4	5.3	1.4	100.0	(527)
Size of Place											
< 5000	66.5	6.5	7.6	6.5	3.4	3.2	1.3	3.6	1.5	100.0	(460)
5000-19999	55.1	9.6	8.7	6.4	6.4	3.2	3.8	4.4	2.2	100.0	(300)
20000 +	53.4	12.7	9.0	7.9	4.5	3.6	2.0	5.5	1.5	100.0	(622)
Age											
15-19	50.8	6.6	14.8	9.8	1.6	4.9	3.3	8.2	0.0	100.0	(51)
20-24	49.2	11.3	9.1	10.4	5.9	4.8	3.4	3.6	2.3	100.0	(418)
25-29	60.6	8.6	8.2	7.7	5.2	2.4	1.5	4.0	1.9	100.0	(533)
30-34	66.0	10.3	6.7	2.4	3.2	2.0	1.6	6.8	1.2	100.0	(249)
35-44	65.6	12.0	7.8	2.1	1.4	4.2	1.4	4.9	0.7	100.0	(131)

TABLE V.28
Most Important Problem after Returning Home Following Delivery* by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Principal Problem after Returning Home									Total	No. of Women
	No Problem	Own Health	Health of the Child	Care of the Child	Home Environ.	Family	Economic	Other	Do Not Rem.		
Marital Status											
Currently Married	59.4	10.4	8.4	7.1	4.7	2.1	2.0	4.4	1.6	100.0	(1274)
Previously Married	50.0	7.4	4.4	5.9	4.4	20.6	2.9	0.0	4.4	100.0	(64)
Single	38.2	3.6	14.5	9.1	1.8	10.9	5.5	16.4	0.0	100.0	(44)
Education											
Primary	58.7	7.9	8.7	4.1	5.8	2.9	4.7	4.7	3.5	100.0	(162)
Secondary No Diploma	58.2	12.3	8.6	6.1	4.3	3.2	1.4	4.5	1.3	100.0	(531)
Secondary Diploma	58.8	8.1	8.1	7.6	4.7	3.8	2.2	4.7	1.9	100.0	(563)
Any University	53.5	12.6	8.7	13.4	3.1	2.4	1.6	4.8	0.0	100.0	(126)
Parity											
Primiparae	46.1	10.5	9.6	15.0	5.1	4.4	3.0	4.3	2.0	100.0	(561)
Multiparae	66.5	9.6	7.6	1.5	4.2	2.6	1.5	4.9	1.4	100.0	(821)

* Limited to the most recent live birth after December 1987.

TABLE V.29
Amount of Prenatal Counseling during Most Recent Pregnancy* by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Characteristics	Number of Prenatal Counseling Sessions				Total	No. of Women
	None	1-3	4 +	Don't Know		
Total	76.0	13.7	7.8	2.5	100.0	(1382)
Region						
Prague	78.0	12.1	8.3	1.5	100.0	(128)
Central Bohemia	79.2	10.1	8.3	2.4	100.0	(162)
South Bohemia	70.9	11.7	14.6	2.9	100.0	(101)
West Bohemia	87.9	5.3	4.5	2.3	100.0	(125)
North Bohemia	84.2	6.5	5.4	3.8	100.0	(182)
East Bohemia	76.4	16.1	5.6	1.9	100.0	(157)
South Moravia	66.5	21.7	7.7	4.0	100.0	(258)
North Moravia	73.2	16.8	8.9	1.1	100.0	(269)
Size of Place						
< 5000	74.9	16.0	6.1	3.0	100.0	(460)
5000-19999	77.2	12.8	7.1	2.9	100.0	(300)
20000 +	76.3	12.4	9.3	2.0	100.0	(622)
Age						
15-19	73.8	14.8	6.6	4.9	100.0	(51)
20-24	76.9	12.5	7.9	2.7	100.0	(418)
25-29	74.8	15.7	6.7	2.8	100.0	(533)
30-34	77.1	13.0	8.3	1.6	100.0	(249)
35-44	76.1	12.4	9.7	1.8	100.0	(105)
Education						
Primary	84.9	7.0	4.7	3.5	100.0	(162)
Secondary No Diploma	77.1	13.0	6.7	3.2	100.0	(531)
Secondary Diploma	72.5	16.4	9.3	1.7	100.0	(563)
Any University	75.6	13.4	9.4	1.6	100.0	(126)
Parity						
Primiparae	72.1	15.4	10.3	2.2	100.0	(561)
Multiparae	78.8	12.5	6.0	2.7	100.0	(821)

*Limited to the most recent pregnancy resulting in a live birth after December 1987.

TABLE V.30
Percent of Respondents' Children Who Were Breastfed
by Selected Characteristics
1993 Czech Republic Reproductive Health Survey

Characteristics	Percent Breastfed	Number of Children
Total	90.6	(1,692)
Region		
Bohemia	89.5	(1,031)
Moravia	92.3	(661)
Size of Place		
Less Than 5,000	90.5	(584)
5,000(19,999	91.5	(364)
20,000 +	90.3	(744)
Age		
15-19	84.1	(53)
20-24	90.8	(539)
25-29	94.3	(665)
30-34	87.4	(290)
35-39	85.8	(117)
40-44	71.0	(28)
Education		
Primary Only	88.9	(207)
Secondary, No Diploma	87.9	(656)
Secondary, Diploma	93.5	(672)
Any University	92.4	(157)
Religion		
None	89.7	(1,126)
Catholic	92.9	(501)
Other	89.1	(61)

*Includes all live births to respondents after December 1987

TABLE V.31
Mean Duration of Breastfeeding* for Children
Born since January 1988, by Selected Characteristics
1993 Czech Republic Reproductive Health Survey

Characteristics	Mean Duration (Months)	
	All Children	Children Ever Breastfed
Total	3.7	4.2
Region		
Bohemia	3.6	3.9
Moravia	4.0	4.5
Size of Place		
Less Than 5,000	3.9	4.3
5,000-19,999	4.4	4.8
20,000 +	3.3	3.7
Age		
15-24	3.4	3.9
25-44	4.1	4.6
Education		
No Secondary Diploma	2.7	3.1
Secondary Diploma/University	4.9	5.3

*Calculated by means of the prevalence/incidence method, using current breastfeeding status data.

VI. YOUNG ADULT SEXUAL BEHAVIOR

One section of the CRRHS questionnaire was devoted to examining issues of reproduction and sexuality among young adults, those between the ages of 15 and 24 years. The purpose of the questions asked of these young women was to describe the age at which women first have sexual intercourse, use of contraception, circumstances surrounding any premarital pregnancy, and related information. Despite the sensitive nature of the information requested, cooperation from respondents was extremely high: Only 2% of young women refused to answer questions regarding sexual activity.

Initiation of Sexual Intercourse

Using life table procedures, we estimated the median age at which females first had sexual intercourse and the proportion who had ever had intercourse before each age from 15 to 23 ([Table VI.1](#) and [Figure VI.1](#)). The estimated median age at first intercourse was 17.5 years. It appears that the age at which females begin having sexual relations may have been decreasing, since the median age was 0.5 years less for those 15-19 years of age than for those five years older. It can also be seen that the percentage reporting sexual experience by each age from 15 to 18 was higher for the younger (15-19 year) cohort than for the older (20-24 year) cohort. Sexual intercourse before age 15 was still quite rare for girls, with only 3 percent reported sexually experienced before their fifteenth birthday, although it may have been becoming more common. By about age 21, only 3% of women had not yet had sexual intercourse.


[Table VI.2](#) examines differences in the age at first sexual intercourse according to geographic variables, religion, and household income. (Education is not included in this analysis because so many young women were still attending school and for those respondents current educational attainment would be misleadingly low.) The overall percentages reporting sexual experience were 36% for 15-17 year-olds, 82% for 18-19 year-olds, and 98% for 20-24 year-olds. There were some noteworthy differences between subgroups for ages 15-17 and 18-19, but for 20-24 year-olds there was virtually no variation according to the characteristics examined. Bohemian women tended to become sexually active before Moravian women (39% experienced, as opposed to 31% among 15-17 year-olds). Likewise, women espousing no religion started sexual relations earlier than Catholics (37% compared with 32%). There was no clear relationship between household income and sexual experience.

Clearly premarital sexual intercourse has become the norm throughout the Czech Republic. Over 99% of sexually experienced women (all but six respondents) first had intercourse before they were married ([Table VI.3](#), upper panel). Eighty-four percent of these women said their first sexual relations were with their "boyfriend". Another 10% described their first contact as a "friend", while only 5% first had sexual relations with

Figure VI.1

Percent of Women Who Ever Had Sexual Intercourse before Given Age, by Age at Interview

15-24 Year-Old Respondents, 1993 CRRHS



their fiancé. As age at first intercourse increases, the likelihood that the first experience was with a "friend" decreases, and with a fiancé increases.

Of sexually experienced young adults, about two-thirds first had sexual intercourse less than four months or four to six months after their relationship began ([Table VI.3](#), lower panel). Relatively small proportions had intercourse the first time they met their first partner (5%) or more than 12 months after the relationship started (9%). As a rule, the younger the age at which the woman began sexual activity, the shorter was the length of the relationship when that activity started. Among those first having intercourse before age 16, 13% had sex the first time they met their first partner. This figure was only 2% for those who became sexually active after age 17.

Contraception at First Intercourse


Forty-two percent of young adult respondents or their partner reportedly used no form of contraception the first time they had sexual intercourse ([Table VI.4](#)). Those who first had sex before age 16 and those without a secondary school diploma were the least likely to use contraception ([Figure VI.2](#)). Residence and religion were not related to whether contraception was used. Among those who did use contraception, over 90% used either withdrawal or condoms, which is not surprising, given that these methods are readily accessible and do not require contact with a clinic or physician. Withdrawal and condoms were the predominant method used by young women, regardless of their characteristics.

When asked why they did not use contraception at first intercourse, 44% of non-users said the reason was that they did not expect to have sex ([Table VI.5](#)). Another 27% said they did not think it was possible for them to become pregnant. The percentage responding that they did not expect to have sex decreased with increasing age at first sex, from 57% for those under 16 years to 35% for those over 17 years. The proportion saying they did not contracept because they did not have any contraceptive method with them increased with age, from 2% to 13%.

Premarital Pregnancy

It is important to keep in mind that underreporting of induced abortions (and, therefore, pregnancies terminated by induced abortion) is likely to have had a significant effect on the results presented on premarital pregnancies, especially if premarital abortions were even more likely than abortions within marriage to go unreported. We expect that the proportion of females experiencing premarital pregnancy was somewhat higher than reported and that the proportion of those women who went on to marry the man by whom they became pregnant may also have been understated. About one-half of 15 to 24 year-old respondents reportedly had ever been pregnant at the time of interview and

Figure VI.2
 Contraceptive Method Used at First Sexual Intercourse,
 by Age at First Sex and by Education,
 15-24 Year-Old Respondents, 1993 CRRHS



about half of those said they were not married at the time they first became pregnant.

The top panel of [Table VI.6](#) reveals that in about three-fourths of instances of premarital pregnancy the man by whom she became pregnant was her "boyfriend", with most of the remainder being a respondent's fiancé. Among those respondents who were 18 or older the man was more likely to be a fiancé than among those under 18. Seventy-three percent of unmarried women who became pregnant said they went on to marry the man by whom they became pregnant. The bottom panel of [Table VI.6](#), shows that in about seven of every ten cases, respondents reported that their partner wanted to get married after learning of the pregnancy. Fourteen percent of men preferred that the woman get an abortion. Eight percent wanted the respondent to have the baby, but did not want to get married. Finally, 2% of women said that the man never learned of the pregnancy.

When asked about the reaction of their parents after finding out about their pregnancy, 43% of women said they were encouraged to get married ([Table VI.7](#)). This proportion was highest among those women who were under 18 when they became pregnant. Thirty percent of parents did not interfere, i.e., they let the women/couple make the decision on their own. As might be expected this percentage was lowest for those women who were less than 18 years of age, 16%. Eleven percent thought the woman should have the baby without getting married, 9% encouraged abortion, and 6% did not know about the pregnancy. The proportion encouraging abortion was much higher for those becoming pregnant before age 18.

Family Planning Discussions with Parents

Finally, all 15 to 24 year-old respondents were asked whether their parents had ever discussed the use of pregnancy prevention methods with them. Sixty percent of women said they had never had such discussions ([Table VI.8](#)). Among the minority whose parents did talk to them about contraception, discussions first took place at a fairly late age: only 14% of respondents' parents discussed the issue with their daughter before she was 15 years old. There were no strong relationships apparent between whether such talks took place and respondent characteristics. Even where such relationships might have been expected, for example, between religious Catholics and others, and according to whether the respondent used a method when she first had sexual intercourse, no strong correlations appear. Women who used contraception the first time they had intercourse were slightly more likely than non-users to have had discussions about family planning with their parents. Those women who had never had sex prior to the time of interview tended to have discussions about family planning with their parents earlier than sexually experienced women.

TABLE VI.1
Life Table Estimates of Percent of Women Who Have Ever Had Sexual Intercourse
before Given Ages and Median Age at First Intercourse, according to Age at Interview
15 to 24 Year-Old Females
1993 Czech Republic Reproductive Health Survey

Age	Total	Age at Interview	
		15-19	20-24
15	2.8	3.9	1.5
16	12.9	16.6	8.4
17	35.7	44.1	27.1
18	64.4	68.7	59.9
19	82.3	82.1	81.0
20	91.9	---	91.7
21	97.0	---	96.9
22	97.7	---	97.6
23	98.6	---	98.6
Median Age at First Intercourse	17.5	17.2	17.7

TABLE VI.2
Percent of 15 to 24 Year-Old Females Who Have Ever Had
Sexual Intercourse, by Selected Characteristics and Age at Interview
1993 Czech Republic Reproductive Health Survey

Characteristics	Age at Interview					
	15-17		18-19		20-24	
	%	<i>N</i>	%	<i>N</i>	%	<i>N</i>
Total	35.5	(359)	81.6	(287)	97.6	(737)
Region						
Bohemia	39.1	(197)	85.1	(179)	97.6	(457)
Moravia	30.8	(162)	76.4	(108)	97.5	(280)
Size of Place						
Less than 5000	31.5	(111)	77.4	(95)	98.9	(237)
5000-20000	44.6	(73)	79.8	(56)	96.9	(149)
20000 +	34.0	(175)	85.9	(136)	97.0	(351)
Religion						
None	37.0	(253)	86.0	(205)	97.5	(501)
Catholic	31.5	(89)	70.9	(68)	98.1	(188)
Monthly Household Income*						
0-7000	31.9	(128)	83.6	(121)	97.9	(412)
7001-10000	46.3	(80)	79.3	(73)	97.8	(162)
10001 and more	38.9	(52)	85.2	(55)	98.5	(109)
Don't Know	28.1	(97)	73.1	(35)	91.9	(45)

*Fourteen women refused to report their monthly household income.

TABLE VI.3
Relationship to First Sexual Partner and
Length of Relationship at the Time of First Sexual Intercourse
according to Age at First Sexual Intercourse
Sexually Experienced Females, 15 to 24 Years Old
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Relationship to First Partner/ Length of Relationship	Total*	Age at First Intercourse		
		Before 16	16-17	18-24
Relationship				
Husband	0.4	0.0	0.3	1.1
Fiance	5.0	0.5	4.1	9.9
Boyfriend	83.7	74.3	86.0	84.7
Friend	10.0	20.1	9.5	4.3
Rape	0.9	5.1	0.1	0.0
Total	100.0	100.0	100.0	100.0
Number of Respondents**	(1,080)	(142)	(614)	(302)
Length of Relationship				
First Time They Met	5.1	12.9	4.6	1.5
Less Than 4 Months	33.7	36.6	33.8	32.4
4-6 Months	31.4	28.2	31.1	35.0
7-12 Months	17.2	12.4	18.7	17.3
More Than 12 Months	9.1	6.4	8.7	12.1
Do Not Remember	3.4	3.5	3.2	1.7
Total	100.0	100.0	100.0	100.0
Number of Respondents***	(1,060)	(134)	(609)	(296)

*Includes 22 women who said they did not remember their age at first intercourse.

**Does not include 8 women who did not report their relationship to their first partner.

***Does not include 28 women who did not report the length of the relationship.

TABLE VI.4
Contraceptive Method Used at First Sexual Intercourse
according to Selected Characteristics
Sexually Experienced Females, 15 to 24 Years Old
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Contraceptive Method Used					Total	Number of Women
	None	Withdrwl	Condoms	Other	DK/DNR		
Total	41.6	29.1	23.0	4.5	1.8	100.0	(1,080)
Age, First Sex							
Before 16	47.1	24.5	23.0	4.4	1.0	100.0	(136)
16-17	40.2	30.4	23.2	4.2	2.0	100.0	(614)
18-24	39.9	30.3	23.0	6.0	0.9	100.0	(303)
Don't Know	57.6	12.1	18.2	0.0	12.1	100.0	(27)
Current Age							
15-19	37.8	28.6	27.1	6.0	0.5	100.0	(361)
20-24	44.2	29.4	20.2	3.6	2.6	100.0	(719)
Region							
Bohemia	40.9	28.9	24.1	4.3	1.8	100.0	(676)
Moravia	42.7	29.3	21.2	5.0	1.7	100.0	(404)
Education							
Primary	52.2	19.1	24.3	2.9	1.5	100.0	(98)
Sec, No Dipl	47.9	26.8	19.4	4.1	1.8	100.0	(524)
Sec, Diploma	31.7	35.4	25.0	6.1	1.9	100.0	(393)
University	30.4	26.6	39.2	2.5	1.3	100.0	(65)
Religion							
None	41.9	29.6	22.1	4.6	1.8	100.0	(763)
Catholic	41.4	27.6	23.9	4.9	2.1	100.0	(257)
Other	41.5	29.2	26.2	3.1	0.0	100.0	(54)

TABLE VI.5
Reason for not Using a Contraceptive Method at the Time of First Sexual Intercourse
according to Age at First Sexual Intercourse
Sexually Experienced Females, 15 to 24 Years Old
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Reason for not Using Contraception	Total*	Age at First Intercourse		
		Before 16	16-17	18-24
Did Not Expect to Have Sex	44.0	57.1	46.1	35.4
Did Not Think Pregnancy Possible	26.9	23.5	27.6	27.8
Did Not Have Contraception	8.5	2.0	8.7	12.5
Wanted to Become Pregnant	3.2	4.1	0.9	7.6
Did Not Know Contraceptive Methods	2.4	4.1	2.8	0.7
Other Reasons	3.1	3.1	2.8	4.2
Don't Know	11.9	6.1	11.1	11.8
Total	100.0	100.0	100.0	100.0
<i>Number of Respondents</i>	<i>(479)</i>	<i>(68)</i>	<i>(264)</i>	<i>(128)</i>

*Includes 19 women who said they did not remember their age at first intercourse.

TABLE VI.6
Relationship to Partner at Time of First Pregnancy, Whether She Married That Man, and
Attitude of Partner Regarding the Pregnancy according to Age at First Pregnancy,
Females 15 to 24 Years Old with Any Reported Premarital Pregnancies
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

		Age at First Pregnancy		
	Total	Under 18	18-19	20-24
Relationship to Man				
Boyfriend	75.4	81.5	72.5	72.9
Fiance	22.9	14.1	26.7	27.1
Other	1.7	4.3	0.8	0.0
Whether Married Him				
Yes	73.4	70.0	77.5	70.0
No	26.6	30.0	22.5	30.0
Partner's Attitude				
Wanted to Marry	71.4	68.8	75.6	67.1
Wanted Abortion	13.9	12.9	10.7	21.4
Accepted without Marriage	8.2	9.7	6.1	10.0
Didn't Know about Pregnancy	1.7	3.2	1.5	0.0
Other/Don't Know	4.8	5.4	6.1	1.4
Total	100.0	100.0	100.0	100.0
<i>Number of Respondents</i>	<i>(263)</i>	<i>(84)</i>	<i>(118)</i>	<i>(61)</i>

TABLE VI.7
Attitude of Respondent's Parents Regarding Her First Pregnancy
according to Age at First Pregnancy,
Females 15 to 24 Years Old with Any Reported Premarital Pregnancies
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Parents' Attitude	Total	Age at First Pregnancy		
		Under 18	18-19	20-24
Wanted Her to Marry	42.5	45.2	42.7	38.6
Did Not Interfere	29.6	16.1	35.9	35.7
Accepted without Marriage	10.9	16.1	8.4	8.6
Wanted Her to Have Abortion	8.8	17.2	5.3	4.3
Didn't Know about Pregnancy	5.8	4.3	3.1	12.9
Don't Know	2.4	1.1	4.6	0.0
Total	100.0	100.0	100.0	100.0
<i>Number of Respondents</i>	<i>(263)</i>	<i>(84)</i>	<i>(118)</i>	<i>(61)</i>

TABLE VI.8
Age at Which Respondent's Parents First Discussed
Family Planning Methods with Her, according to Selected Characteristics
Females 15 to 24 Years Old
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Age at Which First Talked about Family Planning						Total	N
	10-12	13-14	15-16	17+	DK/DNR	Never		
Total	2.1	11.4	18.1	6.5	1.3	60.6	100.0	(1,418)
Current Age								
15-19	3.2	15.2	18.2	3.3	0.6	59.5	100.0	(662)
20-24	0.8	6.5	18.0	10.5	2.2	61.9	100.0	(756)
Region								
Bohemia	2.3	11.4	20.9	7.4	1.2	56.8	100.0	(856)
Moravia	1.9	11.3	14.0	5.1	1.4	66.2	100.0	(562)
Education								
Primary	3.1	13.7	9.3	0.3	0.0	73.6	100.0	(185)
Sec, No Dipl	2.5	14.2	17.6	6.0	1.0	58.7	100.0	(742)
Sec, Diploma	1.0	5.1	24.5	10.8	1.4	57.3	100.0	(421)
University	1.2	6.0	16.7	8.3	9.5	58.3	100.0	(70)
Religion								
None	2.0	11.9	19.3	6.8	0.9	59.0	100.0	(978)
Catholic Attend	4.3	11.2	17.2	5.2	3.4	58.6	100.0	(80)
Catholic Not Att	0.8	10.6	15.6	5.6	1.7	65.8	100.0	(279)
Other	5.2	7.3	12.5	6.3	3.1	65.6	100.0	(75)
Age, First Sex								
Before 16	1.9	18.1	11.6	1.9	0.5	66.0	100.0	(136)
16-17	0.7	6.8	23.7	7.4	1.3	60.1	100.0	(614)
18-24	0.6	5.9	17.8	15.3	2.3	58.1	100.0	(303)
Don't Remember	0.0	3.0	6.1	6.1	6.1	78.8	100.0	(27)
No Sex Yet	5.0	20.2	14.2	1.4	0.6	58.5	100.0	(297)
Use at First Sex								
Used	1.0	10.0	19.9	9.0	1.7	58.3	100.0	(601)
Did Not Use	0.5	6.2	20.0	7.6	1.4	64.2	100.0	(459)
Never Had Sex	5.1	20.5	14.4	1.4	0.0	58.5	100.0	(297)

VII. REPRODUCTIVE HEALTH KNOWLEDGE AND ATTITUDES

Each respondent answered a series of questions regarding her knowledge and attitudes about several aspects of reproduction. These questions included opinions about the conditions under which abortion should be allowable, knowledge of the fertile period, opinions about the efficacy, safety, and side effects of oral contraceptives, and views about women's role. The results of these questions should prove useful for policymakers and for developing reproductive health education and promotion activities.

Attitudes about Restrictions on Abortion

As described in Chapter III, the rate of induced abortions in the Czech Republic has been high by world standards, with official statistics for 1992 showing that there were 48 abortions per 1000 15 to 44 year-old women and 78 abortions for every 100 live births. Abortion laws are permissive, although minor restrictions have been added in recent years and the cost of having an abortion performed has increased substantially.

Eighty-five percent of reproductive age women thought that women/couples should have the right to decide whether to have a pregnancy terminated by induced abortion for any reason they choose ([Table VII.1](#)). Every segment of the population examined was opposed, by a wide margin, to legal restrictions on a woman's access to abortion. There were only small differences in opinions according to place of residence, size of place, age, and marital status. As education and household income increased, so did the proportion of women in favor of the right to choose abortion. Even among Catholics who attend mass regularly, seven of every ten women felt there should be no limitations on legal access to abortion ([Figure VII.1](#)).


The denominator for [Table VII.2](#) consists of only the 14 percent (657 women) of respondents who reported that there should be some restrictions on access to abortion. An overwhelming majority of these women felt that abortion was justified under certain circumstances. Only 4% felt that a pregnancy endangering a woman's life was not adequate grounds for abortion. The proportions rose to 11 % if there was risk of a fetal defect, 56% if the couple could not afford another child, and 70% if the woman was not married.

[Table VII.3](#) shows the overall percentages of respondents who felt that abortion was not justified and should not be available under each of the six circumstances listed in the previous table. These figures range from less than 1% when a pregnancy endangers a woman's life to 10% when a woman is not married. For the first four circumstances there were few differences between population segments. For the others restrictiveness decreased as education increased. Even most Catholics who attended religious services regularly, the group with the most restrictive views about abortion availability, felt that abortion was justified under each of the six circumstances asked about.

Figure VII.1

Percent of Respondents Who Think Women Should be Able to Decide Whether to Have an Abortion for Any Pregnancy, by Education and by Religion,

1993 CRRHS



Knowledge of the Menstrual Cycle

Each respondent was asked at what time during the menstrual cycle is a woman at the greatest risk of becoming pregnant. Fifty-five percent of respondents correctly said that the chance of pregnancy is greatest about halfway between menstrual periods ([Table VII.4](#)). This proportion is considerably higher than has been found in reproductive health surveys in developing countries. Knowledge was lowest among 15-19 year-olds (39%). It also increased sharply with increasing education, from 32% to 83%. Fortunately, those women using natural family planning (NFP) methods, the success of which depend on a woman/couple knowing when conception is the most likely to occur, have the best knowledge (75%), however, one of every four NFP users were unaware of the time that conception is most probable.

Opinions about Oral Contraceptives


Respondents were asked a series of questions in order to learn their opinions about oral contraceptives. Recall that in Chapter III it was noted that among women who were using generally less reliable methods of contraception a fear of negative health effects, especially from oral contraceptives, was overwhelmingly the most important factor in the decision to avoid use of modern methods of family planning. It has also been reported anecdotally that many Czech gynecologists still feel that hormonal contraception should be avoided or not used for extended periods because of potential health effects. A recent study found that most Czech gynecologists felt that low dose oral contraceptives were safe for nonsmokers under 35 years of age, but there was a significant minority, especially among older physicians, who thought they were unsafe (Visser, Uzel et al. 1993). The opinions of physicians is almost certain to have an effect on the views of women. The information presented here documents attitudes of all women of reproductive age regarding the reliability, safety, and perceived side effects of oral contraceptives.

On the whole, many women underestimated the ability of oral contraceptives to prevent pregnancy. Only about half of respondents said that a women who took them correctly could be completely or almost sure that she would not become pregnant ([Table VII.5](#) and [Figure VII.2](#)). One of every five women indicated that the reliability of oral contraceptives was quite low. There was relatively little variation across the population in opinions about reliability. Confidence in oral contraceptives increased with education. It was, not surprisingly, highest among oral contraceptive users, but otherwise was unrelated to whether a woman was contracepting.

Overall, about one-fifth of women felt that oral contraceptives posed a substantial risk to women's health and a slightly smaller percentage said they did not know if they posed a risk ([Table VII.6](#) and [Figure VII.3](#)). The percentage who felt that they were unsafe increased slightly with age and was highest among the least educated. One-third of Catholics attending services felt that oral contraceptives were unsafe, the highest

Figure VII.2
Opinions about the Reliability of Oral Contraceptives

Percent Distribution, 1993 CRRHS



proportion of any population segment examined.

Fifty-three percent of women were aware that oral contraceptives make menstrual periods more regular, while 42% felt that oral contraceptives cause weight gain ([Table VII.7](#)). Sizable minorities of respondents agreed with the statements that oral contraceptives: increase the risk of getting cancer (25%); are bad for blood circulation (21%); can cause infertility (16%); and cause nervousness (13%). For each of these four statements about one-half of respondents said they did not know whether they were true. Reliance on methods of low reliability (as well as unintended pregnancy and induced abortion rates) would be likely to decrease if women became more aware of recent evidence regarding the safety of oral contraceptives, especially of newer formulations.

Women's Family and Reproductive Roles

Respondents were asked whether they agreed or disagreed with five statements, shown in [Table VII.8](#), regarding women's roles and reproduction. There was considerable agreement about the first three of these statements. Sizable majorities disagreed with the statements that "Child care is women's work" and "A woman should be a virgin when she marries", while agreeing that it is all right for a woman not to have children if she so desires. These results indicate that most Czech women do not feel that women should be restricted to traditional roles. Ninety-two and ninety-seven percent, respectively, correctly agreed that a woman could become pregnant the first time she has intercourse and that a condom should not be used more than once.

Figure VII.3
Opinions about the Safety of Oral Contraceptives
Percent Distribution, 1993 CRRHS




TABLE VII.1
Percent of Respondents Who Think That Women Should
Be Able to Decide Whether to Have an Abortion for Any Pregnancy,
by Selected Characteristics
1993 Czech Republic Reproductive Health Survey

Characteristics	Percent	No. of Women
Total	85.2	(4,497)
Residence		
Bohemia	86.4	(2,737)
Moravia	83.4	(1,760)
Size of Place		
Less Than 5,000	84.1	(1,472)
5,000-19,999	87.1	(967)
20,000 +	85.2	(2,058)
Current Age		
15-19	82.5	(662)
20-24	84.3	(756)
25-29	86.9	(828)
30-34	88.0	(720)
35-39	84.8	(805)
40-44	85.9	(726)
Union Status		
Currently Married	86.0	(3,217)
Previously Married	87.9	(355)
Never Married	82.5	(925)
Education		
Primary only	78.8	(738)
Secondary, No Diploma	84.0	(1,779)
Secondary, Diploma	88.9	(1,601)
Any University	91.0	(379)
Religion		
None	87.2	(2,913)
Catholic, Attends Services	70.8	(256)
Catholic, Does Not Attend	85.7	(1,134)
Other	71.7	(180)
Monthly Household Income (Crowns)		
0-3000	77.4	(188)
3,001-7,000	83.6	(1,885)
7,001-10,000	87.8	(1,278)
10,000 +	87.8	(803)
Not Stated	88.3	(253)
Do Not Know	80.7	(90)

TABLE VII.2
Opinions on Whether Women Should Be Able to Have an Abortion
under Given Circumstances, among Respondents Who Think
There Should Be Restrictions on Whether Women Can Have Abortions*
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Circumstance	Whether Women Should Be Allowed to Have an Abortion under the Given Circumstance:			Total
	Yes	Depends	No	
Woman's life endangered	90.7	2.5	4.3	100.0
Fetal defect	73.9	9.8	10.9	100.0
Woman's health endangered	71.5	10.8	12.4	100.0
Pregnancy resulted from rape	71.0	7.8	14.9	100.0
Woman/Couple cannot afford a baby	15.6	18.1	55.5	100.0
Woman is not married	8.3	18.2	70.3	100.0

Number of respondents = 657

*This table does not include the 85.2% of respondents who said that women should be able to decide whether to get an abortion under any circumstances.

TABLE VII.3
Percent of Respondents Who Think That Women Should Not Be Able
to Decide Whether to Have an Abortion under Selected Circumstances
by Selected Characteristics,
1993 Czech Republic Reproductive Health Survey

Characteristics	Circumstance (see key below)						No. of Women
	1	2	3	4	5	6	
Total	0.6	1.6	1.8	2.2	8.2	10.4	(4,497)
Residence							
Bohemia	0.5	1.0	1.4	2.0	7.0	9.4	(2,737)
Moravia	0.8	2.6	2.5	2.5	10.0	11.8	(1,760)
Current Age							
15-19	0.6	2.5	2.8	3.4	9.2	13.0	(662)
20-24	0.5	1.8	1.7	1.7	8.5	10.8	(756)
25-29	0.7	1.4	1.7	2.2	7.9	9.3	(828)
30-34	0.5	1.1	1.3	2.0	7.7	8.3	(720)
35-39	0.6	1.4	1.3	2.1	8.5	10.9	(805)
40-44	0.8	1.3	2.0	1.7	7.1	9.2	(726)
Education							
Primary only	0.8	1.1	2.8	2.9	11.0	15.6	(738)
Secondary, No Diploma	0.7	1.9	1.8	2.2	8.6	10.7	(1,779)
Secondary, Diploma	0.3	1.4	1.3	1.6	6.7	8.1	(1,601)
Any University	1.2	2.2	2.2	2.9	6.1	6.6	(379)
Religion							
None	0.3	0.7	0.9	1.2	6.0	8.3	(2,913)
Catholic, Attends Services	5.0	10.9	11.8	11.8	23.0	25.5	(256)
Catholic, Does Not Attend	0.1	0.7	1.0	1.0	8.1	10.1	(1,134)
Other	1.8	6.8	7.8	10.5	21.9	23.3	(180)

Key to circumstances:

- 1) Woman's life is in danger from the pregnancy
- 2) The fetus has a physical deformity
- 3) The woman's health may be harmed by the pregnancy
- 4) The pregnancy resulted from rape
- 5) The couple cannot afford to have a(nother) child
- 6) The woman is not married

TABLE VII.4
Percent of Respondents with Correct Knowledge of When a Woman Is the Most Likely to Conceive
during Her Menstrual Cycle, according to Selected Characteristics
1993 Czech Republic Reproductive Health Survey

Characteristics	Percent with Correct Knowledge	Number of Respondents
Total	55.4	(4,497)
Residence		
Bohemia	54.9	(2,737)
Moravia	56.0	(1,760)
Size of Place		
Less than 5,000	52.5	(1,472)
5,000-19,999	54.7	(967)
20,000 +	57.8	(2,058)
Current Age		
15-19	39.4	(662)
20-24	58.6	(756)
25-29	63.2	(828)
30-34	61.0	(720)
35-39	60.4	(805)
40-44	54.0	(726)
Education		
Primary only	31.6	(738)
Secondary, No Diploma	46.5	(1,779)
Secondary, Diploma	72.5	(1,601)
Any University	82.6	(379)
Religion		
None	56.8	(2,913)
Catholic, Attends Services	48.4	(256)
Catholic, Does Not Attend	53.6	(1,134)
Other	52.5	(180)
Current contraception		
None	46.2	(1,736)
Oral Contraceptives	65.5	(371)
Other Modern Methods	59.9	(1,331)
Natural Family Planning	75.3	(73)
Traditional Methods	61.8	(986)

TABLE VII.5
Opinions about the Reliability of Oral Contraceptives
by Selected Characteristics, Women 15-44 Years of Age
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Certainty of Preventing Pregnancy					Total	Number of Respondents
	Totally Sure	Almost Sure	Fairly Sure	Not Sure At All	Do Not Know		
Total	14.3	38.4	16.7	19.2	11.5	100.0	(4,497)
Residence							
Bohemia	13.9	38.1	17.2	18.7	12.0	100.0	(2,737)
Moravia	14.8	38.8	15.9	19.8	10.8	100.0	(1,760)
Size of Place							
Less Than 5,000	14.5	36.7	15.1	18.9	14.8	100.0	(1,472)
5,000-19,999	16.2	36.2	16.0	19.7	11.9	100.0	(967)
20,000 +	13.3	40.7	18.2	19.1	8.9	100.0	(2,058)
Current Age							
15-19	10.0	41.7	14.1	17.7	16.5	100.0	(662)
20-24	12.5	44.6	16.8	17.7	8.4	100.0	(756)
25-29	16.1	42.1	16.7	15.7	9.4	100.0	(828)
30-34	16.8	34.6	19.1	16.7	12.8	100.0	(720)
35-39	15.5	34.1	17.7	22.5	10.3	100.0	(805)
40-44	15.8	33.2	16.6	23.6	10.8	100.0	(726)
Education							
Primary only	14.6	23.9	12.4	27.6	21.6	100.0	(738)
Secondary, No Diploma	13.3	38.4	14.1	21.7	12.6	100.0	(1,779)
Secondary, Diploma	16.3	43.1	19.9	13.8	6.9	100.0	(1,601)
Any University	10.0	51.6	26.7	9.3	2.4	100.0	(379)
Religion							
None	14.6	40.2	16.6	18.3	10.2	100.0	(2,913)
Catholic, Attends Services	9.3	36.0	14.9	21.1	18.6	100.0	(256)
Catholic, Does Not Attend	15.2	35.5	17.0	19.9	12.5	100.0	(1,134)
Other	11.0	30.6	19.2	25.6	13.7	100.0	(180)
Current contraception							
Not Using	11.5	35.7	14.5	21.5	16.7	100.0	(188)
Oral Contraceptives	34.3	44.5	13.6	7.1	0.5	100.0	(1,885)
Other Modern Method	14.1	39.1	19.7	17.7	9.4	100.0	(1,278)
Natural FP	9.9	42.0	21.0	21.0	6.2	100.0	(803)
Traditional Method	12.9	40.0	17.5	20.9	8.7	100.0	(253)

TABLE VII.6
Opinions about the Safety of Oral Contraceptives
by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Perceived Safety				Total	Number of Respondents
	Completely Safe	Fairly Safe	Unsafe	Do Not Know		
Total	8.0	52.1	21.6	18.4	100.0	(4,497)
Residence						
Bohemia	8.1	53.0	21.1	17.8	100.0	(2,737)
Moravia	7.8	50.7	22.3	19.2	100.0	(1,760)
Size of Place						
Less Than 5,000	8.4	48.7	20.0	22.9	100.0	(1,472)
5,000-19,999	6.7	52.7	21.0	19.5	100.0	(967)
20,000 +	8.2	54.3	23.0	14.4	100.0	(2,058)
Current Age						
15-19	7.1	50.4	18.3	24.1	100.0	(662)
20-24	9.5	61.3	15.0	14.2	100.0	(756)
25-29	10.0	54.6	19.7	15.6	100.0	(828)
30-34	8.5	51.4	22.2	17.9	100.0	(720)
35-39	8.1	47.4	26.2	18.3	100.0	(805)
40-44	5.4	48.7	27.6	18.3	100.0	(726)
Education						
Primary only	6.2	37.8	27.6	28.5	100.0	(738)
Secondary, No Diploma	7.7	49.4	21.7	21.1	100.0	(1,779)
Secondary, Diploma	9.0	60.9	18.0	12.1	100.0	(1,601)
Any University	9.3	60.9	22.5	7.3	100.0	(379)
Religion						
None	8.9	54.5	19.6	17.0	100.0	(2,913)
Catholic, Attends Services	3.7	40.7	34.2	21.4	100.0	(256)
Catholic, Does Not Attend	7.4	49.1	22.3	21.2	100.0	(1,134)
Other	3.7	48.4	31.1	16.9	100.0	(180)
Current contraception						
Not Using	6.6	46.6	22.1	24.9	100.0	(188)
Oral Contraceptives	23.4	70.6	4.4	1.7	100.0	(1,885)
Other Modern Method	7.9	53.9	22.0	16.2	100.0	(1,278)
Natural FP	3.7	51.9	29.6	14.8	100.0	(803)
Traditional Method	5.9	53.6	25.6	14.9	100.0	(253)

TABLE VII.7
 Opinions regarding Various Statements Concerning Oral Contraceptives
 1993 Czech Republic Reproductive Health Survey
 (Percent Distributions)

Statement about Oral Contraceptives	Agreement with Statement			Total
	Agree	Disagree	Do Not Know	
Cause gain weight	41.7	29.0	29.4	100.0
Make menstrual periods more regular	53.3	14.9	31.8	100.0
Cause nervousness	13.1	39.2	47.7	100.0
Taking them too long can cause infertility	16.0	38.6	45.3	100.0
Increase the risk of getting cancer	24.6	33.7	41.7	100.0
Are bad for blood circulation	20.6	23.5	55.9	100.0

Number of Respondents = 4,497

TABLE VII.8
Opinions regarding Various Statements Concerning Women's Roles and Reproduction
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Statement	Agreement with Statement			Total
	Agree	Disagree	Not Sure	
Care of children is women's work	25.0	74.6	0.4	100.0
A woman should be a virgin when she marries	7.1	87.3	5.6	100.0
A woman can become pregnant the first time she has intercourse	92.1	4.1	3.9	100.0
A condom should not be used more than one time	96.5	0.6	2.9	100.0
It is all right for a woman not to have children if she does not want to	86.6	8.6	4.9	100.0

Number of Respondents = 4,497

VIII. WOMEN'S HEALTH

All respondents were asked a series of questions regarding selected knowledge and behaviors related to the general health of women. The topics on which they were questioned included: receipt of gynecologic exams; breast self-examination; cigarette smoking; alcohol consumption; and height and weight. These topics yield information about risk factors for various chronic diseases and the degree to which women are being screened for certain health conditions.

Gynecologic Examinations

It is recommended that women of childbearing age receive complete gynecologic examinations yearly. In the Czech Republic, the proportion of women having such exams is thought to be a good estimate of the proportion who receive Pap smears to screen for cervical cancer, since Pap smears are a routine component of these examinations. Sixty percent of respondents reported that they receive gynecologic exams at least yearly ([Table VIII.1](#)). At the other extreme, 15% of women said that they had never had such an exam. Bohemian women were somewhat more likely to receive yearly exams than Moravian women (63% versus 56%), with the differences between regions within Bohemia and Moravia generally small. The group least likely to be examined was 15-19 year-olds, among whom only 30% had yearly exams and 59% never had them. Beyond ages 20-24, almost all women had exams at least occasionally. Education was strongly correlated with receiving gynecologic exams, although there was little difference between women with a secondary school diploma and those who attended a university.

Two-thirds of women who did not receive yearly exams said the reason for this was that they did not have any gynecologic problems ([Table VIII.2](#)). These women should be educated to the fact that all women ought to be examined regularly, and not just when problems are evident. The second most common reason given (13%) was that women did not like such exams. Another 9% felt that it was not necessary to go as often as yearly for exams.

Breast Self-Examination

Breast self-examination (BSE) can be a valuable tool, if done properly, for detecting possible breast tumors earlier than they might otherwise be found. In the CRRHS women were asked about their knowledge and use of this procedure. Slightly under half of all respondents said that they practiced BSE at least occasionally ([Table VIII.3](#)). Another 28% were familiar with BSE, but never practiced it and 26% were unfamiliar with BSE. Although there was little difference in reported BSE knowledge and practice according to place of residence, there were large differences by age and education. Use of BSE increased sharply with age, from 12% for 15-19 year-olds to 70% for 40-44 year-

olds. Use increased from 37% to 62% from the lowest to highest education categories.

Cigarette Smoking

It has been firmly established that cigarette smoking is an important risk factor for several forms of cancer, respiratory diseases, and cardiovascular disease, among other chronic conditions. Reducing smoking prevalence can play important role in increasing life expectancy and improving the overall health and quality of life of a population.

Thirty percent of CRRHS respondents reported that they currently smoked ([Table VIII.4](#)). Another 12% had smoked in the past, but were no longer smoking. Smoking prevalence was highest in North Bohemia (40%) and Prague (38%) and lowest in Moravia (26%) and adjacent East Bohemia (27%). Rates also tended to be higher in large cities than elsewhere. Smoking prevalence increased with age up to ages 30-34 and decreased sharply as educational level increased ([Figure VIII. 1](#)). Interestingly, cigarette use was strongly related to marital status, with previously married women having the highest prevalence of any population subgroup examined (48%). This finding warrants further investigation to explain the reasons behind it.


Only 19% of smokers were classified as light smokers, that is, they typically smoked fewer than five cigarettes per day ([Table VIII.5](#)). The largest proportion of smokers, 44%, smoked 10 to 19 cigarettes per day, while 16% were heavy smokers, smoking at least one pack per day. As a general rule, tobacco consumption was heavier among Bohemians than Moravians, among older women than younger women, and among poorly educated than well educated women. Previously married women not only had the highest smoking prevalence, but also were the heaviest smokers.

Alcohol Consumption

Because heavy alcohol consumption is also an important risk factor for a large number of health conditions, respondents were asked about the amount of beer, wine, and liquor they typically consumed. According to responses in the CRRHS, relatively few Czech women chronically consume large amounts of alcohol ([Table VIII.6](#)). Only 3% of respondents said that they consumed an average of more than seven alcoholic drinks per week. Moderate drinking is the norm, with almost eight of every ten women consuming either fewer than three or three to seven drinks per week, while one-fifth said they did not drink at all. The percentage of women in the highest consumption category was highest in Prague (5%) and North Bohemia (4%). It was not outstandingly high in any subgroup examined, being highest among previously married women (6%).

These findings should be interpreted cautiously, however. First, alcohol consumption can often be underreported because society frowns on heavy consumption. Second, these figures do not give any information on acute alcohol use, where individuals tend to consume large amounts of alcohol on single occasions.

Figure VIII.1
Cigarette Smoking Status, by Education
Percent Distribution, 1993 CRRHS



Overweight

Being overweight is another factor that substantially increases the risk of many chronic conditions, such as cardiovascular disease, hypertension, and diabetes. Respondents were asked to report their height and weight. These data were used to compute each woman's body mass index (BMI), which is equal to weight (in kilograms) divided by squared height (in meters) to determine the degree to which women were overweight. A woman was considered to be substantially overweight if her BMI was at least 28.3 (Centers for Disease Control, 1988). Overall, 15% of respondents met this criterion ([Table VIII.7](#)). This percentage was lowest in Prague (10%). It tended to be highest in places with a population of less than 5,000 people (21%). As expected the percentage increased with age, from only 3% for women ages 15 to 19 to 29% for those ages 40 to 44. The proportion overweight fell sharply as education increased, such that only 5% of women who attended university were overweight.

TABLE VIII.1
Frequency of Receipt of Gynecologic Exams,
by Region of Residence, Age, and Education
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Frequency of Exams				Total	Number of Women
	At Least Yearly	Every 1-3 Years	Every 3+ Years	Never		
Total	60.3	13.4	11.6	14.7	100.0	(4,497)
Region						
Bohemia	63.0	13.0	11.5	12.4	100.0	(2,737)
Prague	63.3	13.7	11.6	11.4	100.0	(456)
Central Bohemia	67.1	11.2	11.3	10.4	100.0	(460)
South Bohemia	64.6	10.1	14.5	10.7	100.0	(295)
West Bohemia	57.7	14.5	12.1	15.7	100.0	(413)
North Bohemia	63.7	12.0	12.6	11.7	100.0	(582)
East Bohemia	61.9	15.5	8.4	14.2	100.0	(531)
Moravia	56.1	14.0	11.7	18.1	100.0	(1,760)
South Moravia	54.4	15.3	12.9	17.5	100.0	(873)
North Moravia	57.9	12.8	10.5	18.8	100.0	(887)
Age						
15-19	29.5	7.1	4.4	59.0	100.0	(662)
20-24	69.2	13.9	5.1	11.8	100.0	(756)
25-29	74.0	15.1	8.3	2.5	100.0	(828)
30-34	68.8	15.6	13.6	2.0	100.0	(720)
35-39	64.3	15.8	18.6	1.4	100.0	(805)
40-44	64.3	14.3	19.7	1.7	100.0	(726)
Education						
Primary only	49.9	12.2	17.8	20.1	100.0	(738)
Secondary, No Dipl.	57.9	12.2	9.4	20.5	100.0	(1,779)
Secondary, Diploma	67.8	14.7	10.7	6.9	100.0	(1,601)
University	64.1	17.4	12.9	5.6	100.0	(379)

TABLE VIII.2
Primary Reason for Receiving Gynecologic Exams Less Than Yearly
1993 Czech Republic Reproductive Health Survey
(Percent Distribution)

Reason	Percent
No gynecological problems	64.4
Does Not Like Exam	13.3
Feels It Is Not Necessary To Go So Often	8.5
Lack of Time/Inconvenience	3.1
Forgets about It	3.1
Dislikes Staff/Facilities	1.6
Too Young/Still a Virgin	1.5
Waiting Time Is Too Long	1.1
Other Reasons	3.3
Total	100.0
<i>Number of Women</i>	<i>(1,660)</i>

TABLE VIII.3
Knowledge and Use of Breast Self-Examination (BSE)
according to Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristic	Knowledge/Use of Breast Self-Exam			Total	Number of Women
	Does Not Know BSE	Knows BSE, Does Not Use	Uses BSE		
Total	25.9	27.5	46.6	100.0	(4,497)
Region					
Bohemia	25.4	28.5	46.0	100.0	(2,737)
Moravia	26.6	25.9	47.5	100.0	(1,760)
Size of Place					
Less than 5,000	27.7	28.5	43.8	100.0	(1,472)
5,000-19,999	26.2	28.6	45.2	100.0	(967)
20,000 +	24.4	26.2	49.4	100.0	(2,058)
Age					
15-19	61.3	26.9	11.8	100.0	(662)
20-24	35.9	32.4	31.7	100.0	(756)
25-29	19.4	33.1	47.6	100.0	(828)
30-34	15.1	27.1	57.9	100.0	(720)
35-39	9.4	24.8	65.8	100.0	(805)
40-44	8.0	22.0	69.9	100.0	(726)
Education					
Primary only	39.7	23.3	36.9	100.0	(738)
Secondary, No Diploma	33.0	28.3	38.6	100.0	(1,779)
Secondary, Diploma	13.7	28.3	57.9	100.0	(1,601)
Any University	8.8	29.1	62.1	100.0	(379)

TABLE VIII.4
Current Cigarette Smoking Status, according to Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Smoking Status			Total	Number of Women
	Currently Smoke	Previously Smoked	Never Smoked		
Total	30.2	12.1	57.7	100.0	(4,497)
Region					
Bohemia	32.7	11.1	56.2	100.0	(2,737)
Prague	37.8	14.2	47.9	100.0	(456)
Central Bohemia	28.1	10.6	61.3	100.0	(460)
South Bohemia	34.2	12.2	53.6	100.0	(295)
West Bohemia	30.2	10.3	59.5	100.0	(413)
North Bohemia	39.5	8.7	51.8	100.0	(582)
East Bohemia	26.0	11.5	62.5	100.0	(531)
Moravia	26.3	13.7	60.0	100.0	(1,760)
South Moravia	25.6	14.7	59.7	100.0	(873)
North Moravia	27.1	12.7	60.2	100.0	(887)
Size of place					
Less than 5,000	26.9	12.0	61.2	100.0	(1,472)
5,000 to 19,999	27.5	13.6	58.9	100.0	(967)
20,000 +	33.8	11.6	54.6	100.0	(2,058)
Age Groups					
15-19	18.9	4.1	77.0	100.0	(662)
20-24	27.3	12.7	60.0	100.0	(756)
25-29	28.6	13.0	58.4	100.0	(828)
30-34	35.5	13.6	50.9	100.0	(720)
35-39	36.9	14.5	48.6	100.0	(805)
40-44	35.7	16.3	48.0	100.0	(726)
Marital Status					
Currently Married	31.6	14.7	53.6	100.0	(3,217)
Previously Married	47.8	14.1	38.1	100.0	(355)
Never Married	21.2	4.8	74.1	100.0	(925)
Education					
Primary only	40.8	10.7	48.5	100.0	(738)
Secondary, No Diploma	32.0	11.0	57.0	100.0	(1,779)
Secondary, Diploma	25.0	13.5	61.4	100.0	(1,601)
University	18.1	15.4	66.5	100.0	(379)

TABLE VIII.5
Average Number of Cigarettes Smoked per Day
according to Selected Characteristics
Women Who Currently Smoke
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Cigarettes per Day				Total	Number of Women
	<5	5-9	10-19	20+		
Total	18.6	21.6	43.5	16.2	100.0	(1,394)
Region						
Bohemia	16.0	20.5	44.4	19.0	100.0	(925)
Moravia	23.5	23.7	41.9	10.8	100.0	(469)
Size of place						
Less than 5,000	18.5	23.8	42.9	14.8	100.0	(404)
5,000 to 19,999	16.8	23.9	46.1	13.2	100.0	(274)
20,000 +	19.4	19.5	42.9	18.1	100.0	(716)
Age Groups						
15-19	30.1	28.4	32.3	9.0	100.0	(123)
20-24	22.9	26.0	40.7	10.4	100.0	(214)
25-29	20.5	19.7	49.6	10.1	100.0	(237)
30-34	14.4	23.1	43.9	18.6	100.0	(254)
35-39	14.5	20.5	43.0	21.9	100.0	(293)
40-44	15.0	16.1	48.1	20.8	100.0	(273)
Marital Status						
Currently Married	16.9	21.9	45.7	15.4	100.0	(1,024)
Previously Married	14.0	11.9	42.5	31.6	100.0	(168)
Never Married	28.2	28.9	35.7	9.2	100.0	(202)
Education						
Primary only	10.4	19.0	50.1	20.5	100.0	(322)
Secondary, No Diploma	21.5	21.8	41.7	15.0	100.0	(593)
Secondary, Diploma	19.7	22.4	42.3	15.6	100.0	(409)
University	28.4	29.7	33.8	8.1	100.0	(70)

TABLE VIII.6
Average Weekly Alcohol Consumption*, according to Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Average Number of Drinks per Week				Total	Number of Women**
	Non Drinker	0-2	3-7	8 +		
Total	20.8	44.4	32.0	2.8	100.0	(4,469)
Region						
Bohemia	20.2	44.1	32.6	3.1	100.0	(2,720)
Prague	21.4	38.8	35.2	4.7	100.0	(456)
Central Bohemia	19.5	47.2	32.2	1.1	100.0	(454)
South Bohemia	19.5	46.2	32.5	1.8	100.0	(290)
West Bohemia	30.2	42.0	25.3	2.5	100.0	(409)
North Bohemia	15.8	41.9	38.1	4.2	100.0	(581)
East Bohemia	17.3	49.1	30.1	3.5	100.0	(530)
Moravia	21.7	44.9	31.1	2.2	100.0	(1,749)
South Moravia	21.9	46.2	30.3	1.7	100.0	(872)
North Moravia	21.6	43.6	31.9	2.8	100.0	(877)
Size of place						
Less than 5,000	20.1	44.5	33.1	2.3	100.0	(1,465)
5,000 to 19,999	21.9	46.3	29.5	2.2	100.0	(956)
20,000 +	20.8	43.5	32.3	3.4	100.0	(2,048)
Age Groups						
15-19	29.9	44.7	23.0	2.4	100.0	(659)
20-24	21.8	47.9	28.9	1.4	100.0	(751)
25-29	19.0	46.5	32.2	2.2	100.0	(821)
30-34	19.3	44.3	33.6	2.9	100.0	(714)
35-39	14.5	43.7	38.1	3.7	100.0	(802)
40-44	18.9	40.1	37.1	3.9	100.0	(722)
Marital Status						
Currently Married	18.8	43.8	34.9	2.5	100.0	(3,197)
Previously Married	22.1	42.0	29.9	6.0	100.0	(353)
Never Married	25.7	46.7	25.1	2.5	100.0	(919)
Education						
Primary only	28.0	37.1	31.1	3.7	100.0	(736)
Secondary, No Diploma	22.7	45.7	29.0	2.6	100.0	(1,766)
Secondary, Diploma	15.8	46.9	35.5	1.9	100.0	(1,591)
University	15.8	44.3	34.5	5.4	100.0	(376)

* One drink is considered equal to one-half liter of beer, one-fifth liter of wine, or one-half deciliter of liquor.

**28 women did not respond to questions on alcohol consumption.

TABLE VIII.7
Percent of Respondents Who Are Substantially Overweight*
according to Selected Characteristics
1993 Czech Republic Reproductive Health Survey

Characteristics	Percent Overweight	Number of Women
Total	15.0	(4,497)
Region		
Bohemia	14.7	(2,737)
Prague	10.9	(456)
Central Bohemia	13.8	(460)
South Bohemia	18.6	(295)
West Bohemia	15.3	(413)
North Bohemia	17.5	(582)
East Bohemia	13.2	(531)
Moravia	15.3	(1,760)
South Moravia	15.6	(873)
North Moravia	15.0	(887)
Size of place		
Less than 5,000	20.5	(1,472)
5,000 to 19,999	12.3	(967)
20,000 +	12.1	(2,058)
Age Groups		
15-19	2.8	(662)
20-24	8.5	(756)
25-29	12.6	(828)
30-34	16.0	(720)
35-39	21.9	(805)
40-44	28.6	(726)
Marital Status		
Currently Married	18.2	(3,217)
Previously Married	19.3	(355)
Never Married	5.5	(925)
Education		
Primary only	26.2	(738)
Secondary, No Diploma	14.3	(1,779)
Secondary, Diploma	12.1	(1,601)
University	4.9	(379)

* Respondents were considered substantially overweight if their body mass index (weight in kilograms divided by the squared height in meters) was at least 28.3.

IX. TRANSMISSION OF HIV/AIDS

Since acquired immune deficiency syndrome (AIDS) was first recognized in the early 1980s, human immunodeficiency virus (HIV), the organism causing AIDS, has spread to all parts of the world. Even though, according to official statistics, the incidence of HIV infection was not yet very high in the Czech Republic at the time of the CRRHS, a threat of greatly increased spread of the infection existed. This was especially true in light of the increased numbers of people, especially tourists, entering the Czech Republic from other countries and the newfound freedom of Czechs to travel outside their own country following the fall of the communist government. The CRRHS included a series of questions intended to measure the knowledge of Czech women regarding HIV/AIDS in general and the means by which HIV is transmitted between individuals. Such information was designed to help determine where improvements in knowledge about AIDS was necessary in order to enhance the effectiveness of AIDS prevention programs.

All but five respondents (99.9%) reported that they had heard of AIDS. Every respondent familiar with AIDS was asked whether they thought HIV could be transmitted by each of 12 different means. The proportions of women who felt that these were possible modes of transmission are shown in [Table IX. 1](#). In general, there appeared to be a high level of awareness of those means by which HIV is known to be transmissible-sharing needles (98%), male homosexual relations (96%), and heterosexual relations (94%). Receiving a transfusion, which in some settings can lead to transmission, was felt to be a cause of AIDS by 89% of women. Almost half of women thought a person could be infected by giving blood or by going to a physician or dentist, behaviors that cannot transmit infection except under unusual circumstances. Among those methods asked about by which the virus is known not to be transmitted, the proportion who felt that HIV could be spread ranged from 28% for kissing on the mouth down to 3% for shaking hands. For everything except kissing on the mouth the proportion with correct knowledge tended to increase with increasing education.

Only about three-fourths of respondents felt that condoms provided excellent or good protection against transmission of HIV ([Table IX.2](#)). There were only small differentials according to the characteristics examined in women's opinions about the effectiveness of condoms. There was a slight increase in perceived effectiveness as education increased and little relationship with place of residence, age, or marital status.

Finally, even though CRRHS respondents were not asked about their own risk factors for contracting AIDS, they were asked their opinion about their own risk of becoming infected. Eighty-three percent of women felt they had no risk of becoming infected, 3% felt they had a low risk, and 5% felt they had a high risk ([Table IX.3](#)). There was very little variation in these percentages across the population. Well educated women were slightly more likely to see themselves as being at risk, but this is likely to reflect different considerations in assessing risk, rather than a higher incidence of risky behaviors.

TABLE IX.1
Percent of Respondents Who Think That the AIDS Virus Can Be
Transmitted in Various Ways, by Education of Respondent
1993 Czech Republic Reproductive Health Survey

Mode of Transmission	Total*	Education of Respondent			
		Primary only	Secondary, No Diploma	Secondary, Diploma	Any University
Using an <i>already</i> used needle	98.0	94.3	97.9	99.4	99.3
Homosexual relations among men	96.0	90.3	95.8	98.2	98.3
Heterosexual relations	93.7	88.5	93.4	95.6	97.3
Receiving a transfusion	89.2	84.7	89.5	90.6	90.7
Giving blood	46.1	52.8	49.7	40.0	36.9
Going to a physician or dentist	43.7	41.9	41.5	46.0	48.2
Kissing on the mouth	27.6	30.0	25.5	27.8	32.0
Sharing objects with AIDS patient	21.4	31.2	23.2	16.6	10.0
Mosquito bites	18.7	22.3	21.1	15.3	11.2
Using public restrooms	14.3	19.5	14.5	12.1	10.5
Shaking hands	3.0	6.4	2.8	2.1	0.7
<i>Number of Respondents</i>	<i>(4,492)</i>	<i>(738)</i>	<i>(1,779)</i>	<i>(1,601)</i>	<i>(379)</i>

*Five respondents who had not heard of AIDS were excluded.

TABLE IX.2
Opinions about the Effectiveness of Condoms
in Protecting against HIV Infection, by Selected Characteristics
1993 Czech Republic Reproductive Health Survey
(Percent Distributions)

Characteristics	Perceived Effectiveness of Condoms					Total	No. of Women
	Excellent	Good	Moderate	Poor	Don't Know		
Total	32.8	41.0	14.6	5.7	5.8	100.0	(4,497)
Region							
Bohemia	32.3	42.7	14.0	5.1	5.8	100.0	(2,737)
Moravia	33.6	38.5	15.4	6.5	5.9	100.0	(1,760)
Size of place							
Less than 5,000	33.8	39.6	13.9	5.5	7.2	100.0	(1,472)
5,000 to 19,999	31.7	40.9	15.4	5.7	6.3	100.0	(967)
20,000 +	32.5	42.0	14.7	5.8	5.0	100.0	(2,058)
Age							
15-19	30.3	39.4	16.0	6.5	7.8	100.0	(662)
20-24	29.4	42.9	17.9	5.7	4.1	100.0	(756)
25-29	32.7	44.5	14.3	4.7	3.8	100.0	(828)
30-34	32.0	42.0	13.2	4.4	8.3	100.0	(720)
35-39	36.8	38.3	12.2	6.0	6.7	100.0	(805)
40-44	35.2	39.8	13.7	6.4	4.9	100.0	(726)
Marital Status							
Currently Married	34.2	41.1	13.9	5.6	5.2	100.0	(3,217)
Previously Married	32.9	39.9	13.6	5.9	7.7	100.0	(355)
Never Married	29.0	41.1	16.6	5.9	7.4	100.0	(925)
Education							
Primary only	35.1	30.4	13.6	7.1	13.6	100.0	(738)
Secondary, No Dipl.	32.3	39.7	14.9	6.7	6.3	100.0	(1,779)
Secondary, Diploma	33.3	46.0	14.2	4.0	2.5	100.0	(1,601)
Any University	27.1	50.6	16.4	4.2	1.7	100.0	(379)

TABLE IX.3
 Respondents' Opinions about Their Own Risk of Contracting AIDS
 according to Selected Characteristics
 1993 Czech Republic Reproductive Health Survey
 (Percent Distributions)

Characteristics	Perceived Risk of Getting AIDS				Total	No. of Women
	No Risk	Low Risk	High Risk	Do Not Know		
Total	83.4	3.1	5.1	8.4	100.0	(4,497)
Region						
Bohemia	83.6	2.9	4.9	8.6	100.0	(2,737)
Moravia	82.9	3.4	5.4	8.3	100.0	(1,760)
Size of place						
Less than 5,000	84.8	1.9	4.2	9.1	100.0	(1,472)
5,000 to 19,999	83.1	3.1	5.1	8.7	100.0	(967)
20,000 +	82.3	4.0	5.8	7.9	100.0	(2,058)
Age						
15-19	77.4	4.0	6.9	11.6	100.0	(662)
20-24	83.7	3.2	5.7	7.4	100.0	(756)
25-29	86.5	2.3	3.6	7.6	100.0	(828)
30-34	85.6	2.0	3.5	8.9	100.0	(720)
35-39	84.7	3.0	4.5	7.8	100.0	(805)
40-44	83.7	3.7	5.6	7.0	100.0	(726)
Marital Status						
Currently Married	86.0	2.5	4.2	7.3	100.0	(3,217)
Previously Married	80.9	4.2	4.8	10.1	100.0	(355)
Never Married	77.2	5.3	7.1	10.4	100.0	(925)
Education						
Primary only	84.4	2.7	3.9	9.0	100.0	(738)
Secondary, No Dipl.	84.1	2.9	4.4	8.6	100.0	(1,779)
Secondary, Diploma	82.9	3.7	5.6	7.8	100.0	(1,601)
Any University	78.5	2.7	9.3	9.5	100.0	(379)

REFERENCES

- Albert A, Bennett C, Bojar M. 1992. Health care in the Czech Republic: a system in transition. *Journal of the American Medical Association*, 267(18):2461-6.
- Althaus F. 1992. Replacement-level fertility has now become the rule in most of Eastern Europe. *Family Planning Perspectives*, 24(3):140-2.
- Blayo C. 1991. Contraceptive methods in Eastern Europe. *Population*, 46(3):527-46.
- Blayo C. 1993. Le role de l'avortement dans les pays d'Europe Centrale et Orientale. *Proceedings of the 1993 International Population Conference. IUSSP, Liege Belgium*, pp 235-252.
- Centers for Disease Control. 1988. Prevalence of Overweight in Selected States-Behavioral Risk Factor Survey. *Morbidity and Mortality Weekly Report*, 37:9-11.
- Czechoslovakia Federal Statistical Office. 1991. Preliminary Results of the Population and Housing Census. Prague.
- David HP. 1992. Abortions in Europe, 1920-91: a public health perspective. *Studies in Family Planning*, 23(1):1-22.
- David HP, Dytrych Z, Matejcek Z, Schuller V. 1988. Born Unwanted: Developmental Effects of Denied Abortion. Avicenum, Czechoslovakian Medical Press, Prague.
- Festy P. 1991. Fertility in Eastern Europe since 1970. *Population*, 46(3):470-509.
- Fialova L, Pavlik Z, Veres P. 1990. Fertility decline in Czechoslovakia during the last two centuries. *Population Studies*, 44(1):89-106.
- Hatcher RA, Trussell J, Stewart F, et al. Contraceptive Technology, sixteenth revised edition. Irvington Press, New York, pp 633-687.
- Institute of Health Information and Statistics. 1991. CSFR Health Statistics Yearbook 1991. IHSI, Prague.
- Ketting E. 1992. Family planning in Central and Eastern Europe. *Planned Parenthood Europe (special issue)*, 21(2): 1-39.
- Kraus J. 1987. Research on family size planning--I. *Demographie*, 29(1):23-33.

Massaro TA, Nemec J, Kalman I. 1994. Health system reform in the Czech Republic. *Journal of the American Medical Association*, 271:1870.

National Public Health and Hospital Institute. 1994. Partnerships in Health Care Central and Eastern Europe.

Population Reference Bureau. 1994a, World population data sheet.

Population Reference Bureau. 1994b, The World's Youth (data sheet).

Raffel MW, Raffel NK. 1992. Czechoslovakia's changing health care system. *Public Health Reports*, 107(6):636-43.

Romanian Institute for Mother and Child Care, Centers for Disease Control and Prevention. 1995. Reproductive Health Survey Romania 1993.

Trussell J, Kost K. 1987. Contraceptive failure in the U.S.: a critical review of the literature. *Studies in Family Planning*, 18(5):237-283.

United Nations. 1988-1992. Demographic Yearbook (various years). United Nations, New York.

Uzel R, Ketting E, et al. 1992. Contraceptive Practice and attitudes of czech and Slovak women. *Planned Parenthood Europe* 21(2):14-17.

Visser A, Uzel R, et al. 1993. Attitudes of Czech and Slovak gynecologists toward family planning. *Advances in Contraception* 9:351-362.

World Fertility Survey. 1978. The Czechoslovak Fertility Survey. 1972. International Statistical Institute, Voorburg, Netherlands.

APPENDIX A

SAMPLING ERRORS

Survey estimates of percentages, proportions, means, or other measures are subject to two types of errors: sampling error and non-sampling error. Non-sampling errors can arise from many sources, most notably interviewer errors, inaccurate information given by respondents (intentionally or unintentionally), data processing errors, failure to administer the interview to the correct households or individuals, as well as other factors. Although efforts were made during the design, training and implementation of the 1993 CRRHS to minimize the incidence and severity of non-sampling errors, it is virtually impossible to eliminate all sampling error or to evaluate their importance statistically.

Sampling error is a measure of the variability between all possible samples of a given size that could have been selected from the population being studied (in this case, women between the ages of 15 and 44 throughout the Czech Republic) using a given sample design. Sampling error is measured in terms of the *standard error* for a particular statistic, which is the square root of the variance of that statistic. The standard error can be used to calculate *confidence intervals* around estimated statistics. The confidence interval most commonly used is the 95 percent interval, which defines the range within which the actual percentage, mean, etc. falls, in the absence of all non-sampling error. The 95 percent confidence interval for a statistic is the estimated value plus or minus 1.96 (approximately two) times the standard error of the estimate.

The standard errors of statistics estimated using a multistage sample design, such as that used in the 1993 CRRHS are more complicated to calculate and are larger than standard errors based on simple random samples. The software package SUDAAN (Research Triangle Institute, 1993) was used to compute the CRRHS standard errors with the appropriate statistical methodology. In addition to standard errors, SUDAAN was used to compute the *design effect* for each estimate, which is defined as the estimated ratio between the variance using the multistage cluster design actually used and the variance that would have resulted if a simple random sample had been used. Therefore, the design effect demonstrates the increase in the increased variability of an estimate (including the standard error and confidence interval) due to the use of a complex survey design.

Standard errors, design effects, and 95% confidence intervals (the limits of which are $P-2SE$ and $P+2SE$) are presented in [Table A.2](#) for variables considered to be of major interest. For each variable, the definition and the population used for the calculation are presented in [Table A.1](#).

TABLE A.1
Variables for Which Sampling Errors Were Calculated
and Base Population Used for Calculation

Variable	Population Used for Calculation
% with secondary diploma or any university	All respondents
% who have attended university	All respondents
% ever married or in union	All respondents
% currently married or in union	All respondents
% who want no more children	Fecund women currently married/in union
% of pregnancies planned	All reported pregnancies ending after 1987 and current pregnancies
% currently using any contraception	Women currently married/in union
% currently using modern contraception	Women currently married/in union
% currently using traditional contraception*	Women currently married/in union
% currently using an IUD	Women currently married/in union
% currently using condoms only	Women currently married/in union
% currently using oral contraceptives	Women currently married/in union
% currently using withdrawal only	Women currently married/in union
% in need of family planning (definition 1)**	All respondents
% in need of family planning (definition 2)**	All respondents
% for whom health concerns were important	Women currently using traditional methods*
% with prenatal care in first trimester	Last pregnancies ending in live birth after 1987
% with ultrasound examination	Last pregnancies ending in live birth after 1987
% hospitalized during pregnancy	Last pregnancies ending in live birth after 1987
% who smoked during pregnancy	Last pregnancies ending in live birth after 1987
% low birthweight	Last pregnancies ending in live birth after 1987
% breastfed	Last pregnancies ending in live birth after 1987
% of 15-17 year-olds who are sexually experienced	All 15-17 year-old respondents
% of 18-19 year-olds who are sexually experienced	All 18-19 year-old respondents
% who used contraception at first sexual intercourse	Sexually experienced 15-24 year-old respondents
% who think women should decide whether to have abortion	All respondents
% receiving yearly gynecologic examination	All respondents
% who currently smoke cigarettes	All respondents
% who are substantially overweight	All respondents

* Traditional contraception includes withdrawal, natural family planning, and douche.

**For definitions of need for family planning see Table IV.12.

TABLE A.2
Sampling Errors for Selected Variables
1993 Czech Republic Reproductive Health Survey

Variable	Percentage	Standard Error	Design Effect	P-2SE*	P+2SE*	N
With secondary diploma	41.34	1.05	2.05	39.24	43.44	4,497
Attended university	7.58	0.48	1.45	6.62	8.54	4,497
Ever married/in union	74.27	0.94	2.07	72.15	76.39	4,497
Currently married/in union	66.78	0.96	1.86	64.86	68.70	4,497
Want no more children	69.30	1.00	1.43	67.30	71.30	3,025
Pregnancy planned	65.94	1.31	1.99	63.32	68.56	2,591
Currently use any contraception	69.07	1.19	2.13	66.69	71.45	3,217
Currently use modern contraception	42.89	1.01	1.33	40.87	44.91	3,217
Currently use traditional contraception	26.10	1.14	2.16	23.82	28.34	3,217
Currently use IUD	15.32	0.76	1.44	13.80	16.84	3,217
Currently use condoms	16.66	0.76	1.35	15.14	18.18	3,217
Currently use oral contraceptives	8.05	0.55	1.32	6.95	9.15	3,217
Currently use withdrawal	22.18	1.08	2.15	20.02	24.34	3,217
In need of family planning (Def 1)	9.77	0.61	1.90	8.55	11.99	4,497
In need of family planning (Def 2)	30.55	0.85	1.54	28.85	32.25	4,497
Health concerns important factor	74.56	1.97	1.69	70.62	78.50	830
Prenatal care in first trimester	94.13	0.76	1.44	92.61	95.65	1,382
Ultrasound exam	91.06	0.87	1.27	89.32	92.80	1,382
Hospitalized during pregnancy	28.91	1.29	1.13	26.33	31.49	1,382
Smoked during pregnancy	11.52	1.03	1.44	9.46	13.58	1,382
Low birthweight (<2500 grams)	5.38	0.66	1.19	4.06	6.70	1,382
Breastfed	90.66	0.77	1.20	89.12	92.20	1,691
15-17 years, sexually experienced	35.42	2.62	1.08	30.18	40.66	359
18-19 years, sexually experienced	81.64	2.61	1.30	76.42	86.86	287
Used contraception at 1st sex	56.64	1.93	1.64	52.78	60.50	1,080
Women should make abortion choice	85.22	0.77	2.11	83.68	86.76	4,497
Yearly gynecologic exam	60.29	1.02	1.94	58.25	62.33	4,497
Currently smoke cigarettes	30.16	0.87	1.63	28.42	31.90	4,497
Overweight	11.22	0.55	1.37	10.12	12.32	4,497

*P-2SE and P+2SE are the approximate lower and upper 95% confidence limits of the percentage for a given variable, taking the survey design into account.

APPENDIX B

1993 CZECH REPUBLIC REPRODUCTIVE HEALTH SURVEY Household questionnaire

ID NUMBER _ _ _ _

REGION _____

DISTRICT _____

BASIC CENSUS DISTRICT _____

HOUSE NUMBER (red number) _____

BUILDING NUMBER _____

FLAT NUMBER _____

STREET ADDRESS _____

CITY/TOWN/VILLAGE _____

SIZE OF PLACE

- 1 LESS THAN 5,000
- 2 5,000-19,999
- 3 20,000+

VISIT RECORD

Visit number	1		2		3		4	
	Day	Month	Day	Month	Day	Month	Day	Month
Date of visit	—	—	—	—	—	—	—	—
Result*	—		—		—		—	
Interviewer	—	—	—	—	—	—	—	—
Supervisor	—	—	—	—	—	—	—	—

*RESULT CODES

- 1 Completed interview
- 2 No eligible females
- 3 Nobody at home
- 4 Selected respondent not home
- 5 Total refusal
- 6 Refusal by selected respondent
- 7 Unoccupied house
- 8 Respondent incompetent _____
- 9 Other _____

1. How many families live in this household?
 __ families
2. How many people normally live in this household?
 __ __ people
3. How many females between the ages of 15 and 44 live in this household?
 __ females
4. For each of these women could you give me the following information:

Line	First name	Age	Marital status	Education	
				Level	Grade
1	_____	__ __	__	__	__ __
2	_____	__ __	__	__	__ __
3	_____	__ __	__	__	__ __
4	_____	__ __	__	__	__ __
5	_____	__ __	__	__	__ __
6	_____	__ __	__	__	__ __

Marital status:	Level:
1 Married	0 No school
2 Divorced	1 Primary
3 Separated	2 Secondary
4 Widowed	3 University
5 Single	8 Don't know
6 Consensual union	
8 Don't know	

SELECTION OF INDIVIDUAL RESPONDENT:

	LAST DIGIT OF QUESTIONNAIRE									
	0	1	2	3	4	5	6	7	8	9
Eligible Respondents										
2	1	2	1	2	1	2	1	2	1	2
3	3	1	2	3	1	2	3	1	2	3
4	3	4	1	2	3	4	1	2	3	4
5	1	2	3	4	5	1	2	3	4	5
6	6	1	2	3	4	5	6	1	2	3

1993 CZECH REPUBLIC REPRODUCTIVE HEALTH SURVEY
Individual questionnaire

TIME STARTED: __ __ : __ __

ID NUMBER __ __ __ __

Background characteristics

100. In what month and year were you born?

Month __ __

Year 19 __ __

101. How old are you?

__ __ years old

(MAKE SURE THAT AGE AND DATE OF BIRTH CORRESPOND)

102. What is the highest grade in school you completed?

1 Primary 1 2 3 4 5 6 7 8 9-->GO TO Q104

2 Secondary 1 2 3 4 5+ ----- >GO TO Q103

3 University/Faculty 1 2 3 4 5+ ----- >GO TO Q104

103. Did you receive a diploma?

1 Yes

2 No

104. Are you currently married, living with someone as if married, separated, divorced, widowed, or single?

1 Married \
2 Living with a man \
3 Separated /--GO TO Q106
4 Divorced /
5 Widowed /
6 Single --- >GO TO Q105

105. Have you ever been married or lived with a man?

1 Yes--->CORRECT Q104 AND GO TO Q106

2 No-->GO TO 200 SERIES

106. How many times have you been married or lived with a man?

__ times

107. In what month and year did you begin living with your
(first)
husband/partner?

Month __ __

Year 19 __ __

108. What was the highest grade in school that your (first) husband/partner completed?

1 Primary 1 2 3 4 5 6 7 8 9---->GO TO Q110

2 Secondary 1 2 3 4 5+ ----->GO TO Q109

3 University/Faculty 1 2 3 4 5+ ----->GO TO Q110

8 Don't know ----->GO TO Q110

109. Did he receive a diploma?

1 Yes

2 No

110. When you first got married, how many children did you plan to have?

 Children

66 As many as possible

88 Were not sure

Fertility/Pregnancy

200. Have you ever been pregnant?

- 1 Yes
- 2 No
- 3 Not sure

201. Are you currently pregnant?

- 1 Yes ----->GO TO Q201A
- 2 No
- 3 Not sure

IF CURRENTLY PREGNANT (Q201=1) GO TO Q201A

IF PREGNANT IN THE PAST (Q200=1 AND Q201 NOT 1) GO TO 202.

IF NEVER PREGNANT (NEITHER Q200=1 NOR Q201=1) GO TO Q224.

201A. Thinking back to when you became pregnant, did you want to become pregnant then?

- 1 Yes ----->GO TO Q202
- 2 No ----->GO TO Q201B
- 3 Not sure ->GO TO Q202

201B. Was it that you wanted to wait longer to become pregnant or that you wanted no more children?

- 1 Wanted to wait longer
- 2 Wanted no more children
- 3 Not sure

202. Have you ever had any live-born children?

- 1 Yes
- 2 No --->GO TO PREGNANCY HISTORY, Q210

203. How many living children do you have, including those who do not live with you?

__ __ children

IF 0 LIVING CHILDREN, GO TO Q206

204. How many of these are boys?

__ __ boys

205. How many of these are girls?

__ __ girls

MAKE SURE THAT THE NUMBER OF BOYS AND GIRLS ADD UP TO THE TOTAL NUMBER OF CHILDREN (Q204 + Q205 = Q203).

206. Have you ever had any live-born children who later died, including any who lived only a very short time after birth?

- 1 Yes
- 2 No ---->GO TO Q209

207. How many boys have died?

— boys

208. How many girls have died?

— girls

209. So, altogether you have had __ __ (Q203 + Q207 + Q208) live births?

- 1 Yes
- 2 No--->PROBE AND CORRECT Q203, Q207, Q208 AS NECESSARY

PREGNANCY HISTORY

Now I would like to talk to you about your pregnancies. Please make sure you include all past pregnancies, regardless of when they occurred and how they ended, whether in a live birth, an abortion, a miscarriage, or a stillbirth. Starting with your most recent pregnancy, please give me the following information:

210	210A	211	212	213	214	215	216
Order	Was this a multiple pregnancy?	When did this pregnancy end? (Month & year)	How many months did this pregnancy last?	How did this pregnancy end?	Was this child a boy or a girl?	Is this child still alive?	At ¹ what age did he/ she die?
1	1=Single 2=Twins 3 = 3 + 8=DK	Month __ __ Year 19 __ __	__ months 0=Don't know 9 = 9 +	1 Live birth-->Q214 2 Stillbirth-->NEXT PREGNANCY 3 Miscarriage-->NEXT PREGNANCY 4 Extrauterine->NEXT PREGNANCY 5 Induced abortion-->NEXT PREG	1 Boy 2 Girl	1 Alive- >NEXT PREG 2 Dead	__ __ Years __ __ Months
2	1=Single 2=Twins 3 = 3 + 8=DK	Month __ __ Year 19 __ __	__ months 0=Don't know 9 = 9 +	1 Live birth-->Q214 2 Stillbirth-->NEXT PREGNANCY 3 Miscarriage-->NEXT PREGNANCY 4 Extrauterine->NEXT PREGNANCY 5 Induced abortion-->NEXT PREG	1 Boy 2 Girl	1 Alive- >NEXT PREG 2 Dead	__ __ Years __ __ Months
3	1=Single 2=Twins 3 = 3 + 8=DK	Month __ __ Year 19 __ __	__ months 0=Don't know 9 = 9 +	1 Live birth-->Q214 2 Stillbirth-->NEXT PREGNANCY 3 Miscarriage-->NEXT PREGNANCY 4 Extrauterine->NEXT PREGNANCY 5 Induced abortion-->NEXT PREG	1 Boy 2 Girl	1 Alive- >NEXT PREG 2 Dead	__ __ Years __ __ Months
4	1=Single 2=Twins 3 = 3 + 8=DK	Month __ __ Year 19 __ __	__ months 0=Don't know 9 = 9 +	1 Live birth-->Q214 2 Stillbirth-->NEXT PREGNANCY 3 Miscarriage-->NEXT PREGNANCY 4 Extrauterine->NEXT PREGNANCY 5 Induced abortion-->NEXT PREG	1 Boy 2 Girl	1 Alive- >NEXT PREG 2 Dead	__ __ Years __ __ Months

5	1=Single 2=Twins 3 = 3 + 8=DK	Month __ __ Year 19 __ __	__ months 0=Don't know 9 = 9 +	1 Live birth-- >Q214 2 Stillbirth-- >NEXT PREGNANCY 3 Miscarriage-->NEXT PREGNANCY 4 Extrauterine->NEXT PREGNANCY 5 Induced abortion-->NEXT PREG	1 Boy 2 Girl	1 Alive- >NEXT PREG 2 Dead	__ __ Years __ __ Months
6	1=Single 2=Twins 3 = 3 + 8=DK	Month __ __ Year 19 __ __	__ months 0=Don't know 9 = 9 +	1 Live birth-- >Q214 2 Stillbirth-- >NEXT PREGNANCY 3 Miscarriage-->NEXT PREGNANCY 4 Extrauterine->NEXT PREGNANCY 5 Induced abortion-->NEXT PREG	1 Boy 2 Girl	1 Alive- >NEXT PREG 2 Dead	__ __ Years __ __ Months

210	210A	211	212	213	214	215	216
Order	Was this a multiple pregnancy ?	When did this pregnancy end? (Month & year)	How many months did this pregnancy last?	How did this pregnancy end?	Was this child a boy or a girl?	Is this child still alive?	At ¹ what age did he/ she die?
7	1=Single 2=Twins 3 = 3 + 8=DK	Month __ __ Year 19 __ __	__ months 0=Don't know 9 = 9 +	1 Live birth-- >Q214 2 Stillbirth-- >NEXT PREGNANCY 3 Miscarriage-->NEXT PREGNANCY 4 Extrauterine->NEXT PREGNANCY 5 Induced abortion-->NEXT PREG	1 Boy 2 Girl	1 Alive- >NEXT PREG 2 Dead	__ __ Years __ __ Months
8	1=Single 2=Twins 3 = 3 + 8=DK	Month __ __ Year 19 __ __	__ months 0=Don't know 9 = 9 +	1 Live birth-- >Q214 2 Stillbirth-- >NEXT PREGNANCY 3 Miscarriage-->NEXT PREGNANCY 4 Extrauterine->NEXT PREGNANCY 5 Induced abortion-->NEXT PREG	1 Boy 2 Girl	1 Alive- >NEXT PREG 2 Dead	__ __ Years __ __ Months
9	1=Single 2=Twins 3 = 3 + 8=DK	Month __ __ Year 19 __ __	__ months 0=Don't know 9 = 9 +	1 Live birth-- >Q214 2 Stillbirth-- >NEXT PREGNANCY 3 Miscarriage-->NEXT PREGNANCY 4 Extrauterine->NEXT PREGNANCY 5 Induced abortion-->NEXT PREG	1 Boy 2 Girl	1 Alive- >NEXT PREG 2 Dead	__ __ Years __ __ Months
10	1=Single 2=Twins 3 = 3 + 8=DK	Month __ __ Year 19 __ __	__ months 0=Don't know 9 = 9 +	1 Live birth-- >Q214 2 Stillbirth-- >NEXT PREGNANCY 3 Miscarriage-->NEXT PREGNANCY 4 Extrauterine->NEXT PREGNANCY 5 Induced abortion-->NEXT PREG	1 Boy 2 Girl	1 Alive- >NEXT PREG 2 Dead	__ __ Years __ __ Months
11	1=Single 2=Twins 3 = 3 + 8=DK	Month __ __ Year 19 __ __	__ months 0=Don't know 9 = 9 +	1 Live birth-- >Q214 2 Stillbirth-- >NEXT PREGNANCY 3 Miscarriage-->NEXT PREGNANCY 4 Extrauterine->NEXT PREGNANCY 5 Induced abortion-->NEXT PREG	1 Boy 2 Girl	1 Alive- >NEXT PREG 2 Dead	__ __ Years __ __ Months

12	1=Single 2=Twins 3 = 3 + 8=DK	Month __ __ Year 19 __ __	__ months 0=Don't know 9 = 9 +	1 Live birth-- >Q214 2 Stillbirth-- >NEXT PREGNANCY 3 Miscarriage-->NEXT PREGNANCY 4 Extrauterine->NEXT PREGNANCY 5 Induced abortion-->NEXT PREG	1 Boy 2 Girl	1 Alive- >NEXT PREG 2 Dead	__ __ Years __ __ Months
----	--	------------------------------------	--------------------------------------	--	-----------------	----------------------------------	-----------------------------

AFTER FILLING IN ALL PREGNANCIES: IF NO PREGNANCIES ENDED SINCE THE BEGINNING OF 1988 GO TO 300 SERIES.
IF ANY PREGNANCIES ENDED SINCE THE BEGINNING OF 1988 GO TO Q217.

QUESTIONS 217-223 ONLY FOR PREGNANCIES THAT ENDED IN 1988 OR LATER

	217	218	219	220	221	222	223
COPY LINE # FROM Q210	Thinking back to when you became pregnant that time, did you want to become pregnant?	Was it that you wanted to wait longer to become pregnant or that you wanted no more children?		Did you breastfeed him/her?	Are you still breastfeeding?	How many months did you breastfeed?	What was the most important reason that you decided to have an abortion? (CODES BELOW)
— —	1 Yes---- >Q219 2 No.... >Q218 3 Not sure---- >Q219	1 Wait longer 2 Wanted no more 3 Not sure	IF LIVE BIRTH -- >Q220 IF INDUCED ABORTION -- >Q223 IF NOT A LIVE BIRTH -- >NEXT LINE	1 Yes 2 No-- >NEXT LINE	1 Yes --- >NEXT LINE 2 No	___ months- ->NEXT LINE	— —
— —	1 Yes---- >Q219 2 No.... >Q218 3 Not sure---- >Q219	1 Wait longer 2 Wanted no more 3 Not sure	IF LIVE BIRTH -- >Q220 IF INDUCED ABORTION -- >Q223 IF NOT A LIVE BIRTH -- >NEXT LINE	1 Yes 2 No-- >NEXT LINE	1 Yes --- >NEXT LINE 2 No	___ months- ->NEXT LINE	— —
— —	1 Yes---- >Q219 2 No.... >Q218 3 Not sure---- >Q219	1 Wait longer 2 Wanted no more 3 Not sure	IF LIVE BIRTH -- >Q220 IF INDUCED ABORTION -- >Q223 IF NOT A LIVE BIRTH -- >NEXT LINE	1 Yes 2 No-- >NEXT LINE	1 Yes --- >NEXT LINE 2 No	___ months- ->NEXT LINE	— —
— —	1 Yes---- >Q219 2 No.... >Q218 3 Not sure---- >Q219	1 Wait longer 2 Wanted no more 3 Not sure	IF LIVE BIRTH -- >Q220 IF INDUCED ABORTION -- >Q223 IF NOT A LIVE BIRTH -- >NEXT LINE	1 Yes 2 No-- >NEXT LINE	1 Yes --- >NEXT LINE 2 No	___ months- ->NEXT LINE	— —
— —	1 Yes---- >Q219 2 No.... >Q218 3 Not sure---- >Q219	1 Wait longer 2 Wanted no more 3 Not sure	IF LIVE BIRTH -- >Q220 IF INDUCED ABORTION -- >Q223 IF NOT A LIVE BIRTH -- >NEXT PAGE	1 Yes 2 No-- >NEXT PAGE	1 Yes --- >NEXT PAGE 2 No	___ months- ->NEXT PAGE	— —

CODES FOR Q223:

1 Delivery dangerous to her health/life
2 Fetus diagnosed with defect/high risk of

3 Had all the children she/they wanted
4 Wanted to wait longer for next child
5 Could not afford another child
6 Husband/Partner wanted her to have abortion
7 Not married/No partner
8 Relationship ended
9 Parents wanted her to have an abortion
20 Other

88 Don' t know
99 No response

defect

(specify) _____

GO TO NEXT PAGE (300 SERIES)

224. Have you ever had any pregnancies that ended with an abortion, a miscarriage, or a stillbirth?
1 Yes -->RETURN TO PREGNANCY HISTORY, Q202
2 No

Family Planning

For each of the following methods of preventing pregnancy, please tell me:

METHOD	300. Have you ever heard of it?	301. Have you ever used it?	302. Do you know where to get it?
A. Pills	1 Yes-->Q301 2 No--->B	1 Yes 2 No	1 Yes 2 No
B. IUD	1 Yes-->Q301 2 No-->C	1 Yes 2 No	1 Yes 2 No
C. Condoms	1 Yes-->Q301 2 No-->D	1 Yes 2 No	1 Yes 2 No
D. Foam/Jelly/ Cream	1 Yes-->Q301 2 No--->E	1 Yes 2 No	1 Yes 2 No
E. Diaphragm	1 Yes-->Q301 2 No--->F	1 Yes 2 No	1 Yes 2 No
F. Female Sterilization	1 Yes-->Q301 2 No-->G	1 Yes-->G 2 No-->Q302	1 Yes 2 No
G. Male Ster. (Vasectomy)	1 Yes-->Q301 2 No-->H	1 Yes 2 No	
H. Fertile days method (Rhythm)	1 Yes-->Q301 2 No-->I	1 Yes 2 No	
I. Withdrawal	1 Yes-->Q301 2 No-->J	1 Yes 2 No	
J. Other	1 Yes-->Q301 2 No-->Q303	1 Yes 2 No	

303. RECORD WHETHER RESPONDENT REPORTS HAVING USED ANY METHOD (ANY 1 FOR Q301)

- 1 Never used - >GO TO Q304
- 2 Ever used--- >GO TO Q305

304. So, you have never used any of the methods of preventing pregnancy that I just mentioned with any partner?

- 1 Never used - >GO TO Q328
- 2 Ever used - >CORRECT Q301 AND GO TO Q305

305. Are you (or your partner) currently using any method or doing anything to prevent pregnancy?

- 1 Yes
- 2 No--->GO TO Q321

306. What method are you using?

- 1 Pills -- >GO TO Q308
- 2 IUD ---- >GO TO Q308
- 3 Condoms ---- >GO TO Q309
- 4 Foam/Jelly/Cream ---- >GO TO Q308
- 5 Diaphragm ---- >GO TO Q308
- 6 Tubal ligation -- >GO TO Q307
- 7 Vasectomy ----- >GO TO Q307
- 8 Fertile days ---- >GO TO Q311
- 9 Withdrawal _____"
- 10 Other _____"
- 11 Condoms + Foam/Jelly --- >Q309
- 12 Other combination _____"

307. In what month and year was this operation performed?

Month — —
Year 19 — —

GO TO Q321

308. Do you and your partner ever use a condom in addition to the method you are using?

- 1 Yes
- 2 No

309. Where do you get your family planning method?

- 1 Regional/District Physician's office
- 2 Polyclinic physician's office
- 3 Hospital
- 4 Private practice physician
- 5 Pharmacy
- 6 Condom machine
- 7 At a store (not pharmacy)
- 8 Don't know

310. Do (Did) you pay for this method?

- 1 Yes
- 2 No

311. Would you prefer to use a different method of family planning from the one you are currently using?

- 1 Yes
- 2 No--->GO TO Q314

312. What method would you prefer to use?

- 1 Pills
- 2 IUD
- 3 Condoms
- 4 Foam/Jelly/Cream
- 5 Diaphragm
- 6 Female sterilization
- 7 Male sterilization
- 8 Fertile days method
- 9 Withdrawal
- 20 Other _____
- 77 Any method-- ----->GO TO Q314
- 88 Not sure ----->GO TO Q314

313. What is the most important reason that you do not use that method?

- 1 Doctor will not prescribe it
- 2 Cost
- 3 Not available
- 4 Too far away
- 5 Do not know how to obtain it
- 6 Husband objects to it
- 7 Religious reasons
- 20 Other_____
- 88 Don't know

314. IF USING WITHDRAWAL, RHYTHM, OR FERTILE DAYS METHOD(Q306=8, 9, OR 10)
CONTINUE WITH Q314;
USERS OF OTHER METHODS GO TO Q316.

You said that you are now using _____to avoid becoming pregnant, rather than a method you might obtain from a doctor, health facility, or pharmacy. Please tell me whether each of the following was very important, somewhat important, or not important at all in your decision to use this method:

	<u>Very Important</u>	<u>Somewhat Important</u>	<u>Not Important</u>	<u>Not Sure</u>
A. Difficult to get other methods	1	2	3	8
B. Cost of other methods	1	2	3	8
C. Little knowledge of other methods	1	2	3	8
D. Health/Side effects other methods	1	2	3	8
E. Husband/Partner preference	1	2	3	8
F. Religious beliefs	1	2	3	8

315. How effective at preventing pregnancies do you think __ (method currently used) is compared to modern methods, like the pill or the IUD? (READ CHOICES)

- 1 Current method more effective
- 2 About equally effective
- 3 Current method less effective
- 8 Don't know/Not sure

316. Do you have any problems or concerns with using _____?

- 1 Yes
- 2 No--->GO TO Q318

317. What is the most important problem or concern?

- 1 Side effects
- 2 Health concerns
- 3 Access/Availability
- 4 Cost
- 5 Sometimes forget to use
- 6 Sometimes difficult/inconvenient to use
- 7 Husband/partner disapproves
- 20 Other_____

318. Do you plan to have any (more) children?

- 1 Yes
- 2 No --- >GO TO Q321
- 8 Not sure -- >GO TO Q321

319. How many more do you plan to have?

__ __ children

66=As many as possible

77=Up to God/Fate, etc.

88=Not sure

320. When do you think you would like to become pregnant?

1 Within 1 year

2 In 1-2 years

3 In 3-5 years

4 In more than 5 years

7 When I get married

8 Not sure/Don't know

321. Think back to when you first started using a method to prevent pregnancy. What method was it?

1 Pills

2 IUD

3 Condoms

4 Foam/Jelly/Cream

5 Diaphragm

6 Female sterilization

7 Male sterilization

8 Fertile days method

9 Withdrawal

20 Other _____

88 Don't remember

322. How many living children did you have at that time?

__ __ children

8=8 or more

323. How old were you at that time?

__ __ years

328. CONTRACEPTIVE USE/PREGNANCY CALENDAR:

Starting at the beginning of 1988, please try to remember in which months you started and stopped use of contraceptive methods.

(INTERVIEWER: FILL IN ALL PREGNANIES AND BIRTHS FROM PREGNANCY HISTORY BEFORE COLLECTING CONTRACEPTIVE HISTORY).

COLUMN 1	DATE	1	2		DATE	1	2
<u>CONTRACE PT/PREGNANCY</u>	1988				1991		
0 No method	Jan 1				Jan 1		
1 Pills	Feb 2				Feb 2		
2 IUD	Mar 3				Mar 3		
3 Condoms	Apr 4				Apr 4		
4 Vaginal method	May 5				May 5		
5 Fertile days method	Jun 6				Jun 6		
6 Withdrawal	Jul 7				Jul 7		
7 Sterilization	Aug 8				Aug 8		
8 Other _____	Sep 9				Sep 9		
9 Don't remember 2	Oct 10				Oct 10		
0 Pregnant	Nov 11				Nov 11		
21 Live birth	Dec 12				Dec 12		
22 Stillbirth	1989				1992		
23 Miscarriage	Jan 1				Jan 1		
24 Extrauterine preg	Feb 2				Feb 2		
25 Induced abortion	Mar 3				Mar 3		
-----	Apr 4				Apr 4		
	May 5				May 5		
	Jun 6				Jun 6		
	Jul 7				Jul 7		
	Aug 8				Aug 8		
	Sep 9				Sep 9		
	Oct 10				Oct 10		
	Nov 11				Nov 11		
	Dec 12				Dec 12		
	1990				1993		
	Jan 1				Jan 1		
	Feb 2				Feb 2		
	Mar 3				Mar 3		
	Apr 4				Apr 4		
	May 5				May 5		
	Jun 6				Jun 6		
	Jul 7				Jul 7		
	Aug 8				Aug 8		
	Sep 9				Sep 9		
	Oct 10				Oct 10		
	Nov 11				Nov 11		
	Dec 12				Dec 12		

IF NOT USING A METHOD IN JANUARY 1988 SKIP Q329

329. You said you were using ____ in January of 1987. When did you start using that method?

Month ____ Year 19 ____

IF CURRENTLY USING A FAMILY PLANNING METHOD, GO TO Q338

330. Do you think you are able to get pregnant at the present time?

- 1 Yes--->GO TO Q332
- 2 No--- >GO TO Q331
- 3 Not sure--- >GO TO Q332
- 4 Currently pregnant-- >Q332

331. Why not?

- 1 Menopause/Too old to get pregnant \
- 2 Has had an operation for medical reasons \
- that makes pregnancy impossible \
- 3 Husband/partner has had a medical operation \
- 4 Has tried to get pregnant for \ - >GO TO Q345
- at least 2 years without success /
- (or has not gotten pregnant despite /
- 2 years not contracepting) /
- 5 Doctor says she or partner is infertile /
- 6 Not sexually active----- >GO TO Q332
- 7 Postpartum/Breastfeeding----- >GO TO Q332
- 8 Other (specify) _____>GO TO Q332

332. Do you plan to have any (more) children (after this pregnancy)?

- 1 Yes
- 2 No---- >GO TO Q336
- 3 Not sure---- >GO TO Q336

333. How many more children do you plan to have?

- ___ children
- 66=As many as possible
- 77=Up to God/Fate, etc.
- 88=Not sure

IF CURRENTLY PREGNANT SKIP TO Q338

334. Are you trying to become pregnant now?

- 1 Yes--->GO TO Q337
- 2 No

335. When do you think you would like to become pregnant?

- 1 Within 1 year
- 2 In 1-2 years
- 3 In 3-5 years
- 4 In more than 5 years
- 7 When I get married
- 8 Not sure

336. What is the most important reason you are not using a method to avoid pregnancy now?

- 0 Currently pregnant ----->G0 TO Q3 38
- 1 Not sexually active/No partner----->G0 TO Q338
- 2 Want to become pregnant
- 3 Only occasionally sexually active
- 4 Breastfeeding/Postpartum
- 5 Fear of side effects/health effects
- 6 Previously had side effects/health problems
- 7 Husband/Partner objects
- 8 Religious reasons
- 9 Doctor won't prescribe method
- 10 Desired method not available/difficult to get
- 11 Too expensive
- 12 Don't know where to get method
- 13 Methods difficult to use
- 14 Prefer abortion
- 15 Haven't bothered, but would like to use method
- 16 Too old
- 17 Difficulty getting pregnant
- 20 Other (specify)_____
- 88 Don't know

337. Does your husband/partner think that you should be using a method to prevent you from becoming pregnant?

- 1 Yes
- 2 No
- 3 No husband/partner
- 8 Don't know

338. PLANS TO HAVE MORE CHILDREN

- 1 YES (Q318=1 OR Q332=1) ----->CONTINUE WITH Q341
- 2 NO (Q318=2 OR Q332=2) ----->GO TO Q339
- 3 NOT SURE (Q318 = 8 OR Q332 = 3)- -->CONTINUE WITH Q341

339. Are you interested in having an operation to prevent you from having any more children (after this pregnancy)?

- 1 Yes --->GO TO Q341
- 2 No
- 3 Not sure --->GO TO Q341
- 4 STERILIZED (Q306=6 OR 7) ----->GO TO Q341

340. What is the most important reason you wouldn't be/are not interested?

- 1 Health risks
- 2 Fear of operation
- 3 Husband would object
- 4 Religious reasons
- 5 Not culturally/socially acceptable
- 6 Cost/inconvenience of an operation
- 7 Might want another child
- 8 Don't know enough about sterilization
- 9 Haven't thought about it
- 20 Other_____
- 88 Don't know

341. In the past five years have you ever unsuccessfully tried to get birth control pills?

- 1 Yes
- 2 No--->GO TO Q343

342. What **was** the reason you were unable to get them?

- 1 Physician refused because of health/side effects
- 2 Physician refused because of age
- 3 Physician refused because of marital status
- 4 Physician refused, other _____
- 5 Pills unavailable/out of stock _____
- 6 Cost _____
- 7 Other (specify) _____
- 8 Don't remember _____

343. In the past five years have you ever unsuccessfully tried to get an IUD?

- 1 Yes
- 2 No---->GO TO Q345

344. What was the reason you were unable to get it?

- 1 Physician refused because of health/side effects
- 2 Physician refused because of age
- 3 Physician refused because of marital status
- 4 Physician refused, other _____
- 5 IUDs unavailable/out of stock _____
- 6 Cost _____
- 7 Other (specify) _____
- 8 Don't remember _____

IF NEVER MARRIED, GO TO NEXT SECTION (Q400)

345. Have you (or your husband/partner) ever been to a doctor or clinic to talk about ways to help you become pregnant or to prevent a miscarriage?

- 1 Yes
- 2 No--->GO TO NEXT SECTION (Q400)

346. What kinds of medical treatment or advice did you or your husband/partner receive to help you become pregnant or prevent miscarriages? (READ ALTERNATIVES)

	<u>YES</u>	<u>NO</u>	<u>PR</u>
A. Drugs to make you ovulate	1	2	8
B. Surgery or treatment for blocked tubes	1	2	8
C. Infertility tests on you	1	2	8
D. Advice on how to time intercourse	1	2	8
E. Advice to start or stop contraception	1	2	8
F. Tests on husband/partner	1	2	8
G. In vitro fertilization	1	2	8
H. Artificial insemination	1	2	8
I. Bed rest	1	2	8
J. Other (specify)_____	1	2	8

347. When was the last time you or your husband/partner visited a doctor/clinic for this treatment or advice?

Month — —
 Year 19 — —

MATERNAL-CHILD HEALTH

**THE FOLLOWING QUESTIONS DEAL WITH THE MOST RECENT PREGNANCY THAT LED TO A LIVE
BIRTH SINCE 1/88
IF NO LIVE BIRTHS SINCE 1/88, GO TO Q500**

400. Which of the following best describes your reaction to your most recent pregnancy? (READ CHOICES)
- 1 Immediately happy about it
 - 2 Accepted it easily and quickly became happy about it
 - 3 Accepted it eventually
 - 4 Was never able to accept it
 - 5 Other (Specify)_____
401. Which of the following best describes the father's reaction to that pregnancy? (READ CHOICES)
- 1 Immediately happy about it
 - 2 Accepted it easily and quickly became happy about it
 - 3 Accepted it eventually
 - 4 Was never able to accept it/Refused to become involved
 - 5 Did not know about pregnancy
 - 8 Don't know
402. Did your job during that pregnancy involve much physical work, a moderate amount, or only a little?
- 1 Much physical work
 - 2 A moderate amount
 - 3 Little or no physical work
 - 4 Did not work----- >GO TO Q405
403. Did your job during that pregnancy involve much standing, a moderate amount of standing, or little standing?
- 1 Much standing
 - 2 Moderate standing
 - 3 Little standing
404. Would you say that job generally involved much, a moderate amount, or only a small amount of mental stress?
- 1 Much stress
 - 2 Moderate stress
 - 3 Little stress
405. Did you receive any prenatal care from a doctor, nurse, or midwife for that pregnancy?
- 1 Yes
 - 2 No--->GO TO Q418
406. Who provided the most care?
- 1 Physician
 - 2 Midwife/Nurse
 - 3 Physician and midwife equally
407. How many months pregnant were you when you made your first prenatal visit?
____ months

408. How many prenatal visits did you make during that pregnancy?
- ___ visits
 66=As many as doctor/midwife/nurse said to have
 77=Don't remember, but was definitely at least 10
 88=Don't remember
409. Where did you receive your prenatal care? (READ ALL CHOICES)
- 1 District clinic
 2 District clinic and hospital
 3 Hospital only
 4 Other (specify)_____
410. How many minutes did it take you, on average, to reach the place where you received the most prenatal care?
- ___ minutes
 000=At home/At factory -- >GO TO Q413
 888=Don't remember
411. How did you usually get to that place?
- 1 Walk
 2 Bus/Metro
 3 Private vehicle
 4 Other (specify)_____
412. On average, how long did you have to wait to be seen?
- ___ minutes
 888=Don't remember
413. During that pregnancy did you have an ultrasound?
- 1 Yes
 2 No----- >GO TO Q415
 8 Don't know/Don't remember ---- >GO TO Q415
414. Did anyone (e.g., midwife, physician) explain to you or make sure you understood the purpose of this procedure?
- 1 Explained well
 2 Explained only a little
 3 Did not explain
 8 Don't remember
415. During that pregnancy did you have an amniocentesis?
- 1 Yes
 2 No ----- >GO TO Q418
 8 Don't know - >GO TO Q418
416. During what month of pregnancy was it done?
- ___ month
 88=Don't know

417. Why was it done? (READ CHOICES)

- 1 To test for possible birth defect
- 2 For RH immunization
- 3 Metabolic disorder
- 4 Other _____
- 5 Don't know/Was not explained
- 8 Don't remember

418. During your last pregnancy did a doctor (or midwife?) ever tell you to remain in bed/stay off your feet for one or more weeks because of some problem related to your pregnancy?

- 1 Yes
- 2 No--->GO TO Q420

419. Altogether, how long did you stay in bed because of this problem during that pregnancy?

_____ weeks 01=1
week or less

420. During your last pregnancy were you ever hospitalized because of any problem related to your pregnancy?

- 1 Yes
- 2 No--->GO TO Q423

421. Altogether, how many days were you hospitalized because of this problem during that pregnancy?

___ ___ days

422. Were any of the following conditions the reason you were hospitalized?
(READ EACH CONDITION)

	YES	NO	DK/DR
A. Threatened miscarriage; bleeding in first half of pregnancy	1	2	8
B. Bleeding during second half of pregnancy	1	2	8
C. High blood pressure (hypertension) before pregnancy	1	2	8
D. High blood pressure (hypertension) related to pregnancy	1	2	8
E. Early or false labor	1	2	8
F. Swollen ankles, water retention, edema	1	2	8
G. Improper position of fetus	1	2	8
H. Previous cesarean section	1	2	8
I. Other (specify) _____	1	2	8

423. Did you smoke cigarettes just before you became pregnant?

- 1 Yes
- 2 No--->GO TO Q426

424. Did you smoke cigarettes during that pregnancy?
- 1 Yes
 - 2 No -->GO TO Q426
425. About how many cigarettes did you smoke each day, on average, during that pregnancy?
(ONE PACK=20 CIGARETTES)
- — cigarettes
- 77=Less than 1 per day
80=80 or more
88=Don't remember
426. How many times per week did you drink alcoholic beverages during that pregnancy?
- 1 At least 4 times per week
 - 2 1-3 times per week
 - 3 Less than once per week
 - 4 Never
 - 9 No response
427. How much did your last baby weigh when he/she was born?
- — — — grams -->GO TO Q429
- 8888=Don't know/Don't remember -->GO TO Q428
428. Do you remember if he/she weighed less than 2500 grams or was considered to be low birth weight?
- 1 Yes (<2500 grams/low birth weight)
 - 2 No
 - 8 Don't remember/Don't know
429. Was your last delivery normal or cesarean?
- 1 Normal
 - 2 Cesarean
430. Would you rate the place in which you had your last delivery as good, fair, or poor in the following areas?
- | | <u>GOOD</u> | <u>FAIR</u> | <u>POOR</u> | <u>DK/DR</u> |
|---------------------------------------|-------------|-------------|-------------|--------------|
| A. Distance from home | 1 | 2 | 3 | 8 |
| B. Physical facilities | 1 | 2 | 3 | 8 |
| C. Crowdedness | 1 | 2 | 3 | 8 |
| D. Helpfulness/Attentiveness of staff | 1 | 2 | 3 | 8 |
| E. Competence of staff | 1 | 2 | 3 | 8 |
431. Would you say that your delivery was (READ CHOICES):
- 1 Shorter than you expected
 - 2 About as long as you expected
 - 3 Longer than you expected
 - 4 Planned Cesarean section --->GO TO Q433
432. Would you say that your labor was (READ CHOICES):
- 1 Less painful than you expected
 - 2 About as painful as you expected
 - 3 More painful than you expected

433. Do you feel that you received enough information regarding your pregnancy from those providing your care?
- 1 Yes
 - 2 No
434. Do you feel that you received enough information regarding what to expect during your labor and delivery from those providing your care?
- 1 Yes
 - 2 No
435. Who would you have wanted to be present at your delivery?(READ CHOICES)
- 1 Your husband/partner
 - 2 Your mother
 - 3 Other person (specify) _____
 - 4 Only professional staff
436. If you were to have another child, would you be interested in delivering your baby at home, rather than in a hospital?
- 1 Yes, definitely
 - 2 Possibly
 - 3 No, definitely
 - 4 Not sure
437. If you were to have another child, do you think you would prefer to have it by cesarean section or normally?
- 1 Cesarean
 - 2 Normal
 - 3 Not sure

438. What was your biggest problem the first week after delivery?

- 1 Health/Care of the child
- 2 Breastfeeding
- 3 Her own health
- 4 Behavior of professional staff
- 5 Hospital environment
- 6 Visiting/Contact with family
- 7 Other (specify)_____
- 77 No important problems
- 88 Don't remember

439. What was your biggest problem when you returned home after delivery?

- 1 How to care for child
- 2 Health of child
- 3 Her own health
- 4 Lack of professional care
- 5 Home environment (e.g., living with parents, crowded home)
- 6 Family circumstances (e.g., no husband, husband not helpful)
- 7 Economic situation
- 8 Other (specify)_____
- 77 No important problems
- 88 Don't remember

440. Did you know whom to ask for advice or assistance with problems after you returned home?

- 1 Yes
- 2 No
- 3 Only for certain problems

441. Did you attend any prenatal counselling?

- 1 Yes
- 2 No----->GO TO Q443

442. How many sessions did you attend?

- 1 < 4
- 2 4+
- 8 Don't remember

443. Did you receive any analgesia during labor?

- 1 Yes
- 2 No ----->GO TO Q500
- 3 Don't remember--->GO TO Q500
- 4 Cesarean section-->GO TO Q500

444. How effective was the analgesia you received?

- 1 Very effective
- 2 Somewhat effective
- 3 Not effective
- 8 Don't remember

YOUNG ADULT MODULE

500. AGE OF RESPONDENT

1 15-24 -- >CONTINUE WITH Q501

2 25-44 -- >GO TO Q600

501. In what month and year did you first have sexual relations, if ever?

Month

Year 19 --

22=Never had sexual relations -->GO TO Q516

98=Don't remember

99=No response--->GO TO Q509

502. How old were you at that time?

-- years

88=Don't remember

503. At that time what was your relationship to your first partner?

1 Husband

2 Engaged to be married

3 Boyfriend

4 Friend

5 Rape ---- >GO TO Q509

6 Incest -- >GO TO Q509

7 Other _____

9 No response

504. How long were you and your first partner dating when you first had sexual relations?

____ months

00=First time we met

61=More than five years

77=Other _____

88=Don't remember

99=no response

505. Did you or your partner use a method or do anything to prevent pregnancy at that time?

1 Yes

2 No--- >GO TO Q508

8 Don't remember/Don't know --- >GO TO Q508

506. What method?

1 Pills

2 IUD

3 Condoms

4 Foam/Jelly/Cream

5 Diaphragm

6 Fertile days method

7 Withdrawal

8 Other _____

507. Who took the initiative to use this method?

- 1 You
- 2 Partner
- 3 Both
- 8 Don't remember

GO TO Q509

508. Why didn't you or your partner use a contraceptive method?

- 1 Did not expect to have sex
- 2 Did not know any methods
- 3 Wanted to get pregnant
- 4 Health concerns about contraception
- 5 Wanted to use, but didn't have any
- 6 Didn't think she could get pregnant
- 7 Other _____
- 8 Don't know/Don't remember

509. RESPONDENT EVER PREGNANT

- 1 Yes (Q200=1 OR Q201=1)
- 2 No (Q200=2 OR 3) -- >GO TO Q516

510. How old were you when you became pregnant (for the first time)?

_ Years

511. With whom were you living when you first became pregnant?

- 1 Was married--- >GO TO Q516
- 2 Mother and father
- 3 Mother or father only
- 4 Grandparents only
- 5 Other relatives
- 6 Friends
- 7 Alone
- 8 Don't remember

512. When you became pregnant (the first time), what was your relationship to the father?

- 1 Engaged to be married
- 2 Boy friend
- 3 Friend
- 4 Casual acquaintance/Just met
- 5 Was raped/Incest ----->GO TO Q516
- 6 Other _____
- 9 No response

513. What was the attitude of your partner when he learned of your pregnancy?

- 1 Both wanted to get married
- 2 Only he wanted to get married
- 3 Wanted you to get an abortion
- 4 Accepted pregnancy, but did not want to get married
- 5 He did not know of pregnancy ----- >GO TO Q515
- 7 Other _____
- 8 Don't know

514. Did you marry or go live with him?

- 1 Yes
2 No

515. What was the attitude of your family when they learned of the pregnancy?

- 1 Wanted you to get married
2 Insisted that you get married
3 Wanted you to get an abortion
4 Accepted the pregnancy without marriage
5 They didn't interfere
6 They did not know about pregnancy
7 Other _____
8 Don't know

516. Did your mother or father ever talk to you about the use of pregnancy prevention methods?

- ```

1 Yes
2 No----->G0 TO Q600

```

517. How old were you when they told you about it the first time?

— — years  
88=Don't remember

WOMEN'S HEALTH

600. How often do you go for regular (not pregnancy related) gynecologic exams?

- 1 At least once per year ----- >GO TO Q602
- 2 Every 1-2 years
- 3 Every 3-5 years
- 4 Less than every 5 years
- 5 Never

601. Health care professionals recommend that women receive such an exam every year. What is the main reason you go less often than that?

- 1 Doesn't feel it is necessary to go that often
- 2 She is healthy/doesn't have gynecologic problem
- 3 Doesn't have the time
- 4 She forgets about it
- 5 Doesn't like gyn. exams
- 6 Hard to get appointments
- 7 Doesn't like facilities
- 8 Doesn't like staff
- 9 Waiting time is too long
- 10 Other (specify) \_\_\_\_\_
- 88 Don't know

602. Have you heard about breast self-examinations?

- 1 Yes
- 2 No--->GO TO Q604

603. Do you ever do breast self-examinations?

- 1 Yes
- 2 No

604. Have you smoked at least 100 cigarettes during your life?

- 1 Yes
- 2 No--->GO TO Q607

605. Do you currently smoke?

- 1 Yes
- 2 No--- >GO TO Q607

606. How many cigarettes do you smoke per day, on average?

(1 PACK=20 CIGARETTES)

\_\_\_ cigarettes  
 77=<1 cigarette per day  
 80=80+

607. How many times per week do you usually drink alcoholic beverages?

- 1 At least 4 times per week
- 2 1-3 times per week
- 3 Less than once per week
- 4 Never----->GO TO Q611
- 9 No response----->GO TO Q611

608. How much beer do you drink?

\_\_ \_\_ liters per week

609. How much wine do you drink?

\_\_ \_\_ liters per month

610. How much liquor do you drink?

\_\_ \_\_ deciliters per month

611. How tall are you?

\_\_ \_\_ \_\_ centimeters

612. How much do you weigh?

\_\_ \_\_ \_\_ kilograms

## REPRODUCTIVE HEALTH KNOWLEDGE/ATTITUDES

700. What do you think is the ideal number of children for a young family in the Czech Republic?

- children
- 6 6 or more
- 7 As many as possible
- 8 Don't know

701. When is it most likely for a woman to become pregnant?  
(SHOW CARD A)

- 1 A week before menstruation starts
- 2 During menstruation
- 3 A week after menstruation starts
- 4 Halfway between her periods
- 5 It doesn't matter, all times alike
- 7 Other (specify) \_\_\_\_\_
- 8 Don't know

702. Do you think that in any situation a woman should always have the right to decide about her own pregnancy, including whether to have an abortion?

- 1 Yes--->GO TO Q704
- 2 No

703. Under which of the following conditions is it all right for a woman to have an abortion?

|                                             | <u>YES</u> | <u>NO</u> | <u>DEPENDS</u> | <u>DK</u> |
|---------------------------------------------|------------|-----------|----------------|-----------|
| A. Her life is in danger from the pregnancy | 1          | 2         | 3              | 4         |
| B. The fetus has a physical deformity       | 1          | 2         | 3              | 4         |
| C. The pregnancy has resulted from rape     | 1          | 2         | 3              | 4         |
| D. Her health may be hurt by the pregnancy  | 1          | 2         | 3              | 4         |
| E. She is unmarried                         | 1          | 2         | 3              | 4         |
| F. The couple cannot afford to have a child | 1          | 2         | 3              | 4         |

704. If a woman takes the pill correctly, how sure can she be that she will not become pregnant? (READ ANSWERS 1-4)

- 1 Completely sure
- 2 Almost sure
- 3 Fairly sure
- 4 Not sure at all
- 8 Don't know

705. How safe for a woman's health is the pill? (READ ANSWERS 1-3)

- 1 Completely safe
- 2 Fairly safe
- 3 Unsafe
- 8 Don't know

706. Please indicate whether you agree or disagree with the following statements about the pill.

|                                                                    | <u>AGREE</u> | <u>DISAGREE</u> | <u>DK</u> |
|--------------------------------------------------------------------|--------------|-----------------|-----------|
| A. The pill makes you gain weight                                  | 1            | 2               | 8         |
| B. The pill makes periods more regular                             | 1            | 2               | 8         |
| C. The pill makes you nervous                                      | 1            | 2               | 8         |
| D. Taking the pill for too long can<br>cause infertility           | 1            | 2               | 8         |
| E. Women who take the pill have a higher<br>risk of getting cancer | 1            | 2               | 8         |
| F. The pill is bad for blood circulation                           | 1            | 2               | 8         |

707. Now I will read some other statements. Please tell me whether you agree or disagree with each.

|                                                                                | <u>AGREE</u> | <u>DISAGREE</u> | <u>DK</u> |
|--------------------------------------------------------------------------------|--------------|-----------------|-----------|
| A. Care of children is women's work                                            | 1            | 2               | 8         |
| B. A woman should be a virgin when she marries                                 | 1            | 2               | 8         |
| C. A woman can become pregnant the first time<br>that she has intercourse      | 1            | 2               | 8         |
| D. A condom should not be used more than one time                              | 1            | 2               | 8         |
| E. It is all right for a woman not to have<br>children if she does not want to | 1            | 2               | 8         |

## KNOWLEDGE OF AIDS

800. Have you ever heard of the disease called AIDS or HIV infection?
- 1 Yes
  - 2 No -- >GO TO Q900
801. Do you think a person can be infected with the AIDS virus and not have any signs of the disease?
- 1 Yes
  - 2 No
  - 8 Don't know
802. Do you believe a person can become infected with AIDS in the following ways? (READ EACH LINE)
- |                                                 | YES | NO | DK |
|-------------------------------------------------|-----|----|----|
| A. Receiving a blood transfusion                | 1   | 2  | 8  |
| B. Using public bathrooms                       | 1   | 2  | 8  |
| C. Kissing on the mouth                         | 1   | 2  | 8  |
| D. Having heterosexual relations                | 1   | 2  | 8  |
| E. Men having homosexual relations              | 1   | 2  | 8  |
| F. Shaking hands                                | 1   | 2  | 8  |
| G. Using a needle already used by someone else  | 1   | 2  | 8  |
| H. Mosquito bite                                | 1   | 2  | 8  |
| I. Donating blood                               | 1   | 2  | 8  |
| J. Using domestic objects of a person with AIDS | 1   | 2  | 8  |
| K. Going to a physician or dentist              | 1   | 2  | 8  |
803. What kind of protection do you think condoms provide against AIDS infection? (READ CHOICES)
- 1 Excellent
  - 2 Good
  - 3 Fair
  - 4 Poor
  - 8 Don't know
804. If an anonymous test for AIDS infection was easily and cheaply available would you have it done?
- 1 Yes
  - 2 No
  - 8 Don't know
805. Do you think that you have any risk of getting AIDS?
- 1 Yes
  - 2 No ----- >GO TO Q900
  - 8 Don't know ---- >GO TO Q900
806. Would you say that you have a low risk or a high risk?
- 1 Low risk
  - 2 High risk
  - 8 Don't know

## SOCIOECONOMIC CHARACTERISTICS

900. What is your ethnic background?

- 1 Czech
- 2 Slovak
- 3 Moravian
- 4 Hungarian
- 5 Romany (Gypsy)
- 6 Polish/Silesian
- 7 German
- 8 Other (specify) \_\_\_\_\_
- 9 Refused/Not stated

901. What is your religion?

- 0 No religion -- >GO TO Q903
- 1 Roman Catholic
- 2 Czech Evangelical
- 3 Czechoslovak Hus Church
- 4 Slovakian Evangelical
- 5 Pravoslavne
- 6 Seventh Day Adventist
- 7 Greek Catholic
- 8 Other (specify) \_\_\_\_\_
- 9 Refused/Not stated -- >GO TO Q903

902. About how often do you usually attend religious services?  
(READ ANSWERS 1-5)

- 1 At least once a week
- 2 At least once a month, but less than once a week
- 3 Less than once a month
- 4 Only on holidays
- 5 Never/Almost never

903. Do you currently work outside of the home?

- 1 Yes
- 2 Yes, but on maternity/pregnancy leave -- >GO TO Q906
- 3 No--->GO TO Q906

904. Do you currently work one job or more than one job?

- 1 One job
- 2 More than one job

905. How many hours per week do you usually work, in total?

\_\_ \_\_ hours  
80=80+ hours

906. Please tell me whether this household or any member of it has the following items:

|                         | Yes | No |
|-------------------------|-----|----|
| A. Bathroom/Shower      | 1   | 2  |
| B. Local heating        | 1   | 2  |
| C. Color television     | 1   | 2  |
| D. VCR                  | 1   | 2  |
| E. Automobile           | 1   | 2  |
| F. Auto Washing machine | 1   | 2  |
| G. Telephone            | 1   | 2  |
| H. Personal computer    | 1   | 2  |

907. Do you own or have access to a weekend house (chaloupa)?

- 1 Own
- 2 Have access to
- 3 No

908. What was the average monthly gross income of all members of this household during the 1992? (SHOW CARD B)

- 1 LE 3000 KCr
- 2 3001-7000 KCr
- 3 7001-10,000 KCR
- 4 10,001-15,000 KCR
- 5 15,001-20,000 KCR
- 6 GT 20,000 KCR
- 8 Don't know
- 9 Refuse

909. Does more than one family live in this flat/house?

- 1 Yes
- 2 No

910. Which of these best describes this house/flat?  
(READ CHOICES)

- 1 Own home
- 2 Cooperative apartment/flat
- 3 Rental, from private owner
- 4 Rental, state owned

**IF NOT CURRENTLY MARRIED/NOT IN UNION END INTERVIEW**

911. How many jobs does your husband currently have?

- 0
- 1
- 2 2 or more

**END OF INTERVIEW**

**TIME INTERVIEW ENDED**      \_\_ \_\_ : \_\_ \_\_

INTERVIEWED BY: \_\_\_\_\_

QUESTIONNAIRE REVIEWED BY: \_\_\_\_\_

DATA ENTRY OPERATOR NUMBER: \_\_\_\_